


REVIEW

Open Access

# COVID-19 in Africa: care and protection for frontline healthcare workers



Matthew F. Chersich<sup>1\*</sup> , Glenda Gray<sup>2</sup>, Lee Fairlie<sup>1</sup>, Quentin Eichbaum<sup>3,4</sup>, Susannah Mayhew<sup>5</sup>, Brian Allwood<sup>6</sup>, Rene English<sup>7</sup>, Fiona Scorgie<sup>1</sup>, Stanley Luchters<sup>8,9,10,11</sup>, Greg Simpson<sup>12</sup>, Marjan Mosalman Haghighi<sup>13</sup>, Minh Duc Pham<sup>14,15</sup> and Helen Rees<sup>1</sup>

## Abstract

Medical staff caring for COVID-19 patients face mental stress, physical exhaustion, separation from families, stigma, and the pain of losing patients and colleagues. Many of them have acquired SARS-CoV-2 and some have died. In Africa, where the pandemic is escalating, there are major gaps in response capacity, especially in human resources and protective equipment. We examine these challenges and propose interventions to protect healthcare workers on the continent, drawing on articles identified on Medline (Pubmed) in a search on 24 March 2020. Global jostling means that supplies of personal protective equipment are limited in Africa. Even low-cost interventions such as facemasks for patients with a cough and water supplies for handwashing may be challenging, as is 'physical distancing' in overcrowded primary health care clinics. Without adequate protection, COVID-19 mortality may be high among healthcare workers and their family in Africa given limited critical care beds and difficulties in transporting ill healthcare workers from rural to urban care centres. Much can be done to protect healthcare workers, however. The continent has learnt invaluable lessons from Ebola and HIV control. HIV counselors and community healthcare workers are key resources, and could promote social distancing and related interventions, dispel myths, support healthcare workers, perform symptom screening and trace contacts. Staff motivation and retention may be enhanced through carefully managed risk 'allowances' or compensation. International support with personnel and protective equipment, especially from China, could turn the pandemic's trajectory in Africa around. Telemedicine holds promise as it rationalises human resources and reduces patient contact and thus infection risks. Importantly, healthcare workers, using their authoritative voice, can promote effective COVID-19 policies and prioritization of their safety. Prioritizing healthcare workers for SARS-CoV-2 testing, hospital beds and targeted research, as well as ensuring that public figures and the population acknowledge the commitment of healthcare workers may help to maintain morale. Clearly there are multiple ways that international support and national commitment could help safeguard healthcare workers in Africa, essential for limiting the pandemic's potentially devastating health, socio-economic and security impacts on the continent.

**Keywords:** COVID-19, SARS-Cov-2, Africa, Human resources for health, Healthcare workers, Infection control, mental health

\* Correspondence: [mchersich@wrhi.ac.za](mailto:mchersich@wrhi.ac.za)

<sup>1</sup>Wits Reproductive Health and HIV Institute, Faculty of Health Sciences, University of the Witwatersrand, Johannesburg, South Africa  
Full list of author information is available at the end of the article



© The Author(s). 2020 **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

## Methods for review of literature

### Inclusion and exclusion criteria

Studies were included if they presented data or commentaries on the infection risks and mental wellbeing impacts that healthcare workers face during the COVID-19 pandemic. Studies could be in any setting. We also included articles that covered the COVID-19 pandemic in Africa in general (these papers did not have to include infection risks and mental health). We excluded articles on infection control or mental health of COVID-19 if they covered topics that were not relevant to Africa.

### Search strategy

The literature was identified in a search on Medline(-Pubmed) up to 24 March and identified using the following search strategy: ((((((coronavirus) AND ("2020"[Date - Create] : "3000"[Date - Create]))) OR SARS-CoV-2) OR 2019-nCoV) OR COVID).

## Background

Sustaining safe and quality care in the SARS-CoV-2 pandemic hinges on the health and mental wellbeing of frontline healthcare workers. Medical staff face exhaustion, difficult triage decisions, separation from families, stigma and the pain of losing patients and colleagues, in addition to their own risks of infection. In Italy as of 3 April 2020, around 10,000 healthcare workers have been infected and 74 have died, and many others have died in countries across the globe [1, 2]. While the pandemic in Africa is several weeks behind Europe and Asia, the number of cases in Africa is escalating fast [3–5]. Incidence varies considerably between countries in Africa possibly reflecting variations in volumes of air travel and differences in coverage of SARS-CoV-2 testing [6]. While many countries in Africa are stepping up their preparedness for COVID-19 [6], assessments by WHO point to substantial limitations in response capacity [7]. In particular, there are major shortages of human resources, critical care beds and laboratory capacity. For example, in 2018 the numbers of nurses or midwives to 10,000 population was about 6.0 in Côte d'Ivoire and Mozambique, around 11 in the Democratic Republic of the Congo and Kenya [8]. Corresponding figures for the United Kingdom were 81.7 and 132.4 in Germany. Many countries in Africa have fewer than 30 critical care beds to cover the entire population [9]. Notwithstanding these gaps, COVID-19 control efforts can draw on valuable lessons learnt during the recent Ebola outbreaks and from HIV prevention and treatment successes on the continent. In a similar way, countries such as Singapore and South Korea had learnt valuable lessons from their experiences of the SARS and MERS outbreaks, which they then applied to control of SARS-CoV-2.

In this review we describe the infection risks and mental health challenges that healthcare workers face in the COVID-19 pandemic and propose interventions to counter these in Africa. We highlight lessons from previous disease-control efforts on the continent and draw on experiences with SARS-CoV-2 in other parts of the world. We searched Medline (Pubmed) on 24 March (see Additional File Fig. 1 for the search strategy and PRISMA flow chart) and located 1464 articles, of which 88 were on healthcare workers, and 32 considered relevant to this review. Articles on healthcare workers and Ebola or HIV were identified in targeted searches.

### Healthcare workers' vulnerability to COVID-19: critical interventions to minimise risk

In all settings, patients who have unusual symptoms of COVID-19 or very mild general flu-like symptoms pose considerable risk to healthcare workers who may not have a high level of clinical suspicion in these patients and adopt adequate protective measures. Additionally, in the tropics, patients may present with febrile conditions related to vector-borne and other infections, but have SARS-CoV-2 coinfection. In Thailand, for example, a patient was admitted with dengue fever and also had SARS-CoV-2 and the healthcare worker responsible for care became infected [10].

Healthcare workers – and, indeed, patients admitted to a health facility for other reasons – are particularly vulnerable to infection from 'super-spreading events'. In one instance in Wuhan, China a large cluster of infections occurred in healthcare workers and patients [11]. These infections were traced to a patient with abdominal symptoms who had been admitted in the surgical department [11]. Tracing and controlling super-spreading events within hospitals requires resources, which may not be available in many Africa settings.

Risks of SARS-CoV-2 infection may be higher among professionals who work in close physical proximity to patients, such as ophthalmologists and dentists [12–15]. Additionally, some procedures such as non-invasive ventilation, high-flow nasal cannula and bag-mask ventilation may generate higher aerosol volumes, but be key treatment modalities in settings without mechanical ventilators [16].

Risks of healthcare worker infection can be mitigated with adequate precautions within health facilities [17–20]. Primarily, this involves the use of personal protective equipment (PPE) including a gown, gloves, face mask, and a face shield or goggles. Careful donning and doffing of this equipment remains a key defense, but requires considerable training and supervision. Risks for infection may also be highest at the beginning of the outbreak when healthcare workers may not yet be familiar with PPE use. There are major PPE shortages in

high-income countries and it is likely that limited supplies will be allocated to less resourced countries [21]. These scarce PPE resources need to be appropriately used and distributed equitably across the globe – yet hoarding, misuse, intense competition between and within countries, price gouging, and export blocks are threatening to become the norm [21, 22]. Without international support, any reserves of PPE in hospitals are likely to be rapidly depleted in African countries and new supplies will be very difficult to secure [23]. Importantly, digitalized or telemedicine services could potentially reduce patient contact and thus risks for infection, and allow for national or international experts to give advice from a distance and support to less-experienced doctors. While there may be considerable costs in setting up such systems, these may be outweighed by savings in PPE, staff resources and improved patient outcomes. These initiatives may, however, be challenged by infrastructural constraints such as an unstable power supply or limited internet connectivity, and a lack of interoperability between digital systems.

Risks of infection in healthcare workers appears tied to duration of shifts and hand hygiene, among other factors [24]. Water supplies for handwashing, however, may be limited or unavailable in some parts of Africa. Guidance is needed on the range of low-cost cleaning agents that can be used in lieu of commercially-manufactured sanitisers for cleaning and disinfection of work surfaces and items such as pens, stethoscopes and mobile phones [25].

A range of simple low-cost interventions can reduce the likelihood of infection transmission to healthcare workers in hospitals, including facemasks for patients who have respiratory symptoms, tissues for patients, promotion of cough etiquette and hand washing, and maintenance of at least two metres distance from others. ‘Social [physical] distancing’ between patients and administrative, cleaning staff and healthcare workers may, however, be especially challenging in busy overcrowded primary health care clinics on the continent.

Many governments in Africa will require support from international donors to protect frontline healthcare workers from the outset of the pandemic. The government of China has provided personnel and equipment to support COVID-19 services in many parts of the world [26]. Support from China, building on already well-established links with countries across the continent, could play a major role in shaping the trajectory of the pandemic in Africa. Using international staff to provide care, however, presents challenges. In the Ebola outbreak in West Africa, for example, misunderstandings and language barriers were common with international medical teams, while the needs of local lower-level responders were sometimes neglected. The outsiders were viewed

with distrust or suspicion, as, for example ‘patients were seen to be taken away by hazmat-suited strangers to die in unknown locations’ [27].

Mortality rates among healthcare workers who become infected may be especially high in many parts of Africa given the limited number of critical care beds [28]. Moreover, the large geographical distances pose tremendous practical difficulties in transferring ill healthcare workers from rural areas to secondary- or tertiary-level facilities in urban centres. Additionally, many healthcare workers in Africa themselves fall into the category of ‘high-risk’ groups for COVID-19, given their high rates of certain non-communicable diseases, tuberculosis, and HIV, although evidence on the impacts of the latter on COVID-19 is not yet available [29–31]. Importantly, the benefits of continuing to isolate healthcare workers with SARS-CoV-2 infection or high-risk exposures after a predefined time assuming they do not have symptoms, needs to be weighed against the consequent service disruptions in settings with severe human resource constraints [32].

Overall, much more thought is needed about the optimum use of healthcare workers in the COVID-19 response in rural parts of the continent. During the Ebola outbreak, mobile testing units moved between villages, detecting clusters of infection, quarantining those areas and building temporary treatment centres where required. Such efforts helped to target human and other resources to where it was most needed, while allowing economic activities to continue in unaffected areas. Some combination of antigen, antibody testing and symptom screening may be needed to identify affected communities who are then prioritised for COVID-19 services. Testing and later vaccinating of healthcare workers was a key strategy in the Ebola epidemic, but testing constraints and lack of a vaccine hinders that approach with the current pandemic. Serological testing which demonstrates past infection may provide major relief for healthcare workers and potentially allow people who have immunity to return to work.

#### **Securing the mental wellbeing of healthcare workers**

Caring for patients with COVID-19 disease causes considerable mental stress, resulting in high levels of anxiety and post-traumatic stress disorders, especially among nurses [33]. These conditions have a major impact on healthcare workers, but also undermine their decision-making ability and quality of interaction with patients [34]. It is worth bearing in mind that the strain experienced at work is compounded by the very same disruptions and uncertainties felt by members of the general population at this time [35].

Psychological support is key, perhaps drawing on the large numbers of HIV counsellors or retired nurses in

many parts of the African continent. Formal structured interventions, however, may encounter obstacles, given the competing priorities of health staff [36]. Informal mechanisms might be more successful, where, for example, counsellors or retired nurses visit healthcare worker rest areas to listen to difficulties or to the stories recounted by staff about their work. Support among healthcare workers through social media, such as the Vula platform in South Africa, may also relieve stress and could be extended to other countries [36]. WhatsApp groups among healthcare workers could provide advice on clinical decision making, but also be used to circulate messaging on mental health support. Regular communication between healthcare workers in the Ebola response meant that new knowledge could be shared among providers, especially where there were gaps in evidence on the disease [27].

While healthcare workers may accept an increased risk of infection as part of their chosen profession, they may have considerable anxiety about spreading the virus to their children, families and friends, especially those who are elderly or have chronic medical conditions, and perhaps even their pets, despite the evidence for this being limited [37, 38]. Healthcare workers may wish to have alternative accommodation to avoid the risk of household transmission. Student residences or hotels that are currently empty could be re-purposed to serve as places where they can rest and temporarily isolate themselves from their family [36]. Healthcare workers may also prefer to have longer shifts for a week and stay at the hospital during that time, rather than having shorter shifts and returning home each night. Protective actions within the home may include separating living spaces and bathrooms where possible and having a routine when arriving at home after duty, such as taking off shoes, removing and washing clothing, and immediately showering [39]. Many of these precautions may not be possible, however, in overcrowded areas with limited infrastructure and water availability, for example.

In any biological disaster, fear and stigmatisation are heightened, and healthcare workers may be a target of the latter. Many healthcare workers in the recent Ebola and SARS epidemics experienced considerable stigmatization, loneliness and even loss of trust within their own communities [40, 41]. In Singapore during the SARS epidemic in the early 2000s, for example, one nurse in a lift was told that her presence in the lift was spreading the virus to others, and another was scolded by fellow passengers for making trains “dirty”. These factors are of critical importance and healthcare workers need to feel socially supported; this may affect their self-efficacy, sleep quality and anxiety levels [42]. During the SARS epidemic, over time, some healthcare workers became reluctant to continue working [43, 44]. Maintaining staff motivation may be

especially challenging where levels of trust in the health system and in the government are low, as may be the case in areas of Africa where governments are perceived to be corrupt [44]. A ‘risk allowance’ may be one strategy for motivating and retaining healthcare workers, although there might be jealousy as other staff will continue to care for non-COVID-19 patients and hence miss out on these benefits. Risk allowances were used during the recent Ebola epidemic in Western Africa where the benefits were apparent, but also the jealousies of those not receiving these allowances [45]. It is also important to acknowledge that healthcare workers are powerful agents for change and need to be involved in decision making and in shaping the outbreak response. Healthcare workers in Africa hold an authoritative voice in local communities and society more broadly. They can help ensure that local governing structures, national governments and the international community adopt effective control measures and treatment strategies, and prioritize their safety [46].

Initiatives to increase the number of healthcare workers available to meet the burden of care have included moving staff from other disciplines to medical wards, fast-tracking medical students to join the workforce, cancelling healthcare worker leave and drawing on retired healthcare workers [21, 47]. Healthcare workers can also be relieved from administrative tasks, freeing up their time for patient care [48]. Mobilizing existing HIV counselors and community healthcare workers may be a key strategy in the COVID-19 response in Africa. These cadres, who have long served as ‘first responders’ in Africa could be key information sources to dispel myths in the community, but also perform symptom screening and contact tracing. Community health volunteers and health committee members also provide important supports and linkages to the formal health system in many settings. During the Ebola epidemic, for example, these groups helped build isolation structures, performed tasks such as screening and contact tracing, and assisted healthcare workers to better understand the needs of the community. For their part, community volunteers’ main concerns included inadequate communication with the health system and concerns regarding compensation for their work [49]. Policymakers will, however, have to weigh up the benefits of ‘task shifting’ these community-based cadres against the risks of placing them in situations of heightened risk of acquiring the virus, and with little or no protective equipment.

## Conclusion

Healthcare workers in Africa face considerable challenges during the pandemic. We present 10 priority interventions to safeguard frontline healthcare workers in Africa (Table 1). The degree to which we protect their health and mental wellbeing will shape the pandemic in

**Table 1** 10 priority interventions for securing the health and mental wellbeing of frontline healthcare workers in the COVID-19 response in Africa

1. Secure supplies of PPE of good quality.
2. Capacity development on PPE use and infection control more generally.
3. Research examining protection and support interventions for healthcare workers in Africa, including feasible approaches to repeated SARS-CoV-2 testing.
4. Prioritize healthcare workers for SARS-CoV-2 testing, beds in ICU and other medical wards, drug and vaccine trials, and therapeutics when these become available.
5. Politicians and other public figures to visit healthcare workers, acknowledge their commitment and sacrifices, and address any negative perceptions towards providers [50].
6. Provide food and daily living supplies for healthcare workers, which save them time, but also demonstrate society's appreciation of their work [51].
7. Incorporate a range of healthcare worker cadres into the response, especially community healthcare workers and HIV counsellors.
8. Creative interventions to reduce risks of infection, such as 'Eagle-Eye Observers' who are dedicated full-time staff charged with observing and correcting infection control errors, [52] a task which lay people could potentially do as this is largely based on checklists.
9. Finances to cover the costs of funerals for healthcare workers.
10. 'Risk allowances' to compensate healthcare workers for the risks they take and to motivate staff to continue to work.

Africa, and thus its long-run impacts on social stability, economic growth, and security [53]. While there are clearly a range of priorities for the COVID-19 response on the continent, we strongly urge WHO, national governments, the private sector, and the general populace to pay concerted attention to healthcare worker safety and mental wellbeing.

### Supplementary information

Supplementary information accompanies this paper at <https://doi.org/10.1186/s12992-020-00574-3>.

#### Additional file 1.

#### Abbreviation

PPE: Personal Protective Equipment

#### Acknowledgements

Not applicable.

#### Authors' contributions

MFC conceptualized the article and wrote the first draft. All co-authors contributed to writing the drafts of the paper, provided critical review of each draft, and approved the final manuscript.

#### Funding

No funding was used for the review.

#### Availability of data and materials

Not applicable as it is a review. Data sharing is not applicable to this article as no datasets were generated or analysed during the current study.

#### Ethics approval and consent to participate

Not applicable as it is a review.

#### Consent for publication

Not applicable as it is a review.

#### Competing interests

The authors declare that they have no competing interests.

#### Author details

<sup>1</sup>Wits Reproductive Health and HIV Institute, Faculty of Health Sciences, University of the Witwatersrand, Johannesburg, South Africa. <sup>2</sup>South African Medical Research Council, Cape Town, South Africa. <sup>3</sup>Department of Pathology, Microbiology and Immunology, Vanderbilt University Medical Center, Nashville, TN, USA. <sup>4</sup>Division of Medical Education and Administration, Vanderbilt University School of Medicine, Nashville, TN, USA. <sup>5</sup>Department of Global Health and Development, Faculty of Public Health and Policy, London School of Hygiene & Tropical Medicine, London WC1H 9SH, UK. <sup>6</sup>Division of Pulmonology, Department of Medicine, Stellenbosch University and Tygerberg Hospital, Cape Town, South Africa. <sup>7</sup>Division of Health Systems and Public Health, Department of Global Health, Faculty of Medicine and Health Sciences, Stellenbosch University, Stellenbosch, South Africa. <sup>8</sup>Department of Population Health, Aga Khan University, Nairobi, Kenya. <sup>9</sup>Department of Public Health and Primary Care, International Centre for Reproductive Health (ICRH), Ghent University, Ghent, Belgium. <sup>10</sup>Department of Epidemiology and Preventive Medicine, Monash University, Melbourne, Australia. <sup>11</sup>Burnet Institute, Melbourne, Australia. <sup>12</sup>Wildlife Forensic Academy, Buffelsfontein Nature Reserve, Cape Town, South Africa. <sup>13</sup>The University of Sydney, Faculty of Medicine and Health, Sydney, South Africa. <sup>14</sup>Disease Elimination, Burnet Institute, Melbourne, Australia. <sup>15</sup>Department of Epidemiology and Preventive Medicine, Monash University, Melbourne, Australia.

Received: 15 April 2020 Accepted: 22 April 2020

Published online: 15 May 2020

#### References

1. Regly E. Italian doctors' fatalities reach tragic levels as they fight COVID-19 in overburdened hospitals. *The Globe and Mail*. 2020;<https://www.theglobeandmail.com/world/article-italian-doctors-fatalities-reach-tragic-levels-as-they-fight-covid-1/>.
2. Chen Q, Quan B, Li X, Gao G, Zheng W, Zhang J, et al. A report of clinical diagnosis and treatment of nine cases of coronavirus disease 2019. *J Med Virol*. 2020.
3. Adepoju P. Nigeria responds to COVID-19; first case detected in sub-Saharan Africa. *Nat Med*. 2020.
4. Haider N, Yavilinsky A, Simons D, Osman AY, Ntoumi F, Zumla A, et al. Passengers' destinations from China: low risk of novel coronavirus (2019-nCoV) transmission into Africa and South America. *Epidemiol Infect*. 2020; 148:e41.
5. Nkengasong JN, Mankoula W. Looming threat of COVID-19 infection in Africa: act collectively, and fast. *Lancet (London, England)*. 2020;395(10227): 841–2.
6. Gilbert M, Pullano G, Pinotti F, Valdano E, Poletto C, Boelle PY, et al. Preparedness and vulnerability of African countries against importations of COVID-19: a modelling study. *Lancet (London, England)*. 2020;395(10227): 871–7.
7. WHO. Electronic State Parties Self-Assessment Annual Reporting Tool. <https://extranet.who.int/e-spar>.
8. WHO. The 2018 update, Global Health Workforce Statistics, World Health Organization, Geneva (<http://www.who.int/hrh/statistics/hwfstats/>).
9. Murthy S, Leligdowicz A, Adhikari NK. Intensive care unit capacity in low-income countries: a systematic review. *PLoS One*. 2015;10(1):e0116949.
10. Joob B, Wiwanitkit V. COVID-19 in medical personnel: observation from Thailand. *The Journal of hospital infection*. 2020.
11. Wang D, Hu B, Hu C, Zhu F, Liu X, Zhang J, et al. Clinical Characteristics of 138 Hospitalized Patients With 2019 Novel coronavirus-infected pneumonia in Wuhan, China *JAMA* 2020.
12. Lai THT, Tang EWH, Chau SKY, Fung KSC, Li KKW. Stepping up infection control measures in ophthalmology during the novel coronavirus outbreak: an experience from Hong Kong. Graefes's archive for clinical and

- experimental ophthalmology = Albrecht von Graefes Archiv fur klinische und experimentelle Ophthalmologie. 2020.
13. Zhu H, Wang L, Fang C, Peng S, Zhang L, Chang G, et al. Clinical analysis of 10 neonates born to mothers with 2019-nCoV pneumonia. *Translational pediatrics*. 2020;9(1):51–60.
  14. Lu CW, Liu XF, Jia ZF. 2019-nCoV transmission through the ocular surface must not be ignored. *Lancet*. 2020.
  15. Seah I, Su X, Lingam G. Revisiting the dangers of the coronavirus in the ophthalmology practice. *Eye* (London, England). 2020.
  16. Tran K, Cimon K, Severn M, Pessoa-Silva CL, Conly J. Aerosol generating procedures and risk of transmission of acute respiratory infections to healthcare workers: a systematic review. *PLoS One*. 2012;7(4):e35797.
  17. Ghinai I, McPherson TD, Hunter JC, Kirking HL, Christiansen D, Joshi K, et al. First known person-to-person transmission of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) in the USA. *Lancet* (London, England). 2020.
  18. Marchand-Senechal X, Kozak R, Mubareka S, Salt N, Gubbay JB, Eshaghi A, et al. Diagnosis and Management of First Case of COVID-19 in Canada: lessons applied from SARS. *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America*. 2020.
  19. Ni L, Zhou L, Zhou M, Zhao J, Wang DW. Combination of western medicine and Chinese traditional patent medicine in treating a family case of COVID-19 in Wuhan. *Frontiers of medicine*. 2020.
  20. Cai SJ, Wu LL, Chen DF, Li YX, Liu YJ, Fan YQ, et al. [Analysis of bronchoscope-guided tracheal intubation in 12 cases with COVID-19 under the personal protective equipment with positive pressure protective hood]. *Zhonghua jie he he hu xi za zhi = Zhonghua jiehe he huxi zazhi = Chinese journal of tuberculosis and respiratory diseases*. 2020;43(0):E033.
  21. Mahase E. Covid-19: retired doctors could be asked to return to work, says Hancock. *BMJ* (Clinical research ed). 2020;368:m831.
  22. Mahase E. Covid-19: hoarding and misuse of protective gear is jeopardising the response, WHO warns. *BMJ* (Clinical research ed). 2020;368:m869.
  23. Hopman J, Allegranza B, Mehtar S. Managing COVID-19 in low- and middle-income countries. *JAMA*. 2020.
  24. Ran L, Chen X, Wang Y, Wu W, Zhang L, Tan X. Risk Factors of Healthcare Workers with Corona Virus Disease 2019: A Retrospective Cohort Study in a Designated Hospital of Wuhan in China *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America* 2020.
  25. Chinazzi M, Davis JT, Ajelli M, Gioannini C, Litvinova M, Merler S, et al. The effect of travel restrictions on the spread of the 2019 novel coronavirus (COVID-19) outbreak. *NY: Science* (New York; 2020).
  26. Kapata N, Ihekweazu C, Ntouni F, Raji T, Chanda-Kapata P, Mwaba P, et al. Is Africa prepared for tackling the COVID-19 (SARS-CoV-2) epidemic. Lessons from past outbreaks, ongoing pan-African public health efforts, and implications for the future. *International journal of infectious diseases : IJID : official publication of the International Society for Infectious Diseases*. 2020; 93:233–6.
  27. Ebola Gbalo Research G. Responding to the Ebola virus disease outbreak in DR Congo: when will we learn from Sierra Leone? *Lancet*. 2019;393(10191): 2647–50.
  28. Ren LL, Wang YM, Wu ZQ, Xiang ZC, Guo L, Xu T, et al. Identification of a novel coronavirus causing severe pneumonia in human: a descriptive study. *Chin Med J*. 2020.
  29. Soriano V, Barreiro P. Impact of new coronavirus epidemics on HIV-infected patients. *AIDS Rev*. 2020;22(1):57–8.
  30. Zhu F, Cao Y, Xu S, Zhou M. Co-infection of SARS-CoV-2 and HIV in a patient in Wuhan city, China. *J Med Virol*. 2020.
  31. Grobler L, Mehtar S, Dheda K, Adams S, Babatunde S, van der Walt M, et al. The epidemiology of tuberculosis in health care workers in South Africa: a systematic review. *BMC Health Serv Res*. 2016;16(1):416.
  32. Iacobucci G. Covid-19: GP surgeries close for two weeks after staff test positive. *BMJ* (Clinical research ed). 2020;368:m936.
  33. Huang JZ, Han MF, Luo TD, Ren AK, Zhou XP. [Mental health survey of 230 medical staff in a tertiary infectious disease hospital for COVID-19]. *Zhonghua lao dong wei sheng zhi ye bing za zhi = Zhonghua laodong weisheng zhiyebing zazhi = Chinese journal of industrial hygiene and occupational diseases*. 2020;38(0):E001.
  34. Kang L, Li Y, Hu S, Chen M, Yang C, Yang BX, et al. The mental health of medical workers in Wuhan, China dealing with the 2019 novel coronavirus. *Lancet Psychiatry*. 2020;7(3):e14.
  35. Catton H. Global challenges in health and health care for nurses and midwives everywhere. *Int Nurs Rev*. 2020;67(1):4–6.
  36. Hu Z, Song C, Xu C, Jin G, Chen Y, Xu X, et al. Clinical characteristics of 24 asymptomatic infections with COVID-19 screened among close contacts in Nanjing. *China Science China Life sciences*. 2020.
  37. Rose C, Am I. Part of the cure or Am I part of the disease? Keeping coronavirus out when a doctor comes home. *N Engl J Med*. 2020.
  38. Rodriguez-Morales AJ, Cardona-Ospina JA, Gutierrez-Ocampo E, Villamizar-Pena R, Holguin-Rivera Y, Escalera-Antezana JP, et al. Clinical, laboratory and imaging features of COVID-19: a systematic review and meta-analysis. *Travel Med Infect Dis*. 2020;101623.
  39. Adams JG, Walls RM. Supporting the health care workforce during the COVID-19 global epidemic. *JAMA*. 2020.
  40. McMahon SA, Ho LS, Brown H, Miller L, Ansumana R, Kennedy CE. Healthcare providers on the frontlines: a qualitative investigation of the social and emotional impact of delivering health services during Sierra Leone's Ebola epidemic. *Health Policy Plan*. 2016;31(9):1232–9.
  41. Koh D. Occupational risks for COVID-19 infection. *Occupational medicine (Oxford, England)*. 2020;70(1):3–5.
  42. Xiao H, Zhang Y, Kong D, Li S, Yang N. The effects of social support on sleep quality of medical staff treating patients with coronavirus disease 2019 (COVID-19) in January and February 2020 in China. *Medical science monitor : international medical journal of experimental and clinical research*. 2020;26:e923549.
  43. Schwartz DA. An analysis of 38 pregnant women with COVID-19, their newborn infants, and maternal-fetal transmission of SARS-CoV-2: maternal coronavirus infections and pregnancy outcomes. *Archives of pathology & laboratory medicine*. 2020.
  44. Schwartz J, King CC, Yen MY. Protecting health care workers during the COVID-19 coronavirus outbreak -lessons from Taiwan's SARS response. *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America*. 2020.
  45. Raven J, Wurie H, Witter S. Health workers' experiences of coping with the Ebola epidemic in Sierra Leone's health system: a qualitative study. *BMC Health Serv Res*. 2018;18(1):251.
  46. Parry J. China coronavirus: Hong Kong health staff strike to demand border closure as city records first death. *BMJ* (Clinical research ed). 2020;368:m454.
  47. Iacobucci G. Covid-19: medical schools are urged to fast-track final year students. *BMJ* (Clinical research ed). 2020;368:m1064.
  48. Rimmer A. Covid-19: GPs call for appraisals and CQC inspections to be suspended. *BMJ* (Clinical research ed). 2020;368:m1038.
  49. McMahon SA, Ho LS, Scott K, Brown H, Miller L, Ratnayake R, et al. "We and the nurses are now working with one voice": How community leaders and health committee members describe their role in Sierra Leone's Ebola response. *BMC Health Serv Res*. 2017;17(1):495.
  50. Moberly T. Chinese premier rallies medics in coronavirus fight. *BMJ* (Clinical research ed). 2020;368:m343.
  51. Rimmer A. Covid-19: junior doctor calls on colleagues to gather supplies for staff working long hours. *BMJ* (Clinical research ed). 2020;368:m1072.
  52. Phan LT, Maita D, Mortiz DC, Weber R, Fritzen-Pedicini C, Bleasdale SC, et al. Personal protective equipment doffing practices of healthcare workers. *J Occup Environ Hyg*. 2019;16(8):575–81.
  53. Makoni M. Africa prepares for coronavirus. *Lancet* (London, England). 2020; 395(10223):483.

## Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.