



in the postoperative period and bringing the profession of specialist surgeon into disrepute is provocative and laughable. I'm surprised that this hearing favoured the 'expertise' presented by a retired surgical academic, who by his own admission had done very little laparoscopic surgery, over that of a professor and a specialist intensivist currently at the top of their careers. Surely experienced medical personnel should be appointed to hear public grievances? Those of us who are members of the Medical Protection Society (MPS) are concerned that in a case like this the legal representatives failed dismally. This case should have won hands down.

There is a perception among the lay public and litigation lawyers that as most of us have some form of medical protection there's no harm in 'having a go'! MPS reports suggest that medicolegal claims in South Africa have escalated way above rates in the rest of the world. It is my impression that we in clinical medicine are seen as an easily milked cash cow. We are under continual pressure from medical aids, hospital groups and the media – and now our very own HPCSA.

I sincerely hope that the colleague in question has the stamina to exercise his rights and appeal against the findings of the HPCSA, and that his surgical association reacts strongly to this disgraceful decision.

**Barry Penn**

31 Mountain Road  
Claremont  
7708

**Well done, SAMA's Industrial Relations Unit!**

**To the Editor:** It is reassuring to know that the South African Medical Association, through its Industrial Relations Unit, has the capacity to assist doctors, especially hospital doctors, should any have reason to believe that they have been subjected to unfair labour practices.

My own experience is that about 3 years after retirement I was phoned by the hospital concerned and told that I had received a salary increase some 2 or 3 years before retirement for which I had not been paid. I was told that if I supplied my bank details I would be paid. Having heard nothing for a year I made further enquiries, only to be told that the provincial health department concern had no money.

I had no recourse other than through the SAMA Industrial Labour Unit, which was entirely successful in obtaining my back pay.

I don't hesitate to recommend to all doctors that they should become SAMA members, for this and many other reasons!

**T L Williams**

PO Box 86  
Lidgettton  
3270

**Selective serotonin reuptake inhibitors in children and adolescents**

**To the Editor:** The introduction of the selective serotonin reuptake inhibitors (SSRIs) was widely viewed as an important advance in clinical psychopharmacology, not only because of their broad-spectrum efficacy but also because of their tolerability and safety advantages, particularly compared with the older tricyclic antidepressants (TCAs) and monoamine oxidase inhibitors (MAOIs). Subsequently there has been considerable controversy about this class of agents, partly because of concerns about the extent to which they have been injudiciously prescribed for 'cosmetic' problems rather than for genuine psychopathology,<sup>1</sup> and partly because of concerns regarding their adverse effects. Most recently, attention has been paid to the appropriate use of SSRIs in children and adolescents.

The 'Drug Alert' published by the National Adverse Drug Event Monitoring Centre in the September 2005 SAMJ<sup>2</sup> is singularly unhelpful in this regard. The report takes a far more conservative stance than that taken by regulators in the USA, the UK and the EU; it may be misleading by implication and omission; and (if followed to the letter) it may cause child and adolescent psychiatric patients significant harm.

The 'Drug Alert' warns practitioners on four points. First, 'None of the SSRIs are currently approved in South Africa for any indication in children and adolescents.' It should be pointed out, however, that fluoxetine is registered with the US Food and Drug Administration (FDA) for child and adolescent depression and several of the SSRIs (fluvoxamine, sertraline, and fluoxetine) are also FDA-registered for child and adolescent obsessive-compulsive disorder (OCD).<sup>3</sup> Practitioners should also be aware that decisions about whether to submit pharmaceutical agents to the Medicines Control Council for registration of particular indications may often be made on the basis of cost rather than scientific or clinical considerations.

Second, 'SSRIs have been associated with an increase in the risk of suicidal thinking and behaviour (suicidality) in children and adolescents with MDD [major depressive disorder] and other psychiatric disorders.' However, as the drug alert also states, 'no suicides occurred' in the 24 trials involving over 4 400 patients. In addition, a systematic review<sup>4</sup> published recently found no significant difference in the risk of suicide in patients taking SSRIs compared with those taking TCAs. As several commentators have pointed out, patients with overt suicidal ideation are excluded from clinical trials and the heterogeneous nature of the trial designs employed (use of different definitions and assessments of self-harm in different study populations) further contributes to the difficulty of interpreting the data. The trials quoted were not designed to address the question of whether SSRIs increase suicidal ideation, and cannot in fact do so.<sup>5</sup>



Third, products 'containing citalopram, escitalopram, fluvoxamine, paroxetine, sertraline and venlafaxine are contraindicated in children under 18 years of age'. While many regulatory authorities have cautioned practitioners about possible adverse events of these agents, so far none have taken so inflexible an approach as to say that they are all 'contraindicated' in children and adolescents. The European Medicines Agency in the same press release referred to by the MCC concedes that: 'It is recognised that a doctor may sometimes take a decision based on the individual clinical needs of a child or an adolescent to use these products for the treatment of depression or anxiety. The CHMP [Committee for Medicinal Products for Human Use] is recommending that in these cases patients be monitored carefully for the appearance of suicidal behaviour, self-harm or hostility, particularly at the beginning of treatment.'<sup>6</sup>


It must be remembered that TCAs are not effective in child and adolescent depression, while SSRIs have been shown to be effective (on their own and in combination with cognitive behavioural therapy) for child and adolescent depression, and as a sole intervention in some childhood anxiety disorders, especially OCD.<sup>7-10</sup> Fluoxetine, the only SSRI not on the MCC's 'contraindicated' list, is not always tolerated by younger patients, especially those who have severe anxiety, and is also not available in any form except 20 mg capsules in most of the state sector, making appropriate dosing difficult. The effect

of implementing the MCC recommendations as published would be to deprive child and adolescent patients with these disorders of available, effective and potentially life-saving alternative medications.

Depression is a common, disabling, and potentially fatal disorder that is substantially and unequivocally associated with suicide and deliberate self-harm in children and adolescents. A large recent Swedish forensic study<sup>11</sup> showed that only 13% and 4% respectively of children and adolescents who actually committed suicide had received any antidepressant medication at all; youth suicide is far more likely to occur due to untreated depression than the adverse effects of any antidepressant. It should also be noted that since the advent of the SSRIs in 1988, suicide rates have decreased significantly in many countries.<sup>12,13</sup>

Fourth, 'Discontinuation of SSRIs, especially abrupt discontinuation, commonly leads to significant withdrawal symptoms.' While it is true that discontinuation symptoms can be seen with some SSRIs after abrupt discontinuation, this warning runs the risk of conflating SSRIs, which are not associated with dependence, with medication classes such as the benzodiazepines, where dependence is an acknowledged risk.

In view of the above points we urge practitioners to note that these warnings from the MCC do not seem to reflect a balanced view of current scientific thinking, and may have significant adverse consequences for patients if implemented



## Medical Solutions

### Sony's Components

Sony offers five main components in its Medical Imaging Network solution. These can be used either standalone or as part of a complete networked Solution for hospitals:

### Fast and accurate diagnosis

Sony Medical Imaging Network solutions give 'any time, any place' access to vital patient information and images that they need. With the electronic transfer of high quality digital images throughout a networked environment, delivering fast and accurate diagnosis and treatment, as well as cost savings and enhanced patient experience.

### More hospital efficiency

Designed to cater for the needs of today's demanding hospital environment where integration across all hospital departments is crucial to improving the speed of patient diagnosis and treatment.

- Image acquisition
- Image query & retrieval
- Image printing
- Image archiving & storage
- Image presentation



For the complete range of Medical Solutions by Sony:  
Visit us on: [www.sony.co.za](http://www.sony.co.za) or Contact Will Klapper on: 011-690-3312



inflexibly. We would argue that the SSRIs do have an important role to play in psychiatric practice, including that of child and adolescent psychiatry, and that clinicians should, as always, balance the benefits and risks for any particular patient and keep the interests of the patient paramount. Obtaining an expert opinion from a child and adolescent psychiatrist would be useful in situations where a practitioner is unsure. We hope that the MCC will urgently reconsider the wording of this directive, in the interests of the many young patients who may otherwise be deprived of necessary and effective treatments.

**S M Hawkrige****S Seedat****R Emsley****P Carey****D J Stein**

*Cross-University Brain-Behaviour Initiative  
Departments of Psychiatry and Mental Health  
University of Cape Town and  
Stellenbosch University*

- Stein DJ. Cosmetic psychopharmacology of anxiety: Bioethical considerations. *Current Psychiatry Reports* 2005; 7: 237-238.
- National Adverse Drug Event Monitoring Centre, Medicines Control Council. Drug Alert. Warning: selective serotonin reuptake inhibitors in children and adolescents. *S Afr Med J* 2005; 95: 660.
- FDA Public Health Advisory, 15 October 2004. Suicidality in children and adolescents being treated with antidepressants. <http://www.fda.gov/cder/drug/antidepressants/SSRIPHA200410.htm> (last accessed 28 September 2005).
- Fergusson D, Doucette S, Cranley Glass K, et al. Association between suicide attempts and selective serotonin reuptake inhibitors: systematic review of randomised controlled trials. *BMJ* 2005; 330: 396-403.
- Greenhill LL, Vitiello B, Riddle MA, et al. Review of safety assessment methods used in pediatric psychopharmacology. *J Am Acad Child Adolesc Psychiatry* 2003; 42: 627-633.
- European Medicines Agency/CHMP/12981805/2005: Press Release: European Medicines Agency finalises review of antidepressants in children and adolescents. <http://www.emea.eu.int/pdfs/human/press/pr/12891805en.pdf> (last accessed 28 September 2005).
- Hazell P, O'Connell D, Heathcote D, Henry D. Tricyclic drugs for depression in children and adolescents (Cochrane Review). In: *The Cochrane Library*, issue 1. Chichester: John Wiley & Sons, 2004.
- Treatment for Adolescents With Depression (TADS) Team. Fluoxetine, cognitive-behavioral therapy, and their combination for adolescents with depression. Treatment for Adolescents With Depression Study (TADS) randomized controlled trial. *JAMA* 2004; 292: 807-820.
- Research Unit on Pediatric Psychopharmacology Anxiety Study Group (RUPP). Fluvoxamine for the treatment of anxiety disorders in children and adolescents. *N Engl J Med* 2001; 344: 1279-1285.
- Geller DA, Biederman J, Stewart SE, et al. Which SSRI? A meta-analysis of pharmacotherapy trials in pediatric obsessive-compulsive disorder. *Am J Psychiatry* 2003; 160: 1919-1928.
- Isacson G, Holmgren P, Ahlner J. Selective serotonin reuptake inhibitor antidepressants and the risk of suicide: a controlled forensic database study of 14 857 suicides. *Acta Psychiatr Scand* 2005; 111: 286-290.
- Gibbons RD, Hur K, Bhaumik DK, et al. The relationship between antidepressant medication use and rate of suicide. *Arch Gen Psychiatry* 2005; 62: 165-172.
- Licinio J, Wong ML. Depression, antidepressants and suicidality: a critical appraisal. *Nat Rev Drug Discov* 2005; 4: 165-171.

## Rocephin – the thin end of the wedge

**To the Editor:** I am sure I speak for many colleagues who have been cajoled, squeezed, begged and pressurised by medical aids, hospital administrators, pharmacists and representatives to stick to formularies to the point at which, backs against the wall, we say, 'No more!' The 'thin edge of the wedge' for me is the Rocephin issue.

I have grown up with Rocephin. For me, rightly or wrongly, Rocephin is the drug that diminishes my anxiety, oh so slightly, over the safety of my patients; that may prevent meningitis

secondary to sinus surgery; and that may stop a child's otorrhoea when other drugs have failed and prevent him or her from developing mastoiditis.

But despite evidence that many generics are inferior,<sup>1</sup> the pressure is on me not to use Rocephin, but to prescribe a generic, for the vast saving of about R5 per dose.

When will this pressure stop? If I surrender on the Rocephin issue, what comes next? Are the medical aid administrators, the hospital managers and the pharmaceutical buyers willing to share the medical risk that I face on a daily basis? Will they stand in the dock with me one day, and admit to using medications that are not proven to be equal in efficacy?

We hear so much about 'sharing risk' nowadays. The best way those pressurising me to use 'their' not 'my' choice of drugs can share risk, is by sharing my risk. What about, as a suggestion, paying all or part of my medical indemnity insurance?

I'm not looking for handouts. I'm not looking for perverse incentives. I don't even know, or care, whether the 'local' generic ceftriaxone is equivalent to Rocephin. For a saving of only R5 a dose? I learn from one generic manufacturer that their local factory manufacturing quality is excellent, and then I hear that the drug I am interested in is allegedly manufactured in Turkey, shipped to Germany, and then to South Africa where it is distributed.

For me there is a line I dare not cross, on the other side of which my autonomy stands for nothing. Where I still have a choice, I must fight to maintain it, lest that thin edge of the wedge be pushed in further and further, until it becomes a thick edge, and then a wall. 'Rocephin' for me is that issue, and I will not budge. It is time we as a collective organisation of medical professionals stand up and say to those who would manipulate us 'This far and no further'.

**Martin Young**

*5 Hedge Street  
Knysna  
6571*

- Lambert PA, Conway BR. Pharmaceutical quality of ceftriaxone generic products compared with Rocephin. *J Chemother* 2003; 15: 357-368.

## Problematic childhood atopic eczema consensus document?

**To the Editor:** The childhood atopic eczema consensus document published as part 2 of the June issue of the *SAMJ*<sup>1</sup> is problematic and necessitates the following comments.

A discussion of the controversy over the definition of atopic eczema is necessary, particularly in view of the recommendations that pertain to allergy testing. Without this the rest of the article is open to misinterpretation. When dermatologists speak of atopic eczema they mean a clinical diagnosis. As