

A new model for the pathophysiology of Alzheimer's disease

Aluminium toxicity is exacerbated by hydrogen peroxide and attenuated by an amyloid protein fragment and melatonin

Susan J van Rensburg, Willie M U Daniels,
Felix C V Potocnik, Johann M van Zyl,
Joshua J F Taljaard, Robin A Emsley

Objectives. Although Alzheimer's disease (AD) is the leading cause of dementia in developed countries, there is an as yet unexplained lower prevalence of the disease in parts of Africa. AD is characterised by a catastrophic loss of neurons; free radicals (oxidative toxins) have been implicated in the destruction of the cells through the process of lipid peroxidative damage of cell membranes. Previously aluminium (Al) and a fragment of beta amyloid (A β 25 - 35) were shown to exacerbate free-radical damage, while melatonin reduced this effect. The aim of the present study was: (i) to investigate the conditions determining the toxicity of Al and A β 25 - 35; and (ii) to assess whether melatonin could attenuate the damage done by both aluminium and the amyloid fragment, thus suggesting a pathway for the aetiology of AD.

Design. An *in vitro* model system was used in which free radicals were generated, causing lipid peroxidation of platelet membranes, thus simulating the disease process found in the brain.

Results. 1. Al and A β 25 - 35 caused lipid peroxidation in the presence of the iron (II) ion (Fe²⁺), Al being more toxic than A β 25 - 35. 2. A β 25 - 35 attenuated the lipid peroxidation promoted by Al. 3. Hydrogen peroxide (H₂O₂) greatly exacerbated the toxicity of Al and A β 25 - 35. 4. Melatonin prevented lipid peroxidation by Al and A β 25 - 35 in the absence of H₂O₂, but only reduced the process when H₂O₂ was present.

Conclusions. In the light of the results obtained from the present study, the following hypotheses are formulated.

1. In AD, excessive quantities of Al are taken up into the

brain, where the Al exacerbates iron-induced lipid peroxidation in the lysosomes. 2. In response, the normal synthetic pathway of amyloid protein is altered to produce A β fragments which attenuate the toxicity of Al. In the process of sequestering the Al and iron, immature plaques are formed in the brain. 3. Microglia are activated, in an attempt to destroy the plaques by secreting reactive oxygen species such as H₂O₂. At this point in the disease process, lipid peroxidation causes a catastrophic loss of brain cells. 4. Melatonin, together with other free radical scavengers in the brain, reduces the free-radical damage caused by Al and A β , except in the latter stages of the disease process. Since melatonin is produced by the pineal gland only in the dark, the excess of electric light in developed countries may help explain why AD is more prevalent in these countries than in rural Africa.

S Afr Med J 1997; 87: 1111-1115.

Alzheimer's disease (AD) is characterised by the loss of cells in certain brain areas, as revealed by temporal lobe-orientated X-ray computed tomography (CT) scans.¹ By the time of clinical AD diagnosis, about 90% of the hippocampus will have been destroyed. AD is the leading cause of dementia in developed countries; however, there is an as yet unexplained lower prevalence of the disease in parts of Africa,² even though other dementias such as vascular dementia and alcohol-related dementia are common.

An understanding of the action of certain chemically reactive molecules called free radicals has provided evidence for their involvement in the destruction of brain cells.³ Free radicals are toxic byproducts of cellular biochemical reactions, as found for example when adenosine triphosphate (ATP) is generated in the mitochondria.⁴ Free radicals do, however, serve a purpose in the body, e.g. when cells of the immune system produce free radicals in order to destroy micro-organisms, or digest unwanted cell material.⁵ These reactions are usually carefully controlled for, if left to themselves, free radicals attack living cells, damaging phospholipids (by causing lipid peroxidation), proteins and even DNA.⁶

Chemically speaking, free radicals are ions and molecules which take part in electron transfer reactions.⁷ An example is the iron (II) ion (Fe²⁺), which transfers an electron to molecular oxygen, thus producing the superoxide radical (O₂⁻). Superoxide can dismutate to form hydrogen peroxide and, again through the intervention of Fe²⁺, form hydroxyl radicals (OH[•]).⁷

Superoxide and hydroxyl radicals are extremely damaging to living cells, especially the cell membranes. Membranes play an integral part in cell function and health; besides keeping the cell contents intact, they also determine which molecules enter and leave cells, and in themselves take part in chemical reactions. If the damaged membranes cannot be repaired, the cells die. The principal constituent of membranes, the phospholipids, are particularly vulnerable to free-radical damage (lipid peroxidation) and, as such, provide a model which can be utilised for studying the abovementioned reactions *in vitro*.³

Departments of Chemical Pathology, Medical Physiology, Psychiatry and Pharmacology, University of Stellenbosch, Tygerberg, W Cape

Susan J van Rensburg, PhD

Joshua J F Taljaard, MD

Willie M U Daniels, PhD

Felix C V Potocnik, FFPsych (SA)

Robin A Emsley, MD

Johann M van Zyl, PhD

In living cells, any free radicals formed are normally neutralised by anti-radical defence systems, e.g. free-radical scavengers (anti-oxidants), enzymes such as glutathione peroxidase and superoxide dismutase, and iron-binding proteins such as transferrin.⁸ However, if these defence systems are breached, for example, when transferrin does not bind the iron completely, the iron becomes available for free-radical production. A genetic isoform of transferrin (Tf C2), which has a decreased iron-binding capacity, is more frequently present in AD patients than controls and would therefore abet this process.⁹ But, in order to combat increased concentrations of free radicals, other levels of defence may exist, for example in the form of the hormone, melatonin, which protects membranes against lipid peroxidation.¹⁰

Aluminium (Al)¹¹ and beta amyloid (A β)¹² have both been shown to exacerbate lipid peroxidation and are implicated in the aetiology of AD, although the mechanistic pathway has not been established. In the present study, an *in vitro* system was used: (i) to determine the conditions under which Al and a fragment of the A β protein, A β 25 - 35, cause lipid peroxidation; and (ii) to determine whether melatonin could attenuate the damage done by either the Al or the amyloid fragment.

Methods and materials

Chemicals. Diphenylhexatriene (DPH), 2-thiobarbituric acid (TBA), butylated hydroxytoluene (BHT) and A β 25 - 35 were obtained from Sigma Chemical Co; Al₂(SO₄)₃.18H₂O was obtained from Merck.

Preparation of membranes. Platelets suspended in citrate-dextrose buffer, isolated as described previously,³ were obtained from healthy donors. Erythrocytes were removed by centrifugation at 800 g for 5 minutes. Platelets were then collected by centrifugation at 3 500 g for 15 minutes. The pellets were resuspended in 0.01M phosphate-buffered saline (PBS), pH 7.4. Cell counts were done on a Technicon H2 Blood Analyser, which showed that there were no erythrocytes or white blood cells present. Membranes were prepared by sonication of the suspension with an MSE sonicator for 10 seconds at an amplitude of 5 microns. The fragmented membranes were not washed, in order to maintain their integrity. The protein concentration of the suspension was 0.07 mg/ml, as determined by the method of Lowry *et al.*¹³

Determination of lipid peroxidation. Of the membrane suspension, 100 μ l aliquots were pipetted into silicon glass test tubes (Kimble). The following reagents were then added in sequence, to give final concentrations as indicated: ascorbate (160 μ M), EDTA (10 μ M), Al₂(SO₄)₃.18H₂O (0.1 - 100 μ M), A β 25 - 35 (0.01 - 100 μ M) or melatonin (1 mM); FeSO₄ (40 μ M) and hydrogen peroxide (H₂O₂; 0.5 - 500 μ M). The total volume per tube was adjusted to 1 000 μ l with PBS (pH 5.5, to keep the Al in solution).

The tubes were incubated at 37°C for 30 minutes. The following were then added: 1 ml of a 1% (w/v) TBA solution plus 1 ml of 2.8% (w/v) trichloro-acetic acid. BHT (0.01%) was also included to neutralise metal-catalysed auto-oxidation of lipids during heating of the mixtures at 100°C

for 10 minutes. After cooling, the formation of thiobarbituric reactive substances (TBARS) was determined by reading the chromogen at 535 nm in a Beckman DU 640 spectrophotometer. Appropriate corrections to the absorbance readings were made for the presence of FeSO₄-EDTA and ascorbate in the reaction mixtures. Absorbance readings at 535 nm are proportional to the TBARS formed, and are therefore relative values reflecting lipid peroxidation.

Statistical analysis. The statistical significance of the results was determined by the Mann-Whitney U-test. Results are the mean \pm SEM of 6 observations.

Results

1. **Al caused lipid peroxidation only in the presence of Fe²⁺.** In Fig. 1 (column A), 100 μ M Al₂(SO₄)₃.18H₂O (i.e. 8.1 μ M Al) did not cause lipid peroxidation on its own when iron was absent, but when 40 μ M iron sulphate and Al were present simultaneously (Fig. 1, column B), a significant increase in lipid peroxidation ($P < 0.05$) was observed.




Fig. 1. Lipid peroxidation recorded in the presence of: A — 100 μ M Al sulphate, without iron, B — 100 μ M Al sulphate plus 40 μ M iron sulphate, C — 100 μ M Al sulphate, 40 μ M iron sulphate plus 0.5 mM H₂O₂, D — 100 μ M Al sulphate, 40 μ M iron sulphate, 0.5 mM H₂O₂ plus 1 mM melatonin, E — 100 μ M Al sulphate, 40 μ M iron sulphate, plus 1 mM melatonin. B was significantly higher than A ($P < 0.05$); C was significantly higher than B ($P < 0.005$); D was significantly lower than C ($P < 0.005$); E was significantly lower than B ($P < 0.005$) (Mann-Whitney U-test).

2. **A β 25 - 35 similarly promoted lipid peroxidation when iron was present.** In the presence of Fe²⁺, 0.01 - 1.0 μ M A β 25 - 35 increased the lipid peroxidation significantly ($P < 0.005$) compared with A β 25 - 35 at the same concentrations without iron (Fig. 2).

3. **Al was more toxic than the amyloid protein fraction.** In Fig. 3, the relative toxicity of 0.1 - 100 μ M A β 25 - 35 and 0.1 - 100 μ M Al₂(SO₄)₃.18H₂O (i.e. 0.008 - 8 μ M Al) was compared. The Al produced significantly ($P < 0.005$) more

lipid peroxidation than A β 25 - 35. The amount of Al used in the present study (0.216 - 216 μ g/l) was compatible with the Al concentration recorded previously in the blood of AD patients (8 - 104 μ g/l).¹⁴




Fig. 2. Lipid peroxidation recorded in the presence of increasing concentrations of A β 25 - 35, (●) with and (○) without 40 μ M iron sulphate. A β 25 - 35 with iron produced significantly more lipid peroxidation than A β 25 - 35 without iron ($P < 0.005$).




Fig. 3. Lipid peroxidation recorded in the presence of increasing concentrations of (●) Al sulphate and (○) A β 25 - 35. Al sulphate produced significantly more lipid peroxidation than A β 25 - 35 ($P < 0.005$).

4. The amyloid protein fraction attenuated Al toxicity.

When Al and A β 25 - 35 were present simultaneously (Fig. 4), the lipid peroxidation produced was significantly ($P = 0.005$) less than with Al on its own. The results from two separate experiments were combined. However, when lipid peroxidation was high, as in the presence of H₂O₂ (Fig. 5), this effect was steadily lost.




Fig. 4. Lipid peroxidation recorded in the presence of: (●) increasing concentrations of Al sulphate; and (○) 100 μ M Al sulphate plus increasing concentrations of A β 25 - 35. A β 25 - 35 attenuated the toxicity of Al significantly ($P < 0.005$; Mann-Whitney U-test).




Fig. 5. Lipid peroxidation in the presence of increasing concentrations of H₂O₂: (□) with 100 μ M Al sulphate; and (■) with 100 μ M Al sulphate plus 100 μ M A β 25 - 35.

5. **H₂O₂ greatly exacerbated the toxicity of Al and A β 25 - 35.** Fig. 5 shows that there is a significant increase in lipid peroxidation with increasing H₂O₂ concentrations ($P < 0.005$). In addition, the attenuating effect of A β 25 - 35 in the presence of Al is not seen in the presence of H₂O₂, although the difference between the effects of Al on its own and the combination of Al and A β 25 - 35 was not significant (Fig. 5). Fig. 1 (column C) demonstrates significantly higher lipid peroxidation in the presence of 100 μ M Al sulphate, iron sulphate, and 0.5 mM H₂O₂, compared with iron on its own (Fig. 1, column B).

6. **Melatonin eliminated lipid peroxidation caused by Al and A β 25 - 35 in the absence of H₂O₂.** In Fig. 1 (column E) melatonin (1 mM) eliminates lipid peroxidation caused by Al and A β 25 - 35, but in the presence of H₂O₂ (0.5 mM; Fig. 1, column D), the toxicity was only reduced, not eliminated, even though the concentration of melatonin was twice that of H₂O₂.

Discussion

In the current *in vitro* system, both Al and A β 25 - 35 exacerbated lipid peroxidation in the presence of iron, Al being more toxic than A β 25 - 35. Furthermore, A β 25 - 35 attenuated the toxicity of Al. H₂O₂ greatly increased lipid peroxidation in the presence of both Al and A β 25 - 35, except in the absence of iron. Furthermore, melatonin eliminated lipid peroxidation when H₂O₂ was absent, and decreased lipid peroxidation in the presence of H₂O₂.

The above results, in conjunction with previously published data, suggest the following sequence of events in the aetiology of AD. Firstly, it has previously been established that there is an increased flow of Al into the brain in AD. Taylor *et al.*¹⁴ demonstrated an increase in blood Al after an Al citrate drink was ingested by AD patients. In this group the mean blood Al concentrations increased from the baseline value of 8.6 μ g/l to 104 μ g/l within 60 minutes, whereas in age-matched controls, the blood Al level rose from 7.7 μ g/l to only 37.9 μ g/l over the same time period. The threshold allowing Al entry into the brain may also be lower in AD patients, given that Ward and Mason demonstrated that the concentration of Al in both the hippocampus and cortex of post-mortem brain samples from AD patients was 10 times higher than that of non-demented controls.¹⁵ This suggests that the increased absorption of Al from the gut also results in more Al being deposited in the brains of these AD patients.

Although Al crosses the blood-brain barrier attached to molecules such as citrate, the principal carrier is transferrin (Tf).¹⁶ In the brain Al is freed from Tf in the lysosomes⁸ where the ambient pH is low,¹⁷ which keeps Al in solution, i.e. in a more neurotoxic form. The Al ions then attach themselves to the headgroups of the phospholipid molecules of cellular membranes where, on account of their positive charge, they create artificial openings.⁸ Normally, the headgroups would form a barrier against attack by free radicals, protecting the fatty acid side-chains inside the lipid bilayer, but the 'pores' allow hydroxyl radicals to reach, attack and damage the lipids (lipid peroxidation), thus changing the fluidity of the membranes.¹⁸ It is recognised that abnormalities of the lysosomal system precede the appearance of tangles and

plaques in the neurons of AD patients.¹⁹

The role of amyloid in AD has been a subject of debate. The functions of the amyloid precursor protein (APP) include tissue repair in the brain, cell adhesion, and survival and growth of neurons *in vitro*.²⁰ Shigematsu and McGeer found that APP accumulated in neuronal processes and microglia following intracerebral administration of Al salts in rat brain.²¹ It has also been shown that amyloid fragments are generated in lysosomes.²²

The abovementioned lipid peroxidation causes a 4 Angstrom units reduction in the lipid bilayer width,²³ exposing the transmembrane portion of the APP to proteolytic secretase enzymes which cleave the APP at a site which is normally not accessible to these enzymes,²⁴ producing insoluble A β fragments. Increased production (i.e. not only an accumulation²¹) of APP may well occur in response to increased Al levels in neurons. We hypothesise that APP, being a 'sticky protein', is meant to bind Al and sequester it. Owing to the membrane damage and faulty cleavage of the APP, large amounts of insoluble A β are produced instead, although A β also binds to metals such as Al and iron.²⁵ Sequestering of these metals by amyloid would result in a temporary decrease in lipid peroxidation. Increasing levels of Al result in the formation of immature amyloid plaques, which 'hold' the Al, because the brain cannot destroy a metal. At this stage AD progresses slowly.

The plaques represent foreign material in the brain. We hypothesise that this activates the microglia (brain phagocytes), which start to secrete reactive oxygen species such as H₂O₂.⁵ Microglia, whose presence in the centre of mature plaques has been demonstrated, are capable of destroying nervous system tissue for scavenging purposes.²⁶ However, since the plaques consist of insoluble amyloid in the β -sheet conformation, they will be resistant to degradation by the microglia, which continue to produce more H₂O₂. As is evident from the results of the present study, H₂O₂ greatly exacerbates lipid peroxidation, and in these circumstances, A β does not have a protective effect against Al-initiated lipid peroxidation (Fig. 5). This may be the cause of the catastrophic phase of cell loss seen in the brains of AD patients (15.1% per year, v. 1.5% per year in healthy controls).²⁷

Finally, it was demonstrated that melatonin eliminates lipid peroxidation by Al and A β 25 - 35, but in the absence of H₂O₂. When 0.5 mM H₂O₂ is present, melatonin attenuates but does not neutralise lipid peroxidation.

Melatonin, a hormone secreted by the pineal gland, is of interest in diseases associated with ageing, such as AD, since it is known to decrease in concentration with age.²⁸ The functions of melatonin include regulation of secretion of other hormones (i.e. regulation of the body's biological clock),²⁹ and it has recently been implicated as a further level of defence against lipid peroxidation.¹⁰ Melatonin is secreted only in the dark. Provided the person is in total darkness, melatonin levels start to rise at 21h00, peaking between 02h00 and 04h00, whereafter they decline. In the presence of electric light of sufficient intensity at night time, no melatonin is produced;²⁹ hence the availability of melatonin as free-radical scavenger may be compromised in Western societies, where the use of electric light is excessive. This may help explain the finding of a low prevalence of AD in rural communities in Africa.

Conclusions

In the light of the above hypothesis, a few suggestions may be offered as to how we should combat AD.

1. It could be suggested that Al be removed from the diet. Al in solution has been found to be more toxic than Al in food,³⁰ and drinking tea ('rooibos' herbal tea excluded) has been shown to increase Al levels!³¹ However, it may be more profitable in the end to find the reason why the natural barriers against Al uptake in the gut and the blood-brain barrier are lifted in AD. In AD, more than anything else, prevention will be better than cure.

2. During the phase where AD progresses slowly, it will be of use to ensure that all protective systems against free-radical damage are functioning optimally. This would include taking a wide range of anti-oxidant supplements along with cognition-enhancing drugs. In addition, the damaged phospholipids would have to be replaced by supplementation with essential fatty acids.³²

3. In the catastrophic phase, where the brain is in an inflammatory state, anti-inflammatory drugs may be of use.³³

4. Melatonin, together with other free-radical scavengers in the brain, reduces free radical damage. Since melatonin is produced by the pineal gland only in the dark, we may have to reconsider the effects late-night living may have on our health in old age. In addition, supplementation with melatonin could be considered.¹⁰

We gratefully acknowledge the financial support given by the Cape Provincial Administration and the Medical Research Council of South Africa.

REFERENCES

- Jobst KA, Hindley NJ, King E, Smith AD. The diagnosis of Alzheimer's disease: A question of image? *J Clin Psychiatry* 1994; **55** (11): suppl, 22-31.
- Hendrie HC, Osuntokun BO, Hall KS, et al. Prevalence of Alzheimer's disease and dementia in two communities: Nigerian Africans and African Americans. *Am J Psychiatry* 1995; **152**: 1485-1492.
- Van Rensburg SJ, Daniels WMU, Van Zyl JM, Potocnik FCV, Van der Walt BJ, Taljaard JF. Lipid peroxidation and platelet membrane fluidity — implications for Alzheimer's disease? *NeuroReport* 1994; **5**: 2221-2224.
- Sjodin B, Hellsten-Westling Y, Apple FS. Biochemical mechanisms for oxygen free radical formation during exercise. *Sports Med* 1990; **10**: 236-254.
- Babior BM. Oxidants from phagocytes: agents of defense and destruction. *Blood* 1984; **64**: 959-966.
- Hensley K, Butterfield DA, Hall N, et al. Reactive oxygen species as causal agents in the neurotoxicity of the Alzheimer's disease-associated amyloid beta peptide. *Ann NY Acad Sci* 1996; **786**: 120-134.
- Halliwell B, Gutteridge JMC. *Free Radicals in Biology and Medicine*. 2nd ed. Oxford: Clarendon Press, 1989: 200.
- Van Rensburg SJ, Carstens ME, Potocnik FCV, Van der Spuy G, Van der Walt BJ, Taljaard JF. Transferrin C2 and Alzheimer's disease: Another piece of the puzzle found? *Med Hypoth* 1995; **44**: 268-272.
- Van Rensburg SJ, Carstens ME, Aucamp AK, Taljaard JF. Increased frequency of the transferrin C2 subtype in Alzheimer's disease. *NeuroReport* 1993; **4**: 1269-1271.
- Daniels WMU, Van Rensburg SJ, Van Zyl JM, Van der Walt BJ, Taljaard JF. Free radical scavenging effects of melatonin and serotonin: possible mechanism. *NeuroReport* 1996; **7**: 1593-1596.
- Gutteridge JMC, Quinlan GJ, Clark I, Halliwell B. Aluminium salts accelerate peroxidation of membrane lipids stimulated by iron salts. *Biochim Biophys Acta* 1985; **835**: 441-447.
- Butterfield DA, Hensley K, Harris M, Mattson M, Carney J. β -amyloid peptide free radical fragments initiate synaptosomal lipoperoxidation in a sequence-specific fashion: implications to Alzheimer's disease. *Biochem Biophys Res Commun* 1994; **200**: 710-715.
- Lowry OH, Rosebrough NJ, Farr AL, Randall RJ. Protein measurement with the Folin phenol reagent. *J Biol Chem* 1951; **193**: 265-275.
- Taylor GA, Ferrier IN, McLoughlin IJ, et al. Gastrointestinal absorption of aluminium in Alzheimer's disease: response to aluminium citrate. *Age Ageing* 1992; **21**: 81-90.
- Ward NI, Mason JA. Neutron activation analysis techniques for identifying elemental status in Alzheimer's disease. *Journal of Radioanalytical and Nuclear Chemistry* 1987; **113**: 515-526.
- Birchall JD, Chappell JS. Aluminium, chemical physiology, and Alzheimer's disease. *Lancet* 1988; **2**: 1008-1010.
- Orci L, Ravazzola M, Anderson RGW. The condensing vacuole of exocrine cells is more acidic than the mature secretory vesicle. *Nature* 1987; **326**: 77-79.
- Van Rensburg SJ, Carstens ME, Potocnik FCV, Aucamp AK, Taljaard JF, Kock KR. Membrane fluidity of platelets and erythrocytes in patients with Alzheimer's disease and the effect of small amounts of aluminium on platelet and erythrocyte membranes. *Neurochem Res* 1992; **17**: 825-829.
- Nixon RA, Cataldo AM, Paskevich PA, Hamilton DJ, Wheelock TR, Kanaley-Andrews L. The lysosomal system in neurons. Involvement at multiple stages of Alzheimer's disease pathogenesis. *Ann N Y Acad Sci* 1992; **674**: 65-88.
- Ashall F, Goate AM. Role of the beta amyloid precursor protein in Alzheimer's disease. *Trends Biochem Sci* 1994; **19**: 42-46.
- Shigematsu K, McGeer PL. Accumulation of amyloid precursor protein in damaged neuronal processes and microglia following intracerebral administration of aluminum salts. *Brain Res* 1992; **593**: 117-123.
- Haass C, Koo EH, Mellon A, Hung AY, Selkoe DJ. Targeting of cell-surface beta-amyloid precursor protein to lysosomes: alternative processing into amyloid-bearing fragments. *Nature* 1992; **357**: 500-503.
- Mason RP, Shoemaker WJ, Shajenko L, Chambers TE, Herbet LG. Evidence for changes in the Alzheimer's disease brain cortical membrane structure mediated by cholesterol. *Neurobiol Aging* 1992; **13**: 413-419.
- Dyrks T, Mack E, Masters CL, Beyreuther K. Membrane insertion prevents aggregation of precursor fragments containing the β /A4 sequence of Alzheimer's disease. In: Iqbal K, McLachlan DRC, Winblad B, Wisniewski HM, eds. *Alzheimer's Disease: Basic Mechanisms, Diagnosis and Therapeutic Strategies*. Chichester: John Wiley & Sons, 1991: 281-287.
- Mantyh PW, Ghilardi JR, Rogers S, et al. Aluminium, iron, and zinc ions promote aggregation of physiological concentrations of β -amyloid peptide. *J Neurochem* 1993; **61**: 1171-1174.
- McRae A, Dahlstrom A, Ling EA. Microglia in neurodegenerative disorders: emphasis on Alzheimer's disease. *Gerontology* 1997; **43**: 95-108.
- Jobst KA, Smith AD, Szatmari M, et al. Rapidly progressing atrophy of medial temporal lobe in Alzheimer's disease. *Lancet* 1994; **343**: 829-830.
- Reiter RJ, Pablos MI, Agapito TT, Guerrero JM. Melatonin in the context of the free radical theory of aging. *Ann NY Acad Sci* 1996; **786**: 362-378.
- Brzezinski A. Melatonin in humans. *N Engl J Med* 1997; **336**: 186-195.
- Armstrong RA, Anderson J, Cowburn JD, Cox J, Blair JA. Aluminium administered in drinking water but not in the diet influences bipterin metabolism in the rodent. *Biol Chem Hoppe Seyler* 1992; **373**: 1075-1078.
- Koch KR, Pougnet MAB, De Villiers S, Monteagudo F. Increased urinary excretion of Al after drinking tea. *Nature* 1988; **333**: 122.
- Van Rhijn AG, Prior CA, Corrigan FM. Dietary supplementation with zinc sulphate, sodium selenite and fatty acids in early dementia of Alzheimer's type. *Journal of Nutritional Medicine* 1990; **1**: 259-266.
- Breitner JCS. Onset of Alzheimer's disease: Influence of genes and environmental factors, including anti-inflammatory drugs. In: Heston LL, ed. *Progress in Alzheimer's Disease and Similar Conditions*. Washington, DC: American Psychiatric Press, 1997: 189-197.

Accepted 21 July 1997.