

results have proved the study to be of value: (i) an awareness among senior management staff that information is a vital resource which should be personally planned and controlled by them; (ii) a renewed appreciation of the goals of the hospital and its managers, their functions and problems; (iii) an evolutionary approach to systems development, which allows for the sequential phasing in of subsystems, using an information architecture; and (iv) the infusion of potentially far-reaching concepts regarding information systems management, resource allocation and academic development of medical informatics.

In simpler terms, this study has provided a plan which identifies and supports the hospital's short- and long-term information needs. This plan has been completely accepted by members of the hospital management, who were involved at each stage of development and are committed to its implementation.

It is suggested that a similar study would benefit any large health care institution wishing to re-evaluate its data processing functions. Given the required amount of time, commitment and expertise, such a study can offer management a highly relevant and practical information systems plan and a mechanism for gaining maximum benefit from a costly but vital resource.

We wish to express our gratitude to Dr H.-R. Sanders, Chief Medical Superintendent of Groote Schuur Hospital, who sponsored the HIPS, for permission to publish and for her enthusiastic support of the study. We are also grateful to the members of the study team and to the interviewees for their participation.

REFERENCES

1. IBM Corporation. *Business Systems Planning: Information Planning Guide*. 3rd ed. White Plains: IBM Technical Publications Department, 1981.
2. United States Postal Service. *Business Systems Planning Report 1980*: (unpublished report).
3. Schmitz H. *Hospital Information Systems*. Germantown: Aspen Systems Corporation, 1979.
4. Rockart JF. Chief executives define their own data needs. *Harvard Business Review* 1979; March-April: 115-126.
5. Bakker AR. Implementation approach and evaluation of the Leiden University Hospital Information System. In: Shires DB, Wolf HK, eds. *Medinfo 77*. Amsterdam: North-Holland, 1977: 943-947.
6. Watson RJ. A large-scale professionally oriented medical information system — five years later. *J Med Syst* 1977; 1: 3-21.
7. Reichertz PL, Rienhoff O. The Medical System Hannover. In: Shires DB, Wolf HK, eds. *Medinfo 77*. Amsterdam: North-Holland, 1977: 1068-1069.
8. Reichertz PL. Future developments of data processing in health care. *Methods Inf Med* 1982; 21: 55-58.

The role of phosphatidyl-glycerol in the determination of fetal lung maturity

A. J. SMITH

Summary

Phosphatidyl-glycerol (PG) and the lecithin/sphingomyelin (L/S) ratio were determined on amniotic fluid from 140 patients to evaluate the role of PG estimation as a functional fetal lung maturity test. PG was absent without associated respiratory distress syndrome (RDS) in 17 cases (12%) and present in 123; 3 of the infants born to these patients developed slight hyaline membrane disease. There were no neonatal deaths. When both PG and the L/S ratio were determined the accuracy of prediction of the RDS improved from 87% to 93%. PG determination should not be used as a substitute for the L/S ratio, but should be used in conjunction with it in high-risk situations.

S Afr Med J 1983; 63: 45-47.

Department of Obstetrics and Gynaecology, Tygerberg Hospital, Parowvallei, CP
A. J. SMITH, M.B. CH.B.

Date received: 8 January 1982.

Since the introduction of routine amniocentesis with determination of the lecithin/sphingomyelin (L/S) ratio, the incidence of iatrogenic neonatal respiratory distress syndrome (RDS) has shown a marked decrease. Kulovich *et al.*¹ in a compilation of several reports involving more than 1 200 neonates showed that mature L/S ratios predicted absence of the RDS with about 98% accuracy. According to Harvey *et al.*² a low L/S ratio (less than 2,0) is only 54% accurate in predicting the development of the RDS. Infants born to patients with diabetes mellitus and Rh iso-immunization are at risk of developing the RDS, even if the L/S ratio is 2,0 or more. Kulovich *et al.*¹ therefore developed a 'lung profile', which includes the L/S ratio and the results of acetone precipitable lecithin, phosphatidyl-glycerol (PG) and phosphatidyl-inositol (PI) estimation. This lung profile improved the ability to diagnose unsuspected lung maturity in infants born to mothers with low L/S ratios to 93%.

Surfactant is a complex mixture of lipids (80 - 90% by weight), protein (10 - 20% by weight) and carbohydrates (1 - 2% by weight).³ Eighty to ninety per cent of the lipid fraction is phospholipid, of which 70 - 80% is phosphatidylcholine (lecithin). Of the minor phospholipids, PG is the most abundant, accounting for up to 10% of the surfactant phospholipids in the adult. It is synthesized in the microsomal fraction of the type II alveolar cell. The remaining phospholipids are PI, sphingomyelin, phosphatidylethanolamine (PE) and others of minor importance.

The purpose of this study was to evaluate the role of PG determination as a functional fetal lung maturity test.

Patients and methods

Over a period of 13 months (July 1980 - August 1981) L/S ratio and PG determination and 'bubble tests' (foam stability tests) were performed on amniotic fluid from 182 unselected patients. For the patient to qualify for inclusion in this study, delivery of the infant had to occur within 72 hours of the amniocentesis. According to these criteria, 140 patients were found suitable for inclusion. A full record was kept for each patient (name, folder number, date of amniocentesis, diagnosis, gestational age, L/S ratio, bubble test result, presence or absence of PG, method of delivery and date of delivery). The sex, mass, Apgar score and respiratory status of each infant were recorded at birth. Neonates were evaluated by the paediatric staff, and particular attention was paid to the development of the RDS. The diagnosis of RDS was based on clinical features (tachypnoea of 60/min, rib retraction, sternal recession and grunting), the radiological appearance of the lung fields, and capillary blood gas values. The results of L/S ratio and PG determination and the bubble test were correlated with the patient's clinical diagnosis and the development of RDS in the neonate.

The L/S ratio was determined according to a modification of the technique described by Gluck *et al.*,⁴ while the foam stability test was performed according to the method of Clements *et al.*⁵ PG was determined by a modification of the technique used by Hallman and Teramo.⁶

Silica gel H plates, impregnated with 5% (NH₄)₂SO₄ and activated in an oven at 120°C for 15 minutes, were used for thin-layer chromatography. The phospholipids were isolated from the amniotic fluid by centrifugation after addition of methanol and chloroform. Acetone precipitation of the surface-active phospholipids was included in this technique.³ The precipitate was dissolved in chloroform and applied to the activated silica gel plates, along with the standards of the different phospholipid fractions, namely L/S, PG, PI, and PE.

The plates were first run in one direction, in a solvent of chloroform-methanol-acetic acid-H₂O (130 ml-50 ml-16 ml-8 ml), and dried with warm air; subsequently they were run in another direction in a solvent of tetrahydrofuran-methylalcohol-liquid ammoniac (100 ml-74 ml-16 ml-11 ml). They were then dried and left in a bath containing bromothymol blue detector (bromothymol blue 0,150 g, boric acid 0,005 g, Na₂CO₃ 50 g, made up to 1 000 ml with distilled water). Finally, they were blotted dry and dried in an oven.

The plates were interpreted with a Vitatron densitometer. The L/S ratio as well as the PG/PI ratio and the PG/S (PG/sphingomyelin) ratio were plotted in this way. For the purpose of this study PG was only reported as present or absent, along with the L/S ratio.

The majority of the tests were performed on the same day as the amniocentesis. If the amniotic fluid sample was collected too late it was kept in a refrigerator at -4°C and analysed the following day.

Results

The clinical diagnoses of the 140 patients are summarized in Table I. Diagnoses included diabetes mellitus (41 cases), hypertensive diseases (39 cases), intra-uterine growth retardation (13 cases), and Rh iso-immunization (9 cases); there were also 23 patients undergoing elective caesarean section. PG was absent without associated RDS in 17 cases (12,14% of the total) (Table II). Of these patients 3 had diabetes mellitus, 11 had hypertension, and 3 had other problems. PG was found to be present in 123 cases. Nine infants in this group developed the RDS (6,42%), although there were no neonatal deaths. Only 3 of these 9 infants had mild hyaline membrane disease (HMD); none required assisted ventilation. The other 6 had aspiration pneumonia (5 cases) and transient tachypnoea of the newborn (1 case), probably due to hypoglycaemia. These infants were born to mothers with diabetes (3 patients), hypertension (2), Rh sensitization (1) and other diseases (3) (Table II). Of the 123 patients in whom PG was found to be present, 3 had an L/S ratio below 2,0. PG was found to be absent in 3 of the 41 patients with diabetes. None of their infants developed RDS. Conversely, 3 of the infants born to the 38 diabetic mothers in whom PG was present developed RDS, although none had HMD. None of the diabetics had an L/S ratio of less than 2,0. The L/S ratio was 2,0 - 2,5 in 3 cases and 2,6 - 3,0 in 15; the rest had an L/S ratio of less than 3,0.

TABLE I. CLINICAL DIAGNOSES OF THE PATIENTS

| | No. of patients |
|---------------------------------------|-----------------|
| Diabetes mellitus* | 41 |
| Whites class A | 17 |
| Whites class B | 5 |
| Whites class C | 3 |
| Not classified | 16 |
| Hypertension and pre-eclampsia | 39 |
| Rh iso-immunization | 9 |
| Diverse | 51 |
| Elective caesarean section | 23 |
| Postmaturity | 3 |
| Intra-uterine growth retardation | 13 |
| Other | 12 |
| Total | 140 |

*Class A = chemical diabetes; B = maturity-onset diabetes (age over 20 years), duration < 10 years, no vascular lesions; C = age 10-19 years at onset, and diabetes of 10-19 years' duration.

The accuracy of prediction of lung maturity is shown in Table III. As a single factor, the L/S ratio was 87% accurate, compared with the 86% of PG alone. This difference is statistically insignificant. In combination the accuracy of prediction was 93%.

TABLE II. RDS IN RELATION TO AMNIOTIC FLUID L/S RATIO AND PRESENCE OR ABSENCE OF PG IN 140 PATIENTS

| | PG present with RDS | PG absent without RDS | L/S < 2,0 PG present without RDS |
|----------------------------|---------------------|-----------------------|----------------------------------|
| Diabetes | 3 | 3 | 0 |
| Hypertension | 2 | 11 | 2 |
| RH iso-immunization | 1 | 0 | 2 |
| Diverse | 3 | 3 | 4 |
| Total | 9* (6,42%) | 17 (12,14%) | 8 (5,51%) |

*See text for further discussion.

TABLE III. ACCURACY OF PREDICTION OF LUNG MATURITY WITH L/S RATIO, PRESENCE OF PG AND A COMBINATION OF BOTH

| | Mature L/S (≥ 2) | PG present | Mature L/S and/or PG present |
|------------------------------------|-------------------------|---------------------------|------------------------------|
| Diabetes | 39 | 38 | 39 |
| RH iso-immunization | 7 | 9 | 9 |
| Hypertension | 35 | 28 | 37 |
| Diverse | 44 | 48 | 48 |
| Total | 125 | 123 | 133 |
| HMD | | 3 cases (see text) | |
| Mature (no HMD) | 122 | 120 | 130 |
| Accuracy of prediction of maturity | 97,60% (122 of 125) | 97,56% (120 of 123) | 97,74% (130 of 133) |
| Accuracy of prediction of total | 87,14% (122 of 140) | 85,71% (120 of 140) | 92,85% (130 of 140) |

TABLE IV. PREVALENCE OF RDS AND HMD IN RELATION TO AMNIOTIC FLUID L/S RATIO AND PRESENCE OR ABSENCE OF PG

| L/S ratio | All cases | | | | | PG present | | | | | PG absent | |
|--------------|------------|----------|------------|----------|------------|------------|----------|------------|----------|------------|-----------|--------------|
| | Total | RDS | | HMD | | Total | RDS | | HMD | | Total | No. with RDS |
| | | No. | % | No. | % | | No. | % | No. | % | | |
| $\geq 2,0$ | 125 | 6 | 4,8 | 2 | 1,6 | 120 | 6 | 5 | 2 | 1,7 | 17 | 0 |
| $< 2,0$ | 15 | 3 | 20 | 1 | 6,7 | 3 | 3 | 100 | 1 | 33,3 | 0 | 0 |
| Total | 140 | 9 | 6,4 | 3 | 2,1 | 123 | 9 | 7,3 | 3 | 2,4 | 17 | 0 |

Discussion

For predicting the RDS in the newborn the presence of PG had a false-negative value of 12% and a false-positive value of 7,3% (Tables II and IV). Similar results were obtained when the L/S ratio was used alone (Table III). Combining these two improved the accuracy of prediction of the RDS from 87% to 93% (Table III). This compares favourably with the 93% accuracy of Gluck's lung profile.

When compared with the L/S ratio PG determination has the advantage that PG is not affected by contamination of amniotic fluid with blood, vaginal secretions or meconium.^{1,7-9} Disadvantages of PG determination include the fact that it is technically more difficult, more time-consuming and more expensive than determination of the L/S ratio alone. The present study confirms previous findings that the mere presence of PG in amniotic fluid is a good indication of functional biochemical fetal lung maturity.^{1,10,11} However, the absence of PG is not an indication that the RDS will develop. Seventeen patients (12%), of whom 11 had hypertensive disease, had a false-negative PG result. This was not found in previous studies, and needs further investigation.

In a previous study from our laboratory the foam test was compared with the L/S ratio as a determinant of lung maturity.¹² For the foam test a false-positive rate of 8,3% and a false-negative rate of 27,1% were found. This high false-negative rate made the foam test suitable for screening purposes, but the L/S ratio remained the more important diagnostic test. The present study confirmed these findings; it was possible to perform the foam test in 100 cases, and it was found to have a false-positive rate of 2% and a false-negative rate of 20%.

We conclude that PG determination is not a substitute for the L/S ratio but should be used in conjunction with it in high-risk situations. This will improve the accuracy of prediction of fetal lung maturity by 6%. When amniotic fluid is contaminated with blood or meconium determination of PG is still reliable, whereas the L/S ratio and foam test become unreliable.

REFERENCES

- Kulovich MV, Hallman MB, Gluck L. The lung profile. I. Normal pregnancy. *Am J Obstet Gynecol* 1979; **135**: 57-63.
- Harvey D, Parkinson CE, Campbell S. Risk of respiratory distress syndrome. *Lancet* 1975; **i**: 42.
- Hack M, Fanaroff AA, Klaus MH, Mendelawitz BD, Merkatz IR. Neonatal respiratory distress following elective delivery: a preventable disease? *Am J Obstet Gynecol* 1976; **126**: 43-47.
- Gluck L, Kulovich MV, Borer RC. Estimates of fetal lung maturity. *Clin Perinatol* 1974; **1**: 125-139.
- Clements JA, Platzker ACG, Tierney DF *et al.* Assessment of the risk of the respiratory distress syndrome by a rapid test for surfactant in amniotic fluid. *N Engl J Med* 1972; **286**: 1077-1081.
- Hallman M, Teramo K. Amniotic fluid phospholipid profile as predictor of fetal maturity in diabetic pregnancies. *Obstet Gynecol* 1979; **54**: 703-707.
- Bustos R, Kulovich M, Gluck L *et al.* Significance of phosphatidylglycerol in amniotic fluid in complicated pregnancies. *Am J Obstet Gynecol* 1979; **133**: 899-903.
- Whittle MJ, Wilson AI, Whitfield CR, Paton RD, Logan RW. Amniotic fluid phospholipid profile determined by two-dimensional thin-layer chromatography as index of fetal lung maturity. *Br Med J* 1981; **282**: 428-430.
- Tsai M, Marshall JG. Phosphatidylglycerol in 261 samples of amniotic fluid from normal and diabetic pregnancies, as measured by one-dimensional thin-layer chromatography. *Clin Chem* 1980; **25**: 682-685.
- O'Brien WF, Cefalo RC. Clinical applicability of amniotic fluid tests for fetal pulmonary maturity. *Am J Obstet Gynecol* 1980; **136**: 135-143.
- Cunningham MD, Desai NS, Thompson SA, Greene JM. Amniotic fluid phosphatidylglycerol in diabetic pregnancies. *Am J Obstet Gynecol* 1978; **131**: 719-724.
- Van der Merwe HJ, Cranna EJ. Evaluering van die lesitien/sfingomielien verhouding en die skuimtoets as fetale longmatuureitstoets. *SAfr Med J* 1978; **53**: 54-57.