
IS MASS CORRELATED WITH LENGTH AMONG RED MILLIPEDES *CENTROBOLUS* COOK, 1897?

| Mark I. Cooper* |

* University of Stellenbosch, SOUTH AFRICA. E-mail: cm.i@aol.com, ORCID ID: 0000-0001-7861-895X.

[Cooper, M. I. 2023. Is mass correlated with length among red millipedes *Centrobolus* Cook, 1897?. Munis Entomology & Zoology, 18 (1): ...-...]

ABSTRACT: Body mass variation was correlated with male and female length in the red millipede genus *Centrobolus*. Mass and length ($r=0.86$, Z score= 2.87 , $n=8$, $p<0.01$) in males ($r=0.97$, Z score= 2.99 , $n=5$, $p<0.01$) and females ($r=0.96$, Z score= 2.72 , $n=5$, $p<0.01$) were correlated. *C. inscriptus* females have the largest length (67.4 mm) and the highest mass (2.61 g) while *C. digrammus* has the shortest length (49.9 mm) and the lightest mass (0.68 g).

KEY WORDS: Dimorphic, female, length, mass, morphology, ratio

Red millipedes are found in the southern African subregion with northern limits on the east coast being about -17° latitude S and southern limits being -35° latitude S. They are well represented in the littoral forests of the eastern half of the subcontinent (Lawrence, 1967). It consists of taxonomically important species with 12 species considered threatened and includes nine vulnerable and three endangered species (Mailula, 2021). It occurs in all the forests of the coastal belt from the Cape Peninsula to Beira in Mocambique (Lawrence, 1967). In an analysis of four morphometric factors (male and female length and width), four positive correlations were found in the 22 species of *Centrobolus* (Cooper, 2022). Here, body mass is correlated with body length in *Centrobolus* Cook, 1897 (Cook, 1897; Hamer, 1998; Mailula, 2021). These worm-like millipedes have female-biased sexual size dimorphism (Cooper, 1998; 2014; 2015; 2017; 2018; 2021). One hypothesis is body length correlates with body mass.

MATERIALS AND METHODS

Four valid species were identified as belonging to the genus *Centrobolus* Cook, 1897 (Table 1). Millipede collection localities were obtained from the Mating dynamics of South African forest millipedes *Centrobolus* (Diplopoda: Pachybolidae) (Cooper, 1998). Body mass was correlated with body width. Mean body mass was obtained by calculating the weight with a Mettler AC 100 Autobalance. Body length was measured by placing individuals alongside a plastic rule (mm) (Table 1). Body length and body mass were checked for correlations using the Pearson Correlation Coefficient calculator (<https://www.gigacalculator.com/calculators/correlation-coefficient-calculator.php>).

Table 1. Species in the millipede genus *Centrobolus* Cook, 1897, with body length and body mass. Male followed by female body masses are given with sample sizes in parentheses. Two sets of measurements are included for *C. inscriptus*.

| Species | length | Location | Body mass (g) |
|----------------------|-----------|--------------------|---------------|
| <i>C. fulgidus</i> | 56.2 (11) | Richards Bay | 1.29 (11) |
| | 63.5 (11) | | 1.97 (11) |
| <i>C. inscriptus</i> | 67.4 (88) | Mtunzini | 2.48 (88) |
| | 67.0 (56) | | 2.00 (56) |
| | 63.0 (88) | | 2.27 (88) |
| | 65.0 (41) | | 2.61 (41) |
| <i>C. ruber</i> | 57.8 (18) | Port Shepstone | 1.28 (18) |
| | 62.3 (18) | | 2.00 (18) |
| <i>C. digrammus</i> | 49.9 (6) | Admirals Waterfall | 0.68 (6) |
| | 54.5 (6) | | 1.02 (6) |

RESULTS

Body mass and length (Fig. 1: $r=0.93760869$, Z score= 4.54511480 , $n=10$, $p=0.00000275$) in males (Fig. 2: $r=0.97112526$, Z score= 2.98638973 , $n=5$, $p=0.00141153$) and females (Fig. 3: $r=0.95830497$, Z score= 2.72197404 , $n=5$, $p=0.00324471$) were related. Body length was normally distributed ($D=0.21722$, $n=10$, $p=0.65755$). Body mass was normally distributed ($D=0.23289$, $n=10$, $p=0.57311$). Female length was normally distributed ($D=0.36888$, $n=5$, $p=0.40287$). Male length was normally distributed ($D=0.26334$, $n=5$, $p=0.80102$). Male mass was normally distributed ($D=25851$, $n=5$, $p=0.81732$). Body mass was normally distributed ($D=0.23289$, $n=10$, $p=0.57311$). Female mass was normally distributed ($D=0.29699$, $n=5$, $p=0.67584$). Male width was normally distributed ($D=21396$, $n=5$, $p=0.93695$).




Figure 1. Relationship between body length (x-axis) and body mass (y-axis) in *Centrobolus* Cook, 1897.




Figure 2. Relationship between male body length (x-axis) and male body mass (y-axis) in *Centrobolus* Cook, 1897.




Figure 3. Relationship between female body length (x-axis) and female body mass (y-axis) in *Centrobolus* Cook, 1897.

DISCUSSION

A significant positive relationship between mass and length was found. *C. inscriptus* females have the largest length (67.4 mm), and the highest mass (2.61 g) while *C. digrammus* has the shortest length (49.9 mm), and the lightest mass (0.68 g). This study supports body mass and body length as predictable in *Centrobolus*.

Size-assortative mating based on body mass and body length determine the variance in millipede polygynandrous mating systems across a size gradient with lower body length occurring in lighter individuals. Body length increases with heavier body mass which is explained by a combination of copulatory guarding and fecundity selection.

In Danish red foxes *Vulpes vulpes* SSD varies with body mass at different population densities (Pagh et al. 2018). In the wandering albatross *Diomedea exulans* SSD varies with body mass with different adult survival, breeding probability, breeding success, chick mass, and juvenile survival (Cornioley et al. 2017). In microtines mass rather than length is used to measure SSD (Boonstra et al. 1993). In five species of *Drosophila*, the degree of dimorphism in development time was significantly correlated with dry weight and fecundity, with lighter species tending to be more dimorphic for development time as well as more fecund, both in absolute terms and terms of fecundity per unit weight (Bharathi et al. 2004). In Red-Eared Sliders *Trachemys scripta elegans* sexual dimorphism of size in favor of females is maintained for variables related to mass and size (Gradela et al. 2017). Millipedes resemble the patterns of mass and sexual dimorphism in adult dragonflies (Anholt et al. 2011). Sexual dimorphism in mass and protein content of the forelimb muscles is known in the northern leopard frog *Rana pipiens* (Blackburn, 1992). The influence of mass and mating systems has been documented in the most dimorphic mammals (Weckerly, 1998). Morphological variance in the polygynandrous reproductive system of male and female millipedes means lighter individuals have lower body width or heavier individuals have greater body width pointing toward the evolution of condition-dependent SSD (Bonduriansky, 2007).

CONCLUSION

Length varied systematically with mass in *Centrobolus*. Morphological variance in the polygynandrous reproductive systems of *Centrobolus* has lighter individuals with shorter body length, or conversely heavier individuals have greater body length. Mass impacts the moments of inertia.

LITERATURE CITED

- Anholt, B., Marden, J. & Jenkins, D. 2011. Patterns of Mass Gain and Sexual Dimorphism in Adult Dragonflies (Insecta, Odonata). Canadian Journal of Zoology, 69 (5): 1156-1163.
- Bharathi, N. S., Prasad, N. G., Shakarad, M. & Joshi, A. 2004. Correlates of sexual dimorphism for dry weight and development time in five species of *Drosophila*. Journal of Zoology, 264 (1): 87-95.
- Blackburn, D. 1992. Sexual dimorphism in mass and protein content of the forelimb muscles of the northern leopard frog, *Rana pipiens*. Canadian Journal of Zoology, 70 (4): 670-674.
- Bonduriansky, R. 2007. The Evolution of Condition-Dependent Sexual Dimorphism. The American Naturalist, 169 (1): 9-19.
- Boonstra, R., Gilbert, B. S. & Krebs, C. J. R. 1993. Mating Systems and Sexual Dimorphism in Mass in Microtines. Journal of Mammalogy, 74 (1): 224-229.
- Cook, O. F. 1897. New relatives of *Spirobolus giganteus*. Brandtia (A series of occasional papers on Diplopoda and other Arthropoda), 18 (2): 73-75.

- Cooper, M. I.** 1998. Mating dynamics of South African forest millipedes *Centrobolus* (Diplopoda: Pachybolidae). The University of Cape Town, 1-141.
- Cooper, M. I.** 2014. Sexual size dimorphism and corroboration of Rensch's rule in *Chersastus* millipedes. *Journal of Entomology and Zoology Studies*, 2 (6): 264-266.
- Cooper, M. I.** 2015. Competition affected by re-mating interval in a myriapod. *Journal of Entomology and Zoology Studies*, 3 (4): 77-78.
- Cooper, M. I.** 2016. The relative sexual size dimorphism of *Centrobolus inscriptus* compared to 18 congeners. *Journal of Entomology and Zoology Studies*, 4 (6): 504-505.
- Cooper, M. I.** 2017. Copulation and sexual size dimorphism in worm like millipedes. *Journal of Entomology and Zoology Studies*, 5 (3): 1264-1266.
- Cooper, M. I.** 2017. Relative sexual size dimorphism in *Centrobolus digrammus* (Pocock) compared to 18 congeners. *Journal of Entomology and Zoology Studies*, 5 (2): 1558-1560.
- Cooper, M. I.** 2017. Relative sexual size dimorphism in *Centrobolus fulgidus* (Lawrence) compared to 18 congeners. *Journal of Entomology and Zoology Studies*, 5 (3): 77-79.
- Cooper, M. I.** 2017. Relative sexual size dimorphism *Centrobolus ruber* (Attems) compared to 18 congeners. *Journal of Entomology and Zoology Studies*, 5 (3): 180-182.
- Cooper, M. I.** 2017. Re-assessment of Rensch's rule in *Centrobolus*. *Journal of Entomology and Zoology Studies*, 5 (6): 2408-2410.
- Cooper, M. I.** 2018. *Centrobolus anulatus* (Attems, 1934) reversed sexual size dimorphism. *Journal of Entomology and Zoology Studies*, 6 (4): 1569-1572.
- Cooper, M. I.** 2018. Sexual size dimorphism and the rejection of Rensch's rule in Diplopoda. *Journal of Entomology and Zoology Studies*, 6 (1): 1582-1587.
- Cooper, M. I.** 2018. Allometry for sexual dimorphism in millipedes. *Journal of Entomology and Zoology Studies*, 6 (1): 91-96.
- Cooper, M. I.** 2018. Trigonulid size dimorphism breaks Rensch. *Journal of Entomology and Zoology Studies*, 6 (3): 1232-1234.
- Cooper, M. I.** 2021. Mass covaries with volume in forest millipedes *Centrobolus* Cook, 1897. *Journal of Entomology and Zoology Studies*, 9 (6): 190-192.
- Cooper, M. I.** 2022. Length and Width Correlations in *Centrobolus* Cook, 1897. *New Visions in Biological Science*, 9: 39-45.
- Cornioley, T., Jenouvrier, S., Borger, L., Weimerskirch, H. & Ozgul, A.** 2017. Fathers matter: male body mass affects life-history traits in a size-dimorphic seabird. *Proceedings of the Royal Society B*, 284 (1854): 20170397.
- Gradela, A., Santiago, T. O. C., Pires, I. C., de Castro Souza Silva, A., de Souza, L. C., de Faria, M. D., Neto, J. P. & Milanelo, L.** 2017. Sexual Dimorphism in Red-Eared Sliders (*Trachemys scripta elegans*) from the Wild Animal Triage Center of the Tiete Ecological Park, São Paulo, Brazil. *Acta Scientiae Veterinariae*, 45 (1468): 1-10.
- Hamer, M. L.** 1998. Checklist of Southern African millipedes (Myriapoda: Diplopoda). *Annals of the Natal Museum*, 39 (1): 11-82.
- Lawrence, R. F.** 1967. The Spiroboloidea (Diplopoda) of the eastern half of Southern Africa*. *Annals of the Natal Museum*, 18 (3): 607-646.
- Mailula, R. P.** 2021. Taxonomic revision and Red List assessment of the 'red millipede' genus *Centrobolus* (Spirobolida: Pachybolidae) of South Africa. University of Kwazulu natal, xxiii+289.
- Pagh, S., Hansen, M. S., Jensen, B., Pertoldi, C. & Chriél, M.** 2018. Variability in body mass and sexual dimorphism in Danish red foxes (*Vulpes vulpes*) in relation to population density. *Zoology and Ecology*, 28 (1): 1-9.
- Weckerly, F. W.** 1998. Sexual-Size Dimorphism: Influence of Mass and Mating Systems in the Most Dimorphic Mammals. *Journal of Mammalogy*, 79 (1): 33-52.