

Cyclopia Extracts Act as ER α Antagonists and ER β Agonists, *In Vitro* and *In Vivo*

Koch Visser, Morné Mortimer, Ann Louw*

Department of Biochemistry, University of Stellenbosch, Matieland, Stellenbosch, Republic of South Africa

Abstract

Hormone replacement therapy associated risks, and the concomitant reluctance of usage, has instigated the search for new generations of estrogen analogues that would maintain estrogen benefits without associated risks. Furthermore, if these analogues display chemo-preventative properties in breast and endometrial tissues it would be of great value. Both the selective estrogen receptor modulators as well as the selective estrogen receptor subtype modulators have been proposed as estrogen analogues with improved risk profiles. Phytoestrogen containing extracts of *Cyclopia*, an indigenous South African fynbos plant used to prepare Honeybush tea may serve as a source of new estrogen analogues. In this study three extracts, P104, SM6Met, and cup-of-tea, from two species of *Cyclopia*, *C. genistoides* and *C. subternata*, were evaluated for ER subtype specific agonism and antagonism both in transactivation and transrepression. For transactivation, the *Cyclopia* extracts displayed ER α antagonism and ER β agonism when ER subtypes were expressed separately, however, when co-expressed only agonism was uniformly observed. In contrast, for transrepression, this uniform behavior was lost, with some extracts (P104) displaying uniform agonism, while others (SM6Met) displayed antagonism when subtypes were expressed separately and agonism when co-expressed. In addition, breast cancer cell proliferation assays indicate that extracts antagonize cell proliferation in the presence of estrogen at lower concentrations than that required for proliferation. Furthermore, lack of uterine growth and delayed vaginal opening in an immature rat uterotrophic model validates the ER α antagonism of extracts observed *in vitro* and supports the potential of the *Cyclopia* extracts as a source of estrogen analogues with a reduced risk profile.

Citation: Visser K, Mortimer M, Louw A (2013) *Cyclopia* Extracts Act as ER α Antagonists and ER β Agonists, *In Vitro* and *In Vivo*. PLoS ONE 8(11): e79223. doi:10.1371/journal.pone.0079223

Editor: Wei Xu, University of Wisconsin - Madison, United States of America

Received: July 31, 2013; **Accepted:** September 20, 2013; **Published:** November 4, 2013

Copyright: © 2013 Visser et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: The authors would like to thank the Medical Research Council (MRC) and the Cancer Association of South Africa (CANSA) for financial support to A. L. (grant for projects entitled "Cyclopia Phytoestrogens" and "Cyclopia and breast cancer") and for financial support to K. V. and M. M. the National Research Foundation (NRF) and the Harry Crossley foundation is acknowledged. Any opinion, findings and conclusions or recommendations expressed in this material are those of the author(s) and therefore the funders do not accept any liability in regard thereto. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: The authors have declared that no competing interests exist.

* E-mail: al@sun.ac.za

Introduction

Hormone replacement therapy (HRT), estrogens alone or in combination with progestins, is traditionally prescribed to women undergoing menopausal transition to alleviate symptoms associated with menopause [1], such as hot flashes, night sweats, sleeping problems, vaginal dryness, and osteoporosis [2-4]. However, a number of side effects have been associated with the use of HRT, for example, an increased occurrence of breast cancer [5,6], vaginal bleeding [7], and heart disease or strokes [6,8]. These side effects have led to reluctance among concerned consumers to use HRT and instigated a search for new estrogen analogues with an improved risk profile. Furthermore, it would be of great value if these analogues should also display chemo-preventative properties in breast tissue [9,10].

Estrogens elicit their biological effects by binding to transcription factors called estrogen receptors (ERs) in the target organ/tissue (uterus, ovary, vagina, liver, bone, and breast) [11-13]. The ER exists as two subtypes, namely ER α and ER β [14]. Current estrogens in HRT activate both subtypes of ER in all tissues [14-19]. This attribute is beneficial in bone [18,20,21] and for hot flashes [18,21], but detrimental in the breast [6,21,22] and uterus [21,23] as it increases the risk of tumorigenesis. In contrast, the selective estrogen receptor modulators (SERMs), although not ER subtype specific [24,25], act as agonists in certain tissues, such as bone [26-28], and as antagonists in others, such as breast [9,10,29]. Although, the well-known SERMs, raloxifene and tamoxifen [30], have been shown to decrease the risk of breast cancer [18,31,32] and increase bone mineral density [26-28,33], they have also been linked to an increased risk of venous thromboembolism and

occurrence of hot flashes, and can stimulate endometrial growth [28,34–36]. SERMs are thus not considered as suitable alternatives for HRT.

Physiologically, while ER α is associated with the promotion of cell proliferation that contributes to the occurrence of breast and endometrial cancer, several studies have shown that ER β inhibits ER α -dependent cell proliferation and could prevent cancer development [15,22,37–43]. 17 β -estradiol (E₂) has similar binding affinities for the two ER subtypes [44], and the subtypes stimulate the transcription of both common and distinct subsets of E₂ target genes [13,17,39,45]. However, in many cases the degree of activation via ER β is lower [44], despite the high ligand independent transcriptional activity of this subtype [46,47]. In light of the above, it has been suggested that the development of ER subtype specific ligands may herald the arrival of a new generation of estrogen analogues that may present a novel treatment for post-menopausal symptoms, which in addition, may prevent or decrease the occurrence of breast cancer [44,48,49]. An ideal or “designer” estrogen analogue or selective estrogen receptor subtype modulator (SERSM) has been postulated that would have the following attributes: act as an ER α selective antagonist [50], down-regulate ER α protein levels [50,51], selectively activate ER β transcriptional pathways [15,19,24,43], and display anti-inflammatory properties by inhibiting transcription of pro-inflammatory genes to prevent the occurrence of post-menopausal osteoporosis [15,52]. Current examples of subtype specific ligands are, methyl-piperidino-pyrazole (MPP) (ER α antagonist) [53,54], diarylpropionitrile (DPN) (ER β agonist) [55], ERB-041 (ER β agonist) [56,57], liquiritigenin (ER β agonist) [19], isolated from the plant extract MF101 (ER β agonist) [24]. Phytoestrogens have been referred to as natural SERMs and can be both estrogenic as well as antiestrogenic [58–60]. Furthermore, although evidence in the literature shows that phytoestrogens can bind to both ER subtypes, they generally have a higher affinity for the ER β subtype [61–63] as well as a higher transcriptional potency and efficacy via ER β [63]. Despite conflicting evidence regarding doses of phytoestrogens and breast cancer risk [64,65], generally, findings have pointed the search in the direction of phytoestrogens and focused attention on phytoestrogen rich food sources as a possible source of the ideal SERSM.

One such source may be *Cyclopia* (family: Fabaceae), an indigenous fynbos plant from the Western Cape province of South Africa [66,67]. Traditionally, the “fermented” (oxidized) form of *Cyclopia*, has been consumed as a fragrant, caffeine free honeybush tea beverage with the “unfermented” form being introduced to the commercial market more recently [63,67,68]. Studies that investigated the chemical composition of *Cyclopia* have shown that phenolic compounds with estrogenic activity, for example luteolin, eriodictyol, naringenin, and formononetin, are present in various species of *Cyclopia* [63,68–72]. Furthermore, although dried methanol extracts (DMEs) from plant material of two species of *Cyclopia*, *C. genistoides* and *C. subternata*, have been shown to bind to the ERs and are able to transactivate an ERE-containing promoter reporter construct [62,63,68], only the extract from *C. genistoides* was investigated for ER subtype specificity and

found to transactivate only through ER β , despite binding to both subtypes [62,63]. In addition, studies by Verhoog et al. [63] and Mfenyana et al. [68] showed that although extracts of *Cyclopia* are able to induce proliferation of the ER α and ER β positive MCF-7 BUS cells, they antagonise E₂ induced cell proliferation.

The current study was prompted by the findings of Verhoog et al. [62,63] that the *Cyclopia* extract, P104, although binding to both receptors and with a much higher affinity for ER α , was able to activate an ERE-containing promoter reporter construct only via ER β . As the possibility of ER α antagonism by *Cyclopia* extracts had not been addressed in previous studies it appeared essential to evaluate ER α antagonism while also re-evaluating ER β agonism. The combination of ER α antagonism and ER β agonism may be especially relevant for the chemoprevention of breast cancer as ER antagonism serves as the basis of current chemo-preventative agents [29,31,32,73,74], while ER β specific agonists have recently been identified as having potential for the chemoprevention of breast cancer [19,22]. In addition, this combination might be advantageous for the treatment of menopausal symptoms as an ER β agonist has been shown to alleviate both hot flashes and the surge of inflammation related diseases during menopause [24,52], while an ER α antagonist would not result in hyperplasia of the uterus, commonly associated with ER α agonists [15,52]. Thus, in this study, we evaluate the potential of several extracts of *Cyclopia* to act as ER α antagonists and ER β agonists and demonstrate that all extracts display ER α antagonism, while two also display ER β agonism. In addition, all extracts antagonise E₂-induced MCF-7BUS cell proliferation, one extract displays anti-inflammatory activity, and the two tested extracts do not stimulate uterine growth. These results suggest that the *Cyclopia* extracts, which display ER α antagonism and ER β agonism, have positive attributes that could possibly be further exploited for the development of safer drugs for the treatment or prevention of osteoporosis or pre-menopausal symptoms.

Material and Methods

Ethics statement

Animal care and experimental procedures were conducted with strict adherence to the accepted standards for the use of animals in research and teaching as reflected in the South African National Standards 10386: 2008. Stellenbosch University ethics committee approved this study (ethical approval reference: 11NB_LOU01).

Test Compounds

17 β -Estradiol (E₂), genistein, luteolin, enterodiol, phorbol 12-myristate 13-acetate (PMA) and fulvestrant (ICI 182,780) were obtained from Sigma-Aldrich®, South Africa, and coumestrol was obtained from Fluka™ Analytical, Sigma-Aldrich®, South Africa. The *Cyclopia* extracts used for *in vitro* studies, P104 [62], SM6Met [68] and cup-of-tea [68], were previously prepared, while for *in vivo* studies new SM6Met and cup-of-tea extracts were prepared as previously described [68]. E₂,

genistein, luteolin, enterodiol, coumestrol and *Cyclopia* extract stock solutions were prepared in dimethylsulfoxide (DMSO).

High-performance liquid chromatography (HPLC) analysis of *C. subternata* extracts

The newly prepared SM6Met and cup-of-tea extracts were analyzed using HPLC. Extracts and stock solutions of standards were prepared in DMSO and aliquots frozen at -20°C until needed for analysis. For experimental analysis ascorbic acid was added to defrosted standards and extracts to a final concentration of 9.8 mg/ml. The mixtures were then filtered using Millex-HV syringe filters (Millipore) with a 0.22 μm pore size.

Analyses were performed on an Agilent 1200 HPLC consisting of an in line degasser, diode-array detection (DAD), column oven, autosampler and quaternary pump, controlled by Chemstation software (Agilent Technologies, Santa Clara, CA). The HPLC method previously described by De Beer et al. [75] was used to quantify the major phenolic compounds in *C. subternata* extracts: A Gemini-NX C18 (150 \times 4.6 mm; 3 μm ; 110 \AA) column was used in conjunction with 2% acetic acid (A) and acetonitrile (B) as mobile phases. Injection volumes ranged from 10–20 μl for standards and 5–50 μl for the extracts. Separation was performed at a flow rate of 1 ml/min with the following mobile phase gradient: 0–2 min (8% B), 2–27 min (8–38% B), 27–28 min (38–50% B), 28–29 min (50% B), 29–30 min (50–8% B), 30–40 min (8% B); at a temperature of 30°C .

The dihydrochalcones, flavanones and benzophenones were quantified at 288 nm, whereas the xanthenes, flavones and phenolic acids were quantified at 320 nm. A calibration curve consisting of seven points was set up for all the available standards (mangiferin (Sigma-Aldrich®, South Africa), isomangiferin (Chemos GmbH, Germany), luteolin (Extrasynthese, France), eriocitrin (Extrasynthese, France), hesperidin (Sigma-Aldrich®, South Africa), protocatechuic acid (Fluka™ Analytical, Sigma-Aldrich®, South Africa)) and also standards needed to calculate equivalent values (aspalathin (kind gift from Prof. Gelderblom, PROMEC unit, Medical Research Council, Tygerberg, South Africa), apigenin (Fluka™ Analytical, Sigma-Aldrich®, South Africa), and nothofagin (kind gift from Prof. Gelderblom, PROMEC unit, Medical Research Council, Tygerberg, South Africa)). Iriflophenone-3-C- β -glucoside and iriflophenone-di-O,C-hexoside was quantified using iriflophenone-3-C-glucoside isolated from *C. genistoides* (personal communication from Dr. D. de Beer). Scolymoside and vicerin-2 were expressed as luteolin and apigenin equivalents, respectively, as no authentic reference standards were available for these compounds. Also phloretin-3',5'-di-C-glucoside was expressed in terms of nothofagin (3-hydroxyphloretin-3'-C-glucoside) equivalents.

Cell Culture

COS-1, African green monkey kidney fibroblast cells (ATCC, United States of America), and MCF-7BUS human breast cancer cells [76] (a kind gift from A. Soto, Tufts University, Boston, Massachusetts, United States of America) were maintained in high glucose (4.5 g/L) Dulbecco's modified eagle's medium (DMEM) (Sigma-Aldrich®) supplemented with

10% FCS (Highveld Biologicals, South Africa), 100 IU/ml penicillin and 100 $\mu\text{g/ml}$ streptomycin (Gibco, Invitrogen™, South Africa), 2mM glutamine (Merck), 44mM sodium-bicarbonate (Gibco), 1mM sodiumpyruvate (Gibco, Invitrogen Corporation), and 0.1mM non-essential amino acids (Gibco). All cells were maintained in a humidified cell incubator, set at 97% relative humidity and 5% CO₂ at 37°C . For the cell proliferation assays (MTT assay) MCF-7BUS cells were withdrawn from 100 IU/ml penicillin and 100 $\mu\text{g/ml}$ streptomycin for seven days prior to use.

MTT assay

On day one MCF-7BUS cells were seeded into 96-well tissue culture plates at a concentration of 2500 cells/well and allowed 24 hours to settle. The next day cells were washed with 200 μl well pre-warmed PBS and the medium was changed to DMEM without phenol red supplemented with 5% charcoal treated FCS (Highveld Biologicals, South Africa) and incubated for 24 hours. After incubation the cells were treated for 48 hours with increasing concentrations test compounds and *Cyclopia* extracts in the presence or absence of 10^{-9}M E₂ where after the colorimetric MTT (3-(4,5-dimethylthiazolyl-2)-2,5-diphenyltetrazolium bromide) assay, adapted from Verhoog et al. [63] and Mfenyana et al. [68], was performed. Briefly, the MTT assay entails that 4 hours before the end of the incubation period the assay medium is changed to 150 μl DMEM without phenol red, but supplemented with 5% charcoal stripped FCS, and 50 μL of MTT (methylthiazolyl-diphenyl-tetrazolium) (Sigma-Aldrich®) solution (5 mg/ml) is added to each well. Cells are then incubated for four hours at 37°C , the medium removed, and 200 μL of solubilisation solution (DMSO) added to each well. The plate is then covered with foil, shaken at room temperature for 5 min, and the absorbance read at 550 nm on a BioTek® PowerWave 340 spectrophotometer. All assays included a negative solvent control, which consisted of 0.1% (v/v) DMSO only. Results are expressed as fold induction relative to solvent.

Promoter reporter studies

MCF-7BUS and COS-1 cells were seeded in sterile 10 cm tissue culture plates at a concentration of 2×10^6 cells/plate and allowed 24 hours to settle. On day two the cells were rinsed once with sterile phosphate buffered saline (PBS) (pre-warmed to 37°C), medium changed to DMEM without phenol red supplemented with 10% charcoal treated FCS and 1% penicillin and streptomycin mixture, and cells were transfected.

Plasmids. Human (h) ER α (pSG5-hER α [77]) and ER β (pSG5-hER β [78]) expression plasmids were kind gifts from F. Gannon (European Molecular Biology Laboratory, Heidelberg, Germany). The ERE-containing promoter reporter construct (ERE.vit2.luc) was a kind gift from K. Korach, National Institute of Environmental Health Science, U.S. [79] and the NF κ B-containing promoter reporter construct (p(IL6kB)350hu.lL6Pluc + [80]) was a kind gift from G. Haegeman, University of Ghent, Ghent, Belgium. pGL2-Basic (Promega Corporation, Madison, Wisconsin, U.S.A.) was used as an empty vector.

Transactivation. To test transactivation through ER α COS-1 cells were transfected with 150 ng hER α and 6000 ng

of an ERE-containing promoter reporter construct. To test transactivation through ER β COS-1 cells were transfected with 150 ng hER β , 3000 ng of an ERE-containing promoter reporter construct, and 3000 ng empty vector. MCF-7 BUS cells (which contain endogenous hER α and hER β) were transfected with 3000 ng of an ERE-containing promoter reporter construct and 3000 ng empty vector. The amount of promoter reporter construct for each test model that was selected was determined by the highest E₂ induction achieved (Figure S1).

Transrepression. To test transrepression through ER α COS-1 cells were transfected with 150 ng hER α , 1500 ng of an NF κ B-containing promoter reporter construct and 4500 ng empty vector. To test transrepression through ER β COS-1 cells were transfected with 150 ng hER β , 4500 ng of an NF κ B-containing promoter reporter construct and 1500 ng empty vector. MCF-7BUS cells (which contain endogenous hER α and hER β) were transfected with 6000 ng of an NF κ B-containing promoter reporter construct. The amount of promoter reporter construct for each test model that was selected was determined by the most effective E₂ repression of PMA induction achieved (Figure S2)

All transfections were performed using FuGENE™ 6 transfection reagent (Roche Applied Science, South Africa) as described by the manufacturer. Cells were left for 24 hours, replated in sterile 24-well tissue culture plates at a concentration of 5 x 10⁴ cells/well and allowed 24 hours to settle. Cells were treated for 24 hours with test compounds and *Cyclopia* extracts and lysed overnight with 50 μ l lysis buffer [0.2% (vol/vol) Triton, 10% (vol/vol) glycerol, 2.8% (vol/vol) Tris-phosphate-EDTA, and 1.44 mM EDTA] per well at -20 °C. Luciferase activity was determined using the luciferase assay kit (Promega Corporation, Anatech, South Africa) according to the manufacturer's instructions and normalized for protein content (Bradford assay [81]). Results are expressed as fold induction relative to solvent.

Western Blot

Cell lysates from COS-1 cells transfected with either ER α (150 ng hER α /10 cm plate) or ER β (150 ng hER β /10 cm plate) and MCF-7BUS cells were prepared by adding lysis buffer A (10mM Hepes pH 7.5 (Gibco, Invitrogen Corporation), 1.5mM MgCl₂, 10mM KCl, 0.1% NP-40 (Roche Applied Science) and Complete Mini protease inhibitor cocktail (Roche Applied Science), shaking on ice for 15 min and frozen overnight at -20°C.

On thawing, lysate were transferred to 1.5ml Eppendorf tubes on ice, centrifuged for 10 min at 12 000 x g at 4°C and the cleared lysate was transferred to 1.5ml Eppendorf tubes on ice, aliquoted and stored at -20°C until assayed. Lysates (20 μ l) were separated on a 10% SDS-PAGE gel. Following electrophoresis, proteins were electro-blotted and transferred to a Hybond-ECL nitrocellulose membrane (Amersham Biosciences, South Africa), which was probed for ER α (diluted 1:500), ER β (1:250) and GAPDH (1:500). Proteins were visualized using HRP labeled anti-rabbit antibody for ER α (1:2500) and ER β (1:1000), or HRP labeled anti-mouse antibody for GAPDH (1:5000), and ECL Western blotting detection reagents (Pierce®, Thermo Fisher Scientific Inc.,

U.S.A.) and medical x-ray film (Axim (PTY) LTD., South Africa). All antibodies, primary [ER α (HC-20), cat# sc-543, ER β (H-150), cat# sc-8974, and GAPDH (0411), cat# sc-47724] and secondary (anti-rabbit, cat# sc-2005, and anti-mouse, cat# sc-2030), were purchased from Santa Cruz Biotechnology, Inc., U.S.A.

Animal care

Immature female Wistar rats were obtained from the Stellenbosch University, South Africa, breeding unit and were received as weanlings on postnatal day 18. The animals had free access to standard rat feed (Pure Harvest Rat Feed, Afresh Vention (PTY) Ltd, South Africa) and drinking water. The animals were housed in a 12 hour light-dark cycle at a constant temperature of 20 °C in EHRET individually ventilated cages (EHRET, Emmendingen, Germany). The animals were allowed at least 24 hours to acclimatize before the onset of experimental procedures.

Immature rat uterotrophic assay

The immature rat uterotrophic assay was performed according to methods previously described by Kanno et al. [82] and de Lima et al. [83]. Immature Wistar rats (21 days) were randomly assigned to groups (n=10) and treated daily with E₂, genistein, *Cyclopia* extracts, or vehicle control (sterile PBS) by oral gavage for three consecutive days. The dose volume was 200 μ l/day. The test compounds were suspended in sterile PBS and the solution was kept homogenous by stirring before administration. General health, vaginal opening, and body weight was monitored daily and recorded. On day four, approximately 24 hours after last dose, animals were weighed and sacrificed by administration of a high dose of Isoflurane (2-chloro-2-(difluoromethoxy)-1,1,1-trifluoro-ethane), (Safeline pharmaceuticals Pty (Ltd)). Livers were removed and weighed. Uteri were removed, cleaned of excess fat, photographed, weighed, pierced to remove luminal fluids, and blotted uterine weights were obtained immediately.

Evaluation/Monitoring of vaginal opening of Wistar rats for extended period

Immature Wistar rats (21 days) were randomly assigned to groups (n=10) and treated daily with E₂, *Cyclopia* extracts, or vehicle control (sterile PBS) by oral gavage for 30 consecutive days. The dose volume had to be increased gradually from 200 μ l/day to 400 μ l/day as animals increased in body weight. The test compounds were suspended in sterile PBS and the solution was kept homogenous by stirring before administration. General health, vaginal opening, and body weight was monitored daily and recorded. On day 30 animals were weighed and sacrificed by administration of a high dose of Isoflurane.

Data manipulation and statistical analysis

The GraphPad Prism® version 5.10 for Windows (GraphPad Software) was used for graphical representations and statistical analysis. One-way ANOVA and Dunnett's post-test comparing all columns to the solvent control were used for statistical

analysis and significance is displayed on the graphs. For all experiments the error bars represent the SEM of at least two independent experimental repeats, except for *in vivo* studies where the error bars represent the SEM of the number of animals used.

Results

HPLC analyses of extracts of *Cyclopia*

New SM6Met and cup-of-tea extracts were prepared from the same harvesting of *C. subternata* previously used to prepare these extracts [68]. HPLC analysis was performed on these newly prepared SM6Met and cup-of-tea extracts (Table 1). Prior HPLC results of previously prepared P104 [63] and SM6Met [68] extracts are also shown in Table 1. The results indicate the presence of the xanthenes, mangiferin and isomangiferin, the flavones, scolymoside, luteolin, and vicenin-2, the flavanones, eriocitin and hesperidin, the dihydrochalcones, phloretin-3,5-di-C-glucoside and aspalathin, the benzophenones, iriflophenone-3-C- β -glucoside and iriflophenone-di-O,C-hexoside, and the phenolic carboxylic acid, protocatechuic acid. P104, a DME from *C. genistoides*, contained more mangiferin and isomangiferin than SM6Met, a DME from *C. subternata*, while the cup-of-tea extract from the same species contained the least. Luteolin was present in all of the extracts, albeit at small amounts, with the P104 extract containing the largest amount, followed by the SM6Met extracts, and with the cup-of-tea extract containing the least. The luteolin rutinoside, scolymoside, was not evaluated in P104. The DMEs of *C. subternata* contained more scolymoside, eriocitrin, hesperidin, and phloretin-3,5-di-C-glucoside than the cup-of-tea extract. The newly prepared DME, SM6Met, contained higher amounts than the cup-of-tea extract of compounds not previously tested for, namely, iriflophenone-3-C- β -glucoside, iriflophenone-di-O,C-hexoside, aspalathin, vicenin-2, and protocatechuic acid. In general the DMEs contained higher concentrations of the polyphenols quantified (Table 1) than the water extract.

Methanol extracts of *Cyclopia* act as agonists of ER β , while all extracts antagonize E₂-induced activation via ER α

To evaluate ER α antagonism while also re-evaluating ER β agonism COS-1 cells were transiently transfected with either ER α (Figures 1 A, C) or ER β (Figures 1 B, D) and an ERE-containing promoter reporter construct. Agonism was tested in the absence (Figures 1 A, B) and antagonism in the presence (Figures 1 C, D) of 10⁻⁹ M E₂. Three *Cyclopia* extracts, from two species, *C. genistoides* and *C. subternata*, were tested. Two were methanol extracts, P104 and SM6Met, and one was a water extract, cup-of-tea. In addition we investigated an example from each of the major classes of phytoestrogens: genistein, a well-studied isoflavone, enterodiol, a lignin, and coumestrol, a coumestan [84,85]. Luteolin, an estrogenic polyphenol [71], was also included as it was found to be present in all of the *Cyclopia* extracts (Table 1), while E₂ represents the major endogenous estrogen [86,87].

Table 1. Major polyphenols present in previously and newly prepared extracts of *Cyclopia* as determined by HPLC.

Polyphenol [% of dry extract (g/100g dry extract)]	Extract			
	Previously prepared	Newly prepared		
	P104 [63]	SM6Met [68]	SM6Met	Cup-of-tea
	<i>C.genistoides</i>	<i>C.subternata</i>	<i>C.subternata</i>	<i>C.subternata</i>
Mangiferin	3.606	1.850	1.899	1.000
Isomangiferin	5.094	0.750	0.645	0.420
Luteolin	0.096	0.040	0.040	0.018
Scolymoside (luteolin-7-O-rutinoside)	nt ^a	1.820 ^c	1.289	0.876
Vicenin-2 (apigenin-6,8-di-C-glucoside)	nt	nt	0.089	0.065
Eriocitrin (eriodictyol-7-O-rutinoside)	nd ^b	1.250	0.846	0.600
Hesperidin (hesperitin-7-O-rutinoside)	nt	1.870	2.049	0.935
Phloretin-3,5-di-C-glucoside	nt	1.270 ^d	1.278	0.939
Aspalathin (3-hydroxyphloretin-3',5'-di-C-hexoside)	nt	nt	0.700	0.582
Iriflophenone-3-C- β -glucoside	nt	nt	0.669	0.590
Iriflophenone-di-O,C-hexoside	nt	nt	0.958	0.896
Protocatechuic acid	nt	nt	0.113	0.082

^aNot tested

^bNot detected

^cPreviously "Unknown 1" was quantified as luteolin equivalents as it appeared to be a flavone.

^dPreviously "Unknown 2" was quantified as hesperidin equivalents as it appeared to be a flavanone.

doi: 10.1371/journal.pone.0079223.t001

E₂ induced ER α mediated transactivation in a dose dependent manner with significant induction at two concentrations of E₂, 10⁻⁹ M (2.7 x 10⁻⁴ μ g/ml) (2.5 \pm 0.5 fold) and 9.8 μ g/ml (3.6 x 10⁻⁵ M) (3.9 \pm 0.7 fold), but not at the lowest concentration of 10⁻¹¹ M (2.7 x 10⁻⁶ μ g/ml) (Figure 1A). The same trend was seen for ER β (2.5 \pm 0.5 fold at 10⁻⁹ M and 2.7 \pm 0.4 fold at 9.8 μ g/ml) (Figure 1B), although at the highest concentration of E₂ higher induction was observed via ER α than via ER β (3.9 \pm 0.7 vs. 2.7 \pm 0.4 fold). Although the 9.8 μ g/ml E₂ represents a supra-physiological concentration the 10⁻¹¹ M and 10⁻⁹ M E₂ concentrations reflect the pre- and post-menopausal levels of E₂ respectively [88]. At the concentration of 9.8 μ g/ml, genistein (3.6 x 10⁻⁵ M), luteolin (3.4 x 10⁻⁵ M), and coumestrol (3.7 x 10⁻⁵ M) significantly activated gene transcription through both of the ER subtypes (Figures 1A, B).




Figure 1. Evaluation of ER subtype specific agonism and antagonism of transactivation of an ERE-containing promoter reporter construct in COS-1 cells. COS-1 cells were transfected with either (A and C) pSG5-hER α or (B and D) pSG5-hER β and ERE.vit2.luc and treated for 24 hours with a series of test compounds or extracts. To test agonism cells were treated with test compounds or extracts alone, (A and B), while, to test for antagonism cells were treated with test compounds or extracts in the presence of 10⁻⁹ M E₂ (C and D). Statistical analysis was done using One-way ANOVA with Dunnett's post-test comparing all columns to the solvent control (*, P<0.05; **, P<0.01; ***, P<0.001). The dotted line through the bars represents the values for solvent control. Average \pm SEM is of four independent experiments done in triplicate.

doi: 10.1371/journal.pone.0079223.g001

Enterodiol, however, could not significantly activate gene transcription through either of the subtypes at the concentration of 9.8 μ g/ml (3.2 \times 10⁻⁵ M) (Figures 1A, B). None of the *Cyclopia* extracts were able to induce activation through ER α (Figure 1A), but both the methanol extracts, P104 and

SM6Met, were able to significantly activate transcription through ER β (2.4 \pm 0.4 and 2.5 \pm 0.3 fold, respectively).

To address antagonism, transactivation in the presence of 10⁻⁹ M E₂ was evaluated (Figure 1C and D). The phenolic compounds, genistein, luteolin, and coumestrol were not

antagonists but had an additive effect on E₂-induced activation via both receptor subtypes (Figures 1C and D), confirming their agonism through both subtypes (Figure 1A and B). Enterodiol in contrast, however, only displays ER α antagonism (0.7 \pm 0.1 fold vs. E₂ activation set as 1) (Figure 1C). All of the *Cyclopia* extracts significantly antagonized ER α mediated E₂-induction (P104, 0.7 \pm 0.1, SM6Met, 0.7 \pm 0.1, and cup-of-tea, 0.6 \pm 0.1 fold), however, only P104 had an additive effect on the E₂-induced activation through ER β (Figure 1D). In conclusion, the methanol extracts of *Cyclopia* are ER β agonists and all extracts are ER α antagonists.

In MCF-7BUS cells expressing both ER subtypes all extracts of *Cyclopia* transactivate an ERE-driven promoter reporter construct

Most tissues affected by menopause and/or implicated in HRT side effects, such as uterus, bone, and breast, contain both ER subtypes [89]. Thus, having shown that methanol extracts of *Cyclopia* are ER β agonists and all extracts are ER α antagonists in a system where the ER subtypes were evaluated separately, we were interested in investigating the transactivation potential of *Cyclopia* extracts in a system where both subtypes are present.

MCF-7BUS cells, containing both ER α and ER β (Figure 2A), were transfected with an ERE-containing promoter reporter construct and both agonism (Figure 2B) and antagonism (Figure 2C) were tested. Although strong transactivation was seen with E₂, none of the polyphenols on their own were able to significantly activate gene transcription in this system where both ER subtypes are present (Figure 2B), despite the fact that these polyphenols transactivate when the ER subtypes function in isolation (Figure 1A and B). Furthermore, most of the polyphenols, excluding coumestrol, antagonized E₂ induction when both ER subtypes are together (Figure 2C), whereas when the subtypes were expressed separately only enterodiol showed ER α antagonism (Figure 1C). In contrast to the polyphenols, the extracts of *Cyclopia*, P104 (3.4 \pm 0.5 fold), cup-of-tea (3.4 \pm 0.5 fold) and, SM6Met (3.5 \pm 0.6 fold), were able to activate transcription to a similar extent as 10⁻⁹ M E₂ (3.8 \pm 0.3 fold) (Figure 2B). These results, together with the fact that the *Cyclopia* extracts did not antagonize E₂ induction (Figure 2C), suggests that when both ER subtypes are co-expressed the *Cyclopia* extracts act as agonists, whereas when the ER subtypes are expressed separately they only act as agonists through ER β and antagonize ER α induction.

An extract of *C. genistoides* represses NF κ B activation via ER α and ER β whereas the extracts of *C. subternata* are ER β antagonists.

The decline in estrogen levels during menopause leads to a surge in the occurrence of inflammatory disorders [52,90-92]. Furthermore, NF κ B, a pro-inflammatory transcription factor, is involved in the development of breast cancer [93-95]. Taking this into account we wanted to evaluate the ability of *Cyclopia* extracts to repress the activation of an NF κ B-containing promoter reporter construct by transfecting COS-1 cells with said construct and either ER α (Figures 3A, C, E) or ER β (Figures 3B, D, F). In addition, this system would provide information concerning the behavior of *Cyclopia* extracts in a

transrepression model. Agonism was tested in the absence (Figures 3A, B) and antagonism (Figures 3C, D) in the presence of 10⁻⁹ M E₂.

PMA (phorbol 12-myristate 13-acetate, an activator of NF κ B driven gene expression [96,97]) activation of the NF κ B-containing construct was repressed by E₂ via both receptor subtypes (Figure 3A and B) with a more pronounced repression through ER α (38.6% vs. 27.2%). Like E₂, all of the polyphenols, as well as P104 (*C. genistoides* extract), acted as ER α agonists by repressing PMA activation (genistein 52.1%, luteolin 50.6%, enterodiol 57.4%, coumestrol 61.8%, and P104 59.2%) (Figure 3A). Furthermore, genistein (34.8% repression) and P104 (40.7% repression), like E₂, also displayed significant ER β agonism (Figure 3B). Therefore, in our transrepression model P104 is not an ER β selective agonist, but displays agonism via both subtypes. The water extract of *C. subternata*, cup-of-tea, was unable to repress PMA induction through either ER α or ER β (Figures 3A, B) while the methanol extract, SM6Met, also unable to repress PMA induction through either subtype, significantly added to the activation observed with PMA alone via ER α (5.1 \pm 0.5 vs. 3.5 \pm 0.5) (Figure 3A, B).

Antagonism was evaluated in the presence of 10⁻⁹ M E₂ and only genistein (Figure 3C) had a significant effect via ER α by antagonizing E₂ repression of PMA activation. The polyphenols, luteolin, enterodiol, and coumestrol, but not genistein, however, antagonized E₂ repression of PMA activation via ER β (Figure 3D). Although none of the extracts displayed significant antagonism of ER α , the extracts of *C. subternata* displayed ER β antagonism (Figure 3D).

The result for SM6Met in Figure 3A prompted us to investigate whether this effect was via ER α or if SM6Met is able to activate the NF κ B-containing construct through another mechanism of action. Therefore, we repeated the experiment, for both receptor subtypes, with SM6Met, as well as P104, in the presence and absence of an ER antagonist, ICI 162,780 (Figures 3E, F). The observed repression of PMA activation by E₂ and P104 via ER α and ER β is abolished by ICI (Figure 3E, F) and thus, the observed repression is indeed via the ER. SM6Met, like ICI, increases PMA activation through ER α (Figure 3E) and both have no significant effect on PMA activation via ER β (Figure 3F). Furthermore, the increased transactivation observed with SM6Met in Figure 3A may be attributed to residual E₂ remaining after stripping of FCS, as suggested by others [22], which would further support the contention that SM6Met is behaving as an ER α antagonist. In conclusion then the results suggest that for our transrepression model the methanol extract of *C. genistoides* (P104) is behaving like an ER α and ER β agonist, while the methanol extract of *C. subternata* (SM6Met) is an ER α antagonist in the absence of E₂, and an ER β antagonist in the presence of E₂.

In MCF-7BUS cells expressing both ER subtypes all extracts are agonists, while the water extract of *C. subternata* also displays antagonistic activity.

As we have shown that P104 is an ER agonist and SM6Met is an ER antagonist in a transrepression model where the ER subtypes function in isolation (Figure 3), we wanted to test the effect of these extracts in a model where both subtypes are




Figure 2. Evaluation of transactivation of an ERE-containing promoter reporter construct in MCF-7BUS cells expressing both ER α and ER β . MCF-7BUS cells, with endogenous ER α and ER β (A), were transfected with ERE.vit2.luc and treated for 24 hours with a series of test compounds or extracts. To test agonism cells were treated with test compounds or extracts alone (B), while, to test for antagonism cells were treated with test compounds or extracts in the presence of 10⁻⁹M E₂ (C). Statistical analysis was done using One-way ANOVA with Dunnett's post-test comparing all columns to the solvent control (*, P<0.05; **, P<0.01; ***, P<0.001). The dotted line through the bars represents the values for solvent control. Average \pm SEM is of three independent experiments done in triplicate.

doi: 10.1371/journal.pone.0079223.g002

present as most tissues affected by menopause and/or implicated in HRT side effects contain both subtypes.

MCF-7BUS cells were transfected with an NF κ B-containing promoter reporter construct and both agonism (Figure 4A) and antagonism (Figure 4B) evaluated. Strong repression was observed with E₂, the polyphenols, and P104 when both subtypes are present (Figure 4A), which correlates with what was observed previously for ER α alone (Figure 3A). However, for ER β alone (Figure 3B), significant repression was previously seen only with E₂, genistein, and P104 but not with luteolin, enterodiol, and coumestrol. Unlike previous results, SM6Met behaved differently when subtypes were co-expressed than when the subtypes were expressed separately. It displayed agonism when subtypes are expressed together (Figure 4A) while displaying antagonism when expressed separately (Figure 3A and D). Similarly, where no agonist activity via either subtype alone was observed previously, the cup-of-tea extract was able to change its behavior when both subtypes are present by displaying ER agonism. Furthermore,

antagonism in the presence of both subtypes was only seen with the cup-of-tea extract (Figure 4B), while the subtype specific antagonism of genistein, luteolin, enterodiol, coumestrol, and SM6Met (Figures 3C, D) is abrogated in the presence of both subtypes. Taken together, in a transrepression model, the DME of *C. genistoides*, P104, is an ER agonist in all models (Figures 3A, B, and 4A), the DME of *C. subternata*, SM6Met, is an ER β antagonist in the presence of E₂ (Figure 3D), an ER α antagonist in the absence of E₂ (Figure 3A, E), and an agonist in the presence of both ER subtypes (Figure 4A), while the water extract of *C. subternata*, cup-of-tea, is an ER β antagonist (Figure 3D) and an ER agonist/antagonist (Figures 4A, B) in the presence of both subtypes. This differential behavior of the *Cyclopia* extracts in the transrepression model contrasts to similar behavior by the extracts in the transactivation model where all extracts displayed antagonism through ER α (Figure 1) alone, while displaying agonism to ER β (Figure 1) alone or when both subtypes are expressed (Figure 2).




Figure 3. Evaluation of ER subtype specific agonism and antagonism of an NF κ B-containing promoter reporter construct in COS-1 cells. COS-1 cells were transfected with either (A, C, and E) pSG5-hER α or (B, D, and F) pSG5-hER β and p(IL6kB)350hu.II6Pluc+ and treated for 24 hours with a series of test compounds or extracts. To test agonism cells were treated with test compounds or extracts alone, (A and B), while, to test antagonism cells were treated with test compounds or extracts in the presence of 10⁻⁹ M E₂ (C and D). To ascribe the observed effect to the ER we treated cells with P104 and SM6Met in the absence or presence of the ER antagonist ICI 182,870 (E and F). Statistical analysis was done using One-way ANOVA with Dunnett's post-test comparing all columns to either (A, B, E, and F) 10ng/ml PMA or (C and D) 10ng/ml PMA + 10⁻⁹ M E₂ (*, P<0.05; **, P<0.01; ***, P<0.001). The dotted lines through the bars represent the values for either (A, B, E, and F) solvent control, 10ng/ml PMA, or 10ng/ml PMA + 10⁻⁹ M E₂ or (C and D) 10ng/ml PMA + 10⁻⁹ M E₂. Average \pm SEM is of three independent experiments done in triplicate.

doi: 10.1371/journal.pone.0079223.g003




Figure 4. Evaluation of transrepression of an NF κ B-containing promoter reporter construct in MCF-7BUS cells expressing both ER α and ER β . MCF-7BUS cells were transfected with p(IL6kB)350hu.LL6Pluc+ and treated for 24 hours with a series of test compounds or extracts. To test agonism cells were treated with test compounds or extracts alone, (A), while, to test for antagonism cells were treated with test compounds or extracts in the presence of 10⁻⁹M E₂ (B). Statistical analysis was done using One-way ANOVA with Dunnett's post-test comparing all columns to either (A) 10ng/ml PMA or (B) 10ng/ml PMA + 10⁻⁹M E₂ (*, P<0.05; **, P<0.01; ***, P<0.001). The dotted lines through the bars represent the values for either (A) solvent control, 10ng/ml PMA, or 10ng/ml PMA + 10⁻⁹M E₂ or (B) 10ng/ml PMA + 10⁻⁹M E₂. Average \pm SEM is of three independent experiments done in triplicate.

doi: 10.1371/journal.pone.0079223.g004

Cyclopia extracts weakly induce proliferation of breast cancer cells but antagonizes E₂-induced breast cancer cell proliferation

Having shown that *Cyclopia* extracts can modulate both transactivation and transrepression in the presence of both ER subtypes and when the subtypes are expressed alone, we wanted to re-evaluate agonism of P104 [63], SM6Met and cup-of-tea [68] (Figure 5) and antagonism of P104 [63] (Figure 6) and for the first time evaluate antagonism of SM6Met and cup-of-tea (Figure 6) on MCF-7BUS breast cancer cell proliferation. Cell proliferation in MCF-7BUS cells constitutes an integrated model where not only the ER subtypes are co-expressed, but both transactivation and transrepression of endogenous genes contribute towards the final phenotype, whether it is proliferative or anti-proliferative [39,98-100].

The MTT cell proliferation assay using MCF-7BUS cells was used to address agonism (Figure 5A-H). Estrogen induced cell proliferation at a wide range of concentrations (10⁻⁶ M to 10⁻¹⁰ M) with the highest efficacy (2.1 \pm 0.1 fold) observed at 10⁻⁹ M E₂ (2.7 \times 10⁻⁴ μ g/ml) (Figure 5A). Like E₂, all of the polyphenols were also able to induce cell proliferation, but not to the same extent as E₂, with a maximum efficacy of: genistein, 1.5 \pm 0.1 fold at 10⁻⁹ M (2.7 \times 10⁻⁴ μ g/ml) (Figure 5B), luteolin, 1.5 \pm 0.1 fold at 10⁻⁵ M (2.7 μ g/ml) (Figure 5C), coumestrol, 1.6 \pm 0.1 fold at 10⁻⁶ M (3.0 \times 10⁻¹ μ g/ml) (Figure 5D), and enterodiol, 1.3 \pm 0.1 fold at 10⁻⁹ M (3.0 \times 10⁻⁴ μ g/ml) (Figure 5E). Similarly, all

three extracts of *Cyclopia* induced proliferation of cells with a lower efficacy than E₂ with maximum efficacies of 1.5 \pm 0.2 (significantly different from E₂), 1.3 \pm 0.03 (significantly different from E₂), and 1.7 \pm 0.2 (not significantly different from E₂) fold for 9.8 μ g/ml of P104, cup-of-tea and SM6Met, respectively (Figures 5F-H). The potencies, depicted by EC₅₀ values on graphs (Figures 5A-H), of the polyphenols, as well as of the *Cyclopia* extracts, were lower than that of E₂ with coumestrol, P104, and SM6Met significantly lower and may be listed in order of decreasing potency as follow: E₂ > genistein > enterodiol > luteolin > cup-of-tea > P104 > coumestrol >> SM6Met.

To address antagonism (Figure 6A-G), increasing concentrations of the polyphenols and *Cyclopia* extracts were tested in the presence of 10⁻⁹ M E₂ (highest efficacy, Figure 5A). Genistein (Figure 6A) and enterodiol (Figure 6D), significantly repressed E₂-induced cell proliferation (23.3% at 10⁻⁵ M (2.70 μ g/ml) and 24.5% at 10⁻⁵ M (3.02 μ g/ml), respectively). Although, luteolin (Figure 6B) and coumestrol (Figure 6C) displayed no significant antagonism, coumestrol did have a significant additive effect (1.3 \pm 0.1 fold) at 10⁻⁹ M (2.96 \times 10⁻⁴ μ g/ml), suggesting agonism. Similarly, genistein, an antagonist at high concentrations, also had a significant additive effect (1.2 \pm 0.1 fold) at the lower concentration of 10⁻⁹ M (2.70 \times 10⁻³ μ g/ml) (Figure 6A). All extracts of *Cyclopia* were able to antagonize E₂-induced cell proliferation, with P104 repressing 19.8% at 9.8 \times 10⁻¹ μ g/ml, SM6Met 16.8% 9.8 \times 10⁻⁴




Figure 5. Evaluation of agonism of proliferation, a more complex endpoint encompassing both transactivation and transrepression in MCF-7BUS cells expressing both ER α and ER β . MCF-7 BUS cells were treated with increasing concentrations of (A) E_2 , (B-E) polyphenols, and (F-H) *Cyclopia* extracts for 48 hours. After treatment the amount of living cells was determined using a MTT assay. Statistical analysis was done using One-way ANOVA with Dunnett's post-test comparing all columns to the solvent control (*, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$) or to E_2 for EC_{50} values (#, $P < 0.05$; ##, $P < 0.01$; ###, $P < 0.001$). The dotted line through the bars represents the values for solvent control. Average \pm SEM is of two independent experiments done in six replicates, except (A) where average \pm SEM is of three independent experiments done in six replicates.

doi: 10.1371/journal.pone.0079223.g005




Figure 6. Evaluation of antagonism of proliferation, a more complex endpoint encompassing both transactivation and transrepression in MCF-7BUS cells expressing both ER α and ER β . MCF-7 BUS cells were treated with increasing concentrations of (A-D) polyphenols and (E-G) *Cyclopia* extracts for 48 hours in the presence of 10⁻⁹M E₂. After treatment the amount of living cells was determined using a MTT assay. Statistical analysis was done using One-way ANOVA with Dunnett's post-test comparing all columns to the solvent control (*, P<0.05; **, P<0.01; ***, P<0.001). The dotted line through the bars represents the values for solvent control. Average \pm SEM is of two independent experiments done in six replicates.

doi: 10.1371/journal.pone.0079223.g006

$\mu\text{g/ml}$, and cup-of-tea 15.6% repression at $9.8 \times 10^{-4} \mu\text{g/ml}$ (Figures 6E, F, G). Taken together, these results show that although all extracts of *Cyclopia* induced cell proliferation, the P104 and cup-of-tea extracts did so at a significantly lower efficacy and the methanol extracts at a significantly lower potency than E₂, and that all extracts could antagonize E₂-induced cell proliferation.

SM6Met does not stimulate the growth of rat uteri, antagonizes E₂-induced uterine proliferation, and delays vaginal opening

For the *in vivo* studies only extracts from *C. subternata* was used as P104 plant material was not available in bulk. The immature rat uterotrophic assay is used to determine the ability of test compounds to stimulate ER α induced uterine growth as ER β is not highly expressed in the uterus [56,101] and also allows for the detection of antiestrogens [102]. Rats were administered E₂, genistein, and the two *C. subternata* extracts, SM6Met and cup-of-tea, via oral gavage and the effects on uterine growth were evaluated (Figure 7A, B, and Figure S3). Estrogen, as well as genistein, induced uterine growth (2.5 ± 0.2 and 2.0 ± 0.2 fold, respectively) (Figure 7). In contrast, the extracts significantly reduced uterine weight relative to solvent (Figure 7 and Figure S3). SM6Met also significantly repressed E₂-induced uterine growth by 33.0%, a result that is similar, but less pronounced, than that seen with ICI 182,780 (59.7% repression) (Figure 7) suggesting that the extracts behave as antiestrogens in the uterus.

We also addressed body weight changes and toxicity (Figure S4) and found that E₂ significantly increased body weight, whereas genistein significantly decreased body weight. The extracts of *Cyclopia* and ICI 182,780, however, did not lead to significant weight gain or loss as compared to solvent, except for the animals treated with the highest concentrations (2000mg/kg BW) of SM6Met and cup-of-tea extracts which gained significantly less weight than the solvent treated animals. With regards to toxicity, none of the treated animals showed any significant changes in liver weight, except for a decrease in liver weight in animals treated with 200mg/kg BW SM6Met.

Furthermore, as another marker of estrogenic activity, albeit a less sensitive marker [102], we also evaluated time of vaginal opening over an extended period of daily treatments (Figure 8). Estrogen led to premature vaginal opening when compared to solvent (4.2 ± 0.4 vs. 14.3 ± 1.2 days). This correlates with the observed increase in uterine weight in Figure 7. The significantly delayed vaginal opening of SM6Met treated animals (19.0 ± 1.3 days) also correlates with uterine weight results in displaying antiestrogenic behavior. The significant delay in vaginal opening was observed for all three of the concentrations of SM6Met, however, although the cup-of-tea extract showed a similar trend, it was not significant (Figure S5).

To summarize, for the first time we show that the *C. subternata* extracts are absorbed when administered orally and elicit a biological effect *in vivo*. Specifically, *Cyclopia* extracts, in contrast to E₂ and genistein, did not induce uterine growth and SM6Met antagonized E₂-induced uterine growth.

Furthermore, the extracts also delayed vaginal opening in contrast to E₂. These results suggest that the *Cyclopia* extracts display ER α antagonism *in vivo* by retarding uterine growth [56,101].

Discussion

HRT in the form of estrogens provides relief from the plethora of menopause associated symptoms [1]. Although these estrogens provide relief from menopausal symptoms, they introduced new HRT associated risks, including an increased occurrence of breast cancer, heart disease, strokes, and endometrial cancer [1,5,6,8]. These risks, and the associated reluctance of usage, instigated the search for a new generation of estrogen analogues that would provide the benefits of estrogens without the associated risks. In addition, it would be of great value if these new analogues display chemo-preventative properties in breast and endometrial tissues [9,10,29].

The search for new estrogen analogues heralded the era of the SERMs. These SERMs would selectively modulate estrogen receptors in different tissues, acting as antagonists in the breast and uterus (chemo-preventative) and as agonists in the bone (osteoporosis prevention). Tamoxifen, a first generation SERM, provided the desired protective effect in the breast [31,32] and raloxifene, a second generation SERM, had protective properties in breast and bone tissues [26,27,103]. However, as these SERMs have been linked to the increased occurrence of hot flashes and stimulated endometrial growth (tamoxifen), the search continues [28,34,35]. Third generation SERMs, such as lasoxifene and bazedoxifene, are currently in development, but the focus has shifted to osteoporosis treatment with protection against breast cancer as a beneficial side effect [104-106].

Although SERM development continues there is increased interest in SERSMs, analogues that can differentially modulate specific ER subtypes. This was brought on by studies that have shown that ER β inhibits ER α dependent cell proliferation and could prevent cancer development [15,22,37,40-43]. Phytoestrogens have been shown to be both estrogenic as well as antiestrogenic and while they can bind to both ER subtypes, they generally have a higher affinity for ER β as well as a higher transcriptional potency and efficacy via ER β [61-63]. Thus, phytoestrogen rich food sources have become important potential resources of SERSMS.

The current study evaluated previously described extracts of *Cyclopia*, a source of phytoestrogens, for ER agonism and/or antagonism (summarized in Table S1). Specifically, we evaluated the effect of *Cyclopia* extracts on transactivation and transrepression in a model where ER α and ER β were expressed separately. This allows for the evaluation of the modulation of ER subtype specific activity in two transcriptional models: a classical ERE transactivation model and an NF κ B transrepression model. In the transactivation model the methanol extracts, P104 and SM6Met were ER β agonists, while all extracts antagonized ER α . In the transrepression model, however, the behavior of the *Cyclopia* extracts became more complex. P104, which displayed opposite effects via the




Figure 7. Evaluation of the *in vivo* effect of E₂, genistein and SM6Met on immature rat uterine growth. Immature female wistar rats were treated with 100µg/kg body weight E₂, in the presence and absence of 2000mg/kg body weight SM6Met or 10mg/kg body weight ICI 182,780, 100mg/kg body weight genistein, 2000mg/kg body weight SM6Met, and 10mg/kg body weight ICI 182,780 for three consecutive days. Animals were sacrificed on day four, (A) uteri were photographed, and (B) wet uterine/final body weight was determined. One-way ANOVA with Dunnett's post-test comparing all columns to either solvent control (*, P<0.05; **, P<0.01; ***, P<0.001) or E₂ (#, P<0.05; ##, P<0.01; ###, P<0.001). The dotted lines through the bars represent the values for solvent control or E₂. Average ± SEM is of at least five animals/group.

doi: 10.1371/journal.pone.0079223.g007




Figure 8. Evaluation of the effect of E₂ and SM6Met on the timing of vaginal opening. Immature female wistar rats were treated for 30 consecutive days with 100µg/kg body weight E₂ and 2000mg/kg body weight SM6Met and the day of vaginal opening was determined. One-way ANOVA with Dunnett's post-test comparing all columns to solvent control (*, P<0.05; **, P<0.01; ***, P<0.001). The dotted line through the bars represents the values for solvent control. Average ± SEM is of at least eight animals/group.

doi: 10.1371/journal.pone.0079223.g008

subtypes in the transactivation model, acted as an agonist for both subtypes in the transrepression model. The extracts of *C. subternata*, however, did not elicit such uniform effects in the transrepression model. SM6Met, a methanol extract, acting as an ER α antagonist and ER β agonist regarding transactivation, displayed antagonism towards ER α , in the absence of E₂ and towards ER β , in the presence of E₂. Similar antagonism towards ER α in the absence of E₂ has also been seen for the plant extract MF101 regarding IL6 mRNA expression [24]. The water extract, cup-of-tea, also changed its behavior, acting as an ER β antagonist for transrepression as opposed to an ER α antagonist for transactivation. These behavioral changes were not exclusive to the *Cyclopia* extracts as the polyphenols also displayed these characteristics. Luteolin, for example, displayed ER agonism through both subtypes in the transactivation model but was an ER α agonist and an ER β

antagonist in the transrepression model. The occurrence of mixed agonism and antagonism towards ER subtypes has also been observed for the xenoestrogen, Bisphenol A (BPA) [107].

As the current experiments were performed in the same cell line we have to look towards differences between the mechanisms of transactivation and transrepression for clarification of these results. Classically, transactivation is a product of ER dimer binding directly to the DNA sequence, however, tethering to DNA bound transcription factors (TFs) in the promoter region of affected genes has also been described [108-111]. Binding of the ER to DNA, whether it is direct or indirect, initiates the recruitment of co-activators, which then modulates transcription [112]. Regarding transrepression, specifically the repression of NF κ B driven genes, various mechanisms of ER mediated transrepression have been described [109]. The ER can bind to NF κ B and thereby prevent

DNA binding of the transcription factor [113,114], ligand bound ER present at promoter regions can recruit co-repressors [115,116], ligand bound ER α and activated NF κ B can compete for co-activator recruitment [117,118], or ER α , through a non-genomic pathway, inhibits translocation of activated NF κ B to the nucleus [119]. We can use this knowledge of the mechanism of action and combine it with what we know about SERMs and ER subtypes specific ligands to postulate a mechanism of action of *Cyclopia* agonism and antagonism. For the SERMs, three mechanisms of antagonism have been proposed [18]. SERMs can bind to the ER with a higher affinity than E₂ and block the binding of E₂, they can block the binding of co-activators, or SERMs can induce the recruitment of co-repressors [18,120,121]. Not much is known regarding the mechanism of SERM agonism [18], although it has been suggested that they can block the binding of co-repressors [121]. In addition, MF101 and liquiritigenin, both ER β selective agonists, although being able to bind to ER α , cannot recruit co-activators to ER α , and MF101 cannot promote the interaction of ER α with regulatory elements [15,24]. Furthermore, it has been suggested that SERMs may activate cell surface signaling pathways that results in ligand-independent activation of ERs [29,122,123].

Therefore, with regards to transactivation, we may postulate that the extracts of *Cyclopia* cannot transactivate via ER α as they are unable to recruit the necessary co-activators, while for ER β , P104 and SM6Met are able to do so. It is also possible that the extracts of *Cyclopia* cannot induce ER α interaction with regulatory elements. The observed ER α antagonism of E₂-induced transactivation may be due to the extracts binding to ER α and either inhibiting E₂ binding, inhibiting the recruitment of co-activators or stimulating the recruitment of co-repressors. In our transrepression model P104 behaves like E₂ and could be exerting its function by any of the NF κ B repression models discussed earlier. However, SM6Met displays ER α antagonism in the absence of E₂ and this antagonism is lost in the presence of E₂. Therefore, it is possible that SM6Met is unable to recruit co-repressors in the absence of E₂ and is unable to inhibit the E₂-induced recruitment of co-repressors. Furthermore, antagonism of ER β in the transrepression model by SM6Met and cup-of-tea may be due to the recruitment of co-activators to ER β .

Next we evaluated agonism and antagonism of *Cyclopia* extracts in a more complex environment where the ER subtypes are co-expressed. We used the MCF-7BUS cells, a breast carcinoma cell line, not only because it co-expresses the subtypes (Figure 2A), but also to evaluate the activity of the extracts in breast tissue cells. With regards to transactivation, all extracts of *Cyclopia* were agonists and are likely exerting this agonism through ER β as they were ER β agonists and ER α antagonists in COS-1 cells. Also, previously we discussed the possibility that the extracts may be unable to recruit co-activators to ER α or induce ER α -regulatory element interactions, which supports the idea that the *Cyclopia* extracts are mediating their transactivative effects in MCF-7BUS cells via ER β . Interestingly, the polyphenols, genistein and luteolin, having displayed ER agonism in COS-1 cells, in an environment where both ER subtypes are present displayed

only weak agonism, which may be attributed to the fact that lower concentrations were used in MCF-7BUS cells. However, when both subtypes are present these polyphenols display antagonism, which was not apparent when the subtypes were expressed separately. When both ER subtypes are expressed in the transrepression model, all the polyphenols as well as the *Cyclopia* extracts acted as agonists, while the water extract of *C. subternata* also displayed ER antagonism. The ER agonism of P104 in the transrepression model is thus not a cell type selective effect as it is seen in both the COS-1 (kidney) and MCF-7BUS (breast) cells. The ER antagonism of cup-of-tea in MCF-7BUS cells is likely mediated via ER β as ER β antagonism was observed in COS-1 cells transfected with ER β , but not in cells transfected with ER α . However, the SM6Met extract, which displayed antagonism for ER α and ER β in COS-1 cells, changes its behavior in the MCF-7BUS cells and acts as an ER agonist in the transrepression model. Furthermore, a similar switch in behavior is observed with the polyphenols as the subtype specific antagonism is abrogated in the presence of both ER subtypes. These observed behavioral changes of the *Cyclopia* extracts as well as the polyphenols in different tissues have also been observed for the SERM, tamoxifen [18]. Ball et al. [18] found that tamoxifen differentially regulated ER regulated genes in different cell lines and ascribed this phenomenon to the presence, or lack of, co-regulators in different tissues. Therefore, the differential effect of *Cyclopia* extracts as well as the polyphenols in cells from different tissues might be due to changes in the co-regulator environment.

As MCF-7BUS cells express both ER subtypes, we also have to consider the possibility of ER α / β heterodimer formation and the biological relevance thereof as opposed to homodimer formation in COS-1 cells expressing the ER subtypes in isolation. Using two phytoestrogens that are ER α / β heterodimer selective, cosmosiin and angolensin, it was shown that heterodimer formation, in the presence of these ligands, leads to higher activation of an ERE-promoter reporter construct than homodimers and furthermore that heterodimer formation has a growth inhibitory effect in breast and prostate epithelial cells [124]. Previous studies by Powell et al. [46] showed that the ER β selective agonist, liquiritigenin, which can bind to both ER subtypes, induces an ER α conformation that prefers heterodimerization with ER β , as opposed to forming ER α homodimers. Therefore, we cannot exclude heterodimer formation as an explanation for the strong agonist effect of the *Cyclopia* extracts in the transactivation model in MCF-7BUS cells.

Having evaluated the agonist and antagonist activity of *Cyclopia* extracts in a system where the ER subtypes were expressed separately and together, in a transactivation and a transrepression model, we increased the level of complexity by evaluating the effect of the extracts on MCF-7BUS cell proliferation, a system where the final cell phenotype is a product of not only the two ER subtypes but also of an integrated transactivation and transrepression system [39,98-100]. Although the *Cyclopia* extracts, like E₂, induced cell proliferation it was with either a significantly lower potency (P104 and SM6Met) or lower efficacy (P104 and cup-of-tea)

than E₂. Furthermore, in the presence of E₂, all of the *Cyclopia* extracts displayed antagonistic properties. Similarly, the polyphenols also induced cell proliferation with either lower efficacies or potencies than E₂ and some (genistein and enterodiol) also displayed antagonism. Previously, the agonist activity seen in the transactivation model in MCF-7BUS cells was ascribed to ER β activation and this is probably translating into weak induction of MCF-7BUS cell proliferation. Furthermore, liquiritigenin, an ER β selective agonist, although not able to induce significant MCF-7 cell growth in a mouse xenograft model [19,24], was able to induce proliferation of the ER α and ER β positive [125] osteoblast-like murine MC3T3-E1 cells [126]. The antagonism of E₂-induced cell proliferation by extracts of *Cyclopia* could be attributed to ER α antagonism (observed in the transactivation model in COS-1 cells), ER mediated repression of proliferation inducing genes (ER transrepression observed in MCF-7BUS transrepression model), ER β -mediated transcription (observed in the transactivation model in COS-1 cells) of anti-proliferative and anti-apoptotic genes [39,127], or they might favor the formation of ER α / β heterodimers, which has been suggested to have growth inhibitory effects in breast epithelial cells [124].

Furthermore, we also evaluated the estrogenic and antiestrogenic properties of the *Cyclopia* extracts in an *in vivo* model, an immature rat uterotrophic assay. For the first time we show *in vivo* biological activity of the phytoestrogenic extracts of *Cyclopia*. SM6Met and cup-of-tea, unlike E₂ and genistein, did not increase uterine weight and SM6Met, like the ER antagonist ICI 182,780, antagonized E₂-induced uterine growth. The ER α subtype is the major subtype expressed in the uterus with very low levels of ER β expressed [56,101]. Powell et al. [46] show that although ER β homodimers and ER α /ER β heterodimers are favored, genistein is capable of inducing ER α homodimers and activating ER α -induced transcription. Therefore, we can assume that the increase in uterine growth induced by genistein in the uterotrophic assay is a product of increased ER α homodimerization and hence, increased ER α mediated transcription. The ER β selective agonists, liquiritigenin and ERB-041, in contrast, do not induce uterine growth [19,128]. Thus, the findings regarding ER β selective agonists combined with the low levels of ER β in the uterus excludes ER β as the subtype eliciting the effect of *Cyclopia* extracts in the uterus. It is thus likely that the effect of *Cyclopia* extracts is due to ER α antagonism, as seen in the transactivation model in COS-1 cells, or that upon binding to the ER, the *Cyclopia* extracts induce a change in conformation that inhibits co-activator recruitment or activates co-repressor recruitment. The inability of the *Cyclopia* extracts to induce uterine growth, in contrast to MCF-7BUS cell proliferation, might also be attributed to either the differences in the concentration of co-regulators or the differences in co-regulator recruitment in the breast and uterus [129,130].

Having established ER agonist and/or antagonist activity of *Cyclopia* extracts, we look towards HPLC data, from the current and previous studies, to identify the polyphenol(s) responsible for the observed effects. The xanthenes, mangiferin and isomangiferin, were identified in all *Cyclopia* extracts, but as mangiferin has no estrogenic potential, while

isomangiferin has not previously been tested for estrogenicity [71], it is unlikely that the observed ER agonist/antagonist effects of *Cyclopia* can be ascribed to these polyphenols. However, mangiferin has been shown to inhibit the proliferation of breast cancer cells via ER independent mechanisms [131] and therefore, as mangiferin is present in all extracts at relatively high amounts it cannot be excluded as the polyphenol antagonizing E₂-induced MCF-7BUS cell proliferation. Of the remaining polyphenols identified in the extracts the only aglycone present is the flavone, luteolin. *In vitro*, luteolin binds to both of the ER subtypes, is an ER α and ER β agonist, induces MCF-7BUS cell proliferation, and antagonizes E₂-induced MCF-7BUS cell proliferation [62,63,71,132-134]. Therefore, with regards to the *Cyclopia* extracts, the ER β agonism observed in the transactivation model, the induction of MCF-7BUS cell proliferation, and the antagonism of E₂-induced cell proliferation may be ascribed to the presence luteolin in the extracts, however, the observed ER α antagonism in the transactivation model cannot. Although luteolin is present in all extracts, the concentration is low. However, the 7-O-rutinoside of luteolin, scolymoside, is present in substantial amounts in all of the *C. subternata* extracts (presence was not evaluated in P104). This rutinoside of luteolin has not previously been tested for estrogenicity [71], however, as glycosides may be hydrolyzed by intestinal β -glucosidases [135,136], the bioavailability of the aglycone, luteolin, and hence phytoestrogenicity of the extracts may increase upon hydrolysis of scolymoside. Furthermore, luteolin has been shown to have anti-tumor characteristics and can sensitize breast cancer cells to anti-tumor drugs such as tamoxifen [137] and therefore, the presence of luteolin, as well as scolymoside, in *Cyclopia* extracts can be seen as positive regarding chemoprevention as well as breast cancer treatment. Generally, the glycosides of polyphenols either display reduced estrogenic activity compared to the aglycones or have not been evaluated for estrogenicity [71]. Thus, if the hydrolysis of glycosides present in the *Cyclopia* extracts is considered, it allows us to evaluate the phytoestrogenicity of the aglycones alongside their glycosides: apigenin (aglycone of vicenin-2), eriodictyol (eriocitrin), hesperitin (hesperidin), phloretin (phloretin-3,5-di-C-glucoside), hydroxyphloretin (3-hydroxyphloretin-3',5'-di-C-hexoside), and iriflophenone (iriflophenone-2-C- β -glucoside and iriflophenone-di-O,C-hexoside). However, as β -glucosidases are produced by intestinal flora [138,139], consideration of glycoside metabolism will not help to identify the polyphenols responsible for *in vitro* results but may only be relevant for interpretation of *in vivo* results. For example, as luteolin and apigenin have been shown to significantly increase uterine weight, either in the presence or absence of estrogens [140,141], the effect elicited by *Cyclopia* extracts *in vivo* cannot be ascribed to luteolin, scolymoside, or vicenin-2. The effect of the other identified polyphenols has not been evaluated *in vivo* and therefore we cannot definitively attribute the *in vivo* effect of the *Cyclopia* extracts to any of these polyphenols. Of the glycosides, eriocitrin and hesperidin have been tested for phytoestrogenicity *in vitro* [71]. However, hesperidin does not bind to the ER [62] or activate an ERE-containing promoter reporter construct [133]. Eriocitrin, however, has been shown to

bind to only ER β [62], but no work has been done to elucidate the estrogenic effect elicited by this polyphenol. For the first time we identified the dihydrochalcone, aspalathin, in *Cyclopia*. Aspalathin has not been tested for estrogenicity but has been shown to inhibit the proliferation of liver cells [142], however, due to the presence of unique drug metabolizing enzymes in the liver, the possibility of aspalathin metabolites eliciting this effect cannot be excluded nor can the results be extrapolated to breast cancer cells. The phytoestrogenicity of the remaining glycosides and aglycones, as well as protocatechuic acid, has not been tested [71]. In summary, none of the compounds identified in the *Cyclopia* extracts can account for the observed ER α antagonism, some (luteolin and eriocitrin) may explain the observed ER β agonism and others (mangiferin and aspalathin) should not be excluded as possible effectors of ER-independent effects on proliferation. Therefore, thus far, we cannot with certainty ascribe the effects observed with *Cyclopia* extracts in this study to any of the individual constituents of our extracts. Although, further research regarding the polyphenol content, bioavailability, and estrogenic activity of our extracts is required to identify the compound causing the observed effects, we cannot exclude the possibility that a mixture of polyphenols is required to elicit the effects observed with *Cyclopia* extracts.

Physiologically, our results may be assessed both in terms of treatment of menopausal symptoms (hot flashes, osteoporosis, and increased inflammation [2-4,52,90-92]) and prevention of estrogen replacement associated side effects (breast cancer and uterine proliferation [5,6,52]). With regards to menopausal symptoms, the ER β agonist MF101 [24], has been shown in clinical trials to reduce hot flashes and thus, the ER β agonism of the *Cyclopia* extracts may be considered as a positive attribute. Furthermore, with regards to the postmenopausal surge in inflammatory disorders the fact that the *Cyclopia* extracts displayed agonism in the transrepression model in MCF-7BUS cells may also be considered as a positive attribute for the treatment of postmenopausal inflammatory disorders. With respect to the known roles of ER subtypes in breast cancer [15,22,37-43], the fact that extracts of *Cyclopia* antagonize ER α , while being ER β agonists, may be beneficial. In addition, the extracts were able to antagonize the proliferation of breast cancer cells in the presence of E $_2$ at lower concentrations than that required for breast cancer cell proliferation. Furthermore, not only do the *Cyclopia* extracts show potential as protectors against breast cancer development and inflammatory disorders, they also do this without promoting uterine growth, a negative SERM associated side effect [35,143].

Although *Cyclopia* extracts show potential to be developed as SERSMs, further work, which is ongoing, is needed to clarify their mechanism of action. This includes, but is not limited to, directly comparing the *Cyclopia* extracts with the known SERMs tamoxifen and raloxifene, investigating the effect of *Cyclopia* extracts on ER subtype levels, ER homo- or heterodimerization, induction or inhibition of co-regulator recruitment, and the modulation of cancer development and progression in a rat breast cancer model. In addition, further work is needed to identify the polyphenol(s) responsible for

eliciting the observed effects and the possibility that distinct polyphenols present in *Cyclopia*, rather than an individual polyphenol, may be causing the observed ER α agonism and ER β antagonism cannot be excluded.

Supporting Information

Figure S1. Determination of ERE-containing promoter reporter construct concentration. (A & B) COS-1 cells, transfected with equal amounts of (A) ER α and (B) ER β , and (C) MCF-7BUS cells were transfected with increasing amounts of the ERE-containing promoter reporter construct (ERE.vit2.luc) and treated with either solvent or E $_2$ to determine at which concentration of the ERE-containing promoter reporter construct the highest induction of E $_2$ is observed. The dotted line through the bars represents the values for solvent control. Fold induction is indicated in boxes above the E $_2$ columns. Average \pm SEM is of one experiment done with three to four repeats. (TIF)

Figure S2. Determination of NF κ B-containing promoter reporter construct concentration. (A & B) COS-1 cells, transfected with equal amounts of (A) ER α and (B) ER β , and (C) MCF-7BUS cells were transfected with increasing amounts of the NF κ B-containing promoter reporter construct (p(IL6 κ B)350hu.lL6Pluc+) and treated with either solvent, PMA or PMA + E $_2$ to determine at which concentration of the NF κ B-containing promoter reporter construct the highest repression by E $_2$ of PMA induction is observed. The dotted lines through the bars represent the values for either solvent control or 10ng/ml PMA. Percentage repression, where applicable, is indicated in boxes above the PMA + E $_2$ columns. Average \pm SEM is of one experiment done with three repeats. (TIF)

Figure S3. The effect of the SM6Met and cup-of-tea extracts on immature rat uterine growth. Immature female wistar rats were treated with 2000, 200, and 20mg/kg body weight SM6Met and cup-of-tea for three consecutive days. Animals were sacrificed on day four, (A) uteri were photographed and (B) wet and (C) blotted uterine/final body weight was determined. One-way ANOVA with Dunnett's post-test comparing all columns to solvent control (*, P<0.05; **, P<0.01; ***, P<0.001). The dotted line through the bars represents the values for solvent control. Average \pm SEM is of at least eight animals/group. (TIF)

Figure S4. The effect of E $_2$, genistein, extracts of *Cyclopia*, and ICI on body and liver weight. Immature female wistar rats were treated for three consecutive days with 100 μ g/kg body weight (BW) E $_2$, in the presence and absence of 2000mg/kg BW SM6Met or 10mg/kg BW ICI 182,780, 100mg/kg BW genistein, 2000, 200, or 20mg/kg BW SM6Met, 2000, 200, or 20mg/kg BW cup-of-tea, and 10mg/kg BW ICI 182,780 for three consecutive days. Animal were sacrificed on day four and changes in (A) body and (B) liver weights were

determined. One-way ANOVA with Dunnett's post-test comparing all columns to solvent control (*, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$). The dotted line through the bars represents the values for solvent control (A and B) and 100 μ g/kg BW E₂ (A). Average \pm SEM is of at least five animals/group. (TIF)

Figure S5. The effect of different concentration of the SM6Met and cup-of-tea extracts on the onset of vaginal opening. Immature female wistar rats were treated for 30 consecutive days with the SM6Met and cup-of-tea extracts and the day of vaginal opening was determined. One-way ANOVA with Dunnett's post-test comparing all columns to solvent control (*, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$). The dotted line through the bars represents the values for solvent control. Average \pm SEM is of at least eight animals/group. (TIF)

Table S1. Summary of ER agonism and antagonism of Cyclopia extracts.

References

- Ross RK, Paganini-Hill A, Wan PC, Pike MC (2000) Effect of hormone replacement therapy on breast cancer risk: Estrogen versus estrogen plus progestin. *JNCI. J Natl Cancer Inst* 92: 328-332. doi:10.1093/jnci/92.4.328. PubMed: 10675382.
- Burger HG, Hale GE, Dennerstein L, Robertson DM (2008) Cycle and hormone changes during perimenopause: The key role of ovarian function. *Menopause* 15: 603-612. doi:10.1097/gme.0b013e318174ea4d. PubMed: 18574431.
- Dennerstein L, Dudley EC, Hopper JL, Guthrie JR, Burger HG (2000) A prospective population-based study of menopausal symptoms. *Obstet Gynecol* 96: 351-358. doi:10.1016/S0029-7844(00)00930-3. PubMed: 10960625.
- Lindsay R (1996) The menopause and osteoporosis. *Obstet Gynecol* 87: 16S-19S. doi:10.1016/0029-7844(95)00430-0. PubMed: 8559548.
- Million women study collaborators (2003) Breast cancer and hormone-replacement therapy in the million women study. *The Lancet* 362: 419-427.
- Rossouw JE, Anderson GL, Prentice RL, LaCroix AZ, Kooperberg C et al. (2002) Risks and benefits of estrogen plus progestin in healthy postmenopausal women: Principal results from the women's health initiative randomized controlled trial. *JAMA* 288: 321-333. doi:10.1001/jama.288.3.321. PubMed: 12117397.
- Nand SL, Webster MA, Baber R, O'Connor V (1998) Bleeding pattern and endometrial changes during continuous combined hormone replacement therapy: the Ogen/Provera study group. *Obstet Gynecol* 91: 678-684. doi:10.1016/S0029-7844(98)00038-6. PubMed: 9572210.
- Anderson GL, Limacher M, Assaf AR, Bassford T, Beresford SA et al. (2004) Effects of conjugated equine estrogen in postmenopausal women with hysterectomy: The women's health initiative randomized controlled trial. *JAMA* 291: 1701-1712. doi:10.1001/jama.291.14.1701. PubMed: 15082697.
- Lerner LJ, Jordan VC (1990) Development of antiestrogens and their use in breast cancer: Eighth cain memorial award lecture. *Cancer Res* 50: 4177-4189. PubMed: 2194650.
- Jordan VC (1988) Chemosuppression of breast cancer with tamoxifen: Laboratory evidence and future clinical investigations. *Cancer Invest* 6: 589-595. doi:10.3109/07357908809082124. PubMed: 3063338.
- Hillisch A, Peters O, Kosemund D, Müller G, Walter A et al. (2004) Dissecting physiological roles of estrogen receptor alpha and beta with potent selective ligands from structure-based design. *Mol Endocrinol* 18: 1599-1609. doi:10.1210/me.2004-0050. PubMed: 15105439.
- Habel LA, Stanford JL (1993) Hormone receptors and breast cancer. *Epidemiol Rev* 15: 209-219. PubMed: 8405205.
- Stossi F, Barnett DH, Frasor J, Komm B, Lyttle CR et al. (2004) Transcriptional profiling of estrogen-regulated gene expression via estrogen receptor (ER) alpha or ERbeta in human osteosarcoma cells: Distinct and common target genes for these receptors. *Endocrinology* 145: 3473-3486. doi:10.1210/en.2003-1682. PubMed: 15033914.
- Hertrampf T, Seibel J, Laudenbach U, Fritzsche KH, Diel P (2008) Analysis of the effects of oestrogen receptor alpha (ERalpha)- and ERbeta-selective ligands given in combination to ovariectomized rats. *Br J Pharmacol* 153: 1432-1437. PubMed: 18246095.
- Paruthiyil S, Cvorova A, Zhao X, Wu Z, Sui Y et al. (2009) Drug and cell type-specific regulation of genes with different classes of estrogen receptor beta-selective agonists. *PLOS ONE* 4: e6271. doi:10.1371/journal.pone.0006271. PubMed: 19609440.
- Riggs BL, Hartmann LC (2003) Selective estrogen-receptor modulators - mechanisms of action and application to clinical practice. *N Engl J Med* 348: 618-629. doi:10.1056/NEJMra022219. PubMed: 12584371.
- Tee MK, Rogatsky I, Tzagarakis-Foster C, Cvorova A, An J et al. (2004) Estradiol and selective estrogen receptor modulators differentially regulate target genes with estrogen receptors alpha and beta. *Mol Cell Biol* 24: 1262-1272. PubMed: 14699072.
- Ball LJ, Levy N, Zhao X, Griffin C, Tagliaferri M et al. (2009) Cell type- and estrogen receptor-subtype specific regulation of selective estrogen receptor modulator regulatory elements. *Mol Cell Endocrinol* 299: 204-211. doi:10.1016/j.mce.2008.10.050. PubMed: 19059307.
- Mersereau JE, Levy N, Staub RE, Baggett S, Zogovic T et al. (2008) Lignin is a plant-derived highly selective estrogen receptor beta agonist. *Mol Cell Endocrinol* 283: 49-57. doi:10.1016/j.mce.2007.11.020. PubMed: 18177995.
- Felson DT, Zhang Y, Hannan MT, Kiel DP, Wilson PW et al. (1993) The effect of postmenopausal estrogen therapy on bone density in elderly women. *N Engl J Med* 329: 1141-1146. doi:10.1056/NEJM199310143291601. PubMed: 8377776.
- Rymer J, Wilson R, Ballard K (2003) Making decisions about hormone replacement therapy. *BMJ* 326: 322-326. doi:10.1136/bmj.326.7384.322. PubMed: 12574048.
- Paruthiyil S, Parmar H, Kerekatte V, Cunha GR, Firestone GL et al. (2004) Estrogen receptor beta inhibits human breast cancer cell proliferation and tumor formation by causing a G2 cell cycle arrest. *Cancer Res* 64: 423-428. doi:10.1158/0008-5472.CAN-03-2446. PubMed: 14729654.
- Weiderpass E, Adami HO, Baron JA, Magnusson C, Bergström R et al. (1999) Risk of endometrial cancer following estrogen replacement with and without progestins. *J Natl Cancer Inst* 91: 1131-1137. doi:10.1093/jnci/91.13.1131. PubMed: 10393721.
- Cvorova A, Paruthiyil S, Jones JO, Tzagarakis-Foster C, Clegg NJ et al. (2007) Selective activation of estrogen receptor-beta transcriptional pathways by an herbal extract. *Endocrinology* 148: 538-547. PubMed: 17095596.
- Flötotto T, Niederacher D, Hohmann D, Heimerzheim T, Dall P et al. (2004) Molecular mechanism of estrogen receptor (ER)alpha-specific, estradiol-dependent expression of the progesterone receptor (PR) B-isoform. *J Steroid Biochem Mol Biol* 88: 131-142. doi:10.1016/j.jsbmb.2003.11.004. PubMed: 15084345.

26. Delmas PD, Bjarnason NH, Mitlak BH, Ravoux AC, Shah AS et al. (1997) Effects of raloxifene on bone mineral density, serum cholesterol concentrations, and uterine endometrium in postmenopausal women. *N Engl J Med* 337: 1641-1647. doi:10.1056/NEJM199712043372301. PubMed: 9385122.
27. Love RR, Mazess RB, Barden HS, Epstein S, Newcomb PA et al. (1992) Effects of tamoxifen on bone mineral density in postmenopausal women with breast cancer. *N Engl J Med* 326: 852-856. doi:10.1056/NEJM199203263261302. PubMed: 1542321.
28. D'Amelio P, Isaia GC (2013) The use of raloxifene in osteoporosis treatment. *Expert Opin Pharmacother* 14: 949-956. doi:10.1517/14656566.2013.782002. PubMed: 23521229.
29. Jordan VC (2007) Chemoprevention of breast cancer with selective oestrogen-receptor modulators. *Nat Rev Cancer* 7: 46-53. doi:10.1038/nrc2048. PubMed: 17186017.
30. Jordan VC, O'Malley BW (2007) Selective estrogen-receptor modulators and antihormonal resistance in breast cancer. *J Clin Oncol* 25: 5815-5824. doi:10.1200/JCO.2007.11.3886. PubMed: 17893378.
31. MacGregor JL, Jordan VC (1998) Basic guide to the mechanisms of antiestrogen action. *Pharmacol Rev* 50: 151-196. PubMed: 9647865.
32. O'Regan RM, Jordan VC (2002) The evolution of tamoxifen therapy in breast cancer: Selective oestrogen-receptor modulators and downregulators. *Lancet Oncol* 3: 207-214. doi:10.1016/S1470-2045(02)00711-8. PubMed: 12067682.
33. Ettinger B, Black DM, Mitlak BH, Knickerbocker RK, Nickelsen T et al. (1999) Reduction of vertebral fracture risk in postmenopausal women with osteoporosis treated with raloxifene: Results from a 3-year randomized clinical trial. multiple outcomes of raloxifene evaluation (MORE) investigators. *JAMA* 282: 637-645. doi:10.1001/jama.282.7.637. PubMed: 10517716.
34. Cranney A, Adachi JD (2005) Benefit-risk assessment of raloxifene in postmenopausal osteoporosis. *Drug Saf* 28: 721-730. doi:10.2165/00002018-200528080-00006. PubMed: 16048357.
35. Fong CJ, Burgoon LD, Williams KJ, Jones AD, Forgacs AL et al. (2010) Effects of tamoxifen and ethynylestradiol cotreatment on uterine gene expression in immature, ovariectomized mice. *J Mol Endocrinol* 45: 161-173. doi:10.1677/JME-09-0158. PubMed: 20628019.
36. Vosse M, Renard F, Coibion M, Neven P, Nogaret JM et al. (2002) Endometrial disorders in 406 breast cancer patients on tamoxifen: The case for less intensive monitoring. *Eur J Obstet Gynecol Reprod Biol* 101: 58-63. doi:10.1016/S0301-2115(01)00516-4. PubMed: 11803101.
37. Ali S, Coombes RC (2000) Estrogen receptor alpha in human breast cancer: Occurrence and significance. *J Mammary Gland Biol Neoplasia* 5: 271-281. doi:10.1023/A:1009594727358. PubMed: 14973389.
38. Saji S, Jensen EV, Nilsson S, Rylander T, Warner M et al. (2000) Estrogen receptors alpha and beta in the rodent mammary gland. *Proc Natl Acad Sci U S A* 97: 337-342. doi:10.1073/pnas.97.1.337. PubMed: 10618419.
39. Chang EC, Frasier J, Komm B, Katzenellenbogen BS (2006) Impact of estrogen receptor beta on gene networks regulated by estrogen receptor alpha in breast cancer cells. *Endocrinology* 147: 4831-4842. doi:10.1210/en.2006-0563. PubMed: 16809442.
40. Lazennec G, Bresson D, Lucas A, Chauveau C, Vignon F (2001) ER beta inhibits proliferation and invasion of breast cancer cells. *Endocrinology* 142: 4120-4130. doi:10.1210/en.142.9.4120. PubMed: 11517191.
41. Shaaban AM, O'Neill PA, Davies MP, Sibson R, West CR et al. (2003) Declining estrogen receptor-beta expression defines malignant progression of human breast neoplasia. *Am J Surg Pathol* 27: 1502-1512. doi:10.1097/00000478-200312000-00002. PubMed: 14657709.
42. Ström A, Hartman J, Foster JS, Kietz S, Wimalasena J et al. (2004) Estrogen receptor beta inhibits 17 β -estradiol-stimulated proliferation of the breast cancer cell line T47D. *Proc Natl Acad Sci U S A* 101: 1566-1571. doi:10.1073/pnas.0308319100. PubMed: 14745018.
43. Latrich C, Stegerer A, Häring J, Schüller S, Ortmann O et al. (2013) Estrogen receptor beta agonists affect growth and gene expression of human breast cancer cell lines. *Steroids* 78: 195-202. doi:10.1016/j.steroids.2012.10.014. PubMed: 23153457.
44. Enmark E, Gustafsson JA (1999) Oestrogen receptors - an overview. *J Intern Med* 246: 133-138. doi:10.1046/j.1365-2796.1999.00545.x. PubMed: 10447781.
45. Monroe DG, Getz BJ, Johnsen SA, Riggs BL, Khosla S et al. (2003) Estrogen receptor isoform-specific regulation of endogenous gene expression in human osteoblastic cell lines expressing either ERalpha or ERbeta. *J Cell Biochem* 90: 315-326. doi:10.1002/jcb.10633. PubMed: 14505348.
46. Powell E, Xu W (2008) Intermolecular interactions identify ligand-selective activity of estrogen receptor alpha/beta dimers. *Proc Natl Acad Sci U S A* 105: 19012-19017. doi:10.1073/pnas.0807274105. PubMed: 19022902.
47. Tremblay A, Tremblay GB, Labrie F, Giguère V (1999) Ligand-independent recruitment of SRC-1 to estrogen receptor beta through phosphorylation of activation function AF-1. *Mol Cell* 3: 513-519. doi:10.1016/S1097-2765(00)80479-7. PubMed: 10230404.
48. Palmieri C, Cheng GJ, Saji S, Zelada-Hedman M, Wärrä A et al. (2002) Estrogen receptor beta in breast cancer. *Endocr Relat Cancer* 9: 1-13. doi:10.1677/erc.0.0090001. PubMed: 11914179.
49. Williams C, Edvardsson K, Lewandowski SA, Ström A, Gustafsson JA (2008) A genome-wide study of the repressive effects of estrogen receptor beta on estrogen receptor alpha signaling in breast cancer cells. *Oncogene* 27: 1019-1032. doi:10.1038/sj.onc.1210712. PubMed: 17700529.
50. Wardell SE, Nelson ER, Chao CA, McDonnell DP (2013) Bazedoxifene exhibits antiestrogenic activity in animal models of tamoxifen-resistant breast cancer: Implications for treatment of advanced disease. *Clin Cancer Res* 19: 2420-2431. doi:10.1158/1078-0432.CCR-12-3771. PubMed: 23536434.
51. Yeh WL, Shioda K, Coser KR, Rivizzigno D, McSweeney KR et al. (2013) Fulvestrant-induced cell death and proteasomal degradation of estrogen receptor alpha protein in MCF-7 cells require the CSK c-src tyrosine kinase. *PLOS ONE* 8: e60889. doi:10.1371/journal.pone.0060889. PubMed: 23593342.
52. Cvoró A, Tatomer D, Tee MK, Zogovic T, Harris HA et al. (2008) Selective estrogen receptor-beta agonists repress transcription of proinflammatory genes. *J Immunol* 180: 630-636. PubMed: 18097065.
53. Sun J, Huang YR, Harrington WR, Sheng S, Katzenellenbogen JA et al. (2002) Antagonists selective for estrogen receptor alpha. *Endocrinology* 143: 941-947. doi:10.1210/en.143.3.941. PubMed: 11861516.
54. Harrington WR, Sheng S, Barnett DH, Petz LN, Katzenellenbogen JA et al. (2003) Activities of estrogen receptor alpha- and beta-selective ligands at diverse estrogen responsive gene sites mediating transactivation or transrepression. *Mol Cell Endocrinol* 206: 13-22. doi:10.1016/S0303-7207(03)00255-7. PubMed: 12943986.
55. Meyers MJ, Sun J, Carlson KE, Marriner GA, Katzenellenbogen BS et al. (2001) Estrogen receptor-beta potency-selective ligands: Structure-activity relationship studies of diarylpropionitriles and their acetylene and polar analogues. *J Med Chem* 44: 4230-4251. doi:10.1021/jm010254a. PubMed: 11708925.
56. Harris HA (2007) Estrogen receptor-beta: Recent lessons from *in vivo* studies. *Mol Endocrinol* 21: 1-13. PubMed: 16556737.
57. Malamas MS, Manas ES, McDevitt RE, Gunawan I, Xu ZB et al. (2004) Design and synthesis of aryl diphenolic azoles as potent and selective estrogen receptor-beta ligands. *J Med Chem* 47: 5021-5040. doi:10.1021/jm049719y. PubMed: 15456246.
58. Brzezinski A, Debi A (1999) Phytoestrogens: The "natural" selective estrogen receptor modulators? *Eur J Obstet Gynecol Reprod Biol* 85: 47-51. doi:10.1016/S0301-2115(98)00281-4. PubMed: 10428321.
59. Tikkanen MJ, Adlercreutz H (2000) Dietary soy-derived isoflavone phytoestrogens. could they have a role in coronary heart disease prevention? *Biochem Pharmacol* 60: 1-5. doi:10.1016/S0006-2952(99)00409-8. PubMed: 10807939.
60. Oseni T, Patel R, Pyle J, Jordan VC (2008) Selective estrogen receptor modulators and phytoestrogens. *Planta Med* 74: 1656-1665. doi:10.1055/s-0028-1088304. PubMed: 18843590.
61. Kuiper GG, Lemmen JG, Carlsson B, Corton JC, Safe SH et al. (1998) Interaction of estrogenic chemicals and phytoestrogens with estrogen receptor beta. *Endocrinology* 139: 4252-4263. doi:10.1210/en.139.10.4252. PubMed: 9751507.
62. Verhoog NJD, Joubert E, Louw A (2007) Screening of four *Cyclopia* (honeybush) species for putative phyto-oestrogenic activity by oestrogen receptor binding assays. *S Afr J Sci* 103: 13-21.
63. Verhoog NJ, Joubert E, Louw A (2007) Evaluation of the phytoestrogenic activity of *Cyclopia genistoides* (honeybush) methanol extracts and relevant polyphenols. *J Agric Food Chem* 55: 4371-4381. doi:10.1021/jf063588n. PubMed: 17461595.
64. Glazier MG, Bowman MA (2001) A review of the evidence for the use of phytoestrogens as a replacement for traditional estrogen replacement therapy. *Arch Intern Med* 161: 1161-1172. doi:10.1001/archinte.161.9.1161. PubMed: 11343439.
65. Trock BJ, Hilakivi-Clarke L, Clarke R (2006) Meta-analysis of soy intake and breast cancer risk. *J Natl Cancer Inst* 98: 459-471. doi:10.1093/jnci/djj102. PubMed: 16595782.
66. Kies P (1951) Revision of the genus *Cyclopia* and notes on some other sources of bush tea. *Bothalia* 6: 161-176.

67. du Toit J, Joubert E, Britz TJ (1998) Honeybush tea: A rediscovered indigenous South African herbal tea. *J Sustain Agric* 12: 67-84. doi: 10.1300/J064v12n02_06.
68. Mfenyana C, DeBeer D, Joubert E, Louw A (2008) Selective extraction of *Cyclopia* for enhanced in vitro phytoestrogenicity and benchmarking against commercial phytoestrogen extracts. *J Steroid Biochem Mol Biol* 112: 74-86. doi:10.1016/j.jsmb.2008.08.005. PubMed: 18793725.
69. Kamara BI, Brandt EV, Ferreira D, Joubert E (2003) Polyphenols from honeybush tea (*Cyclopia intermedia*). *J Agric Food Chem* 51: 3874-3879. doi:10.1021/jf0210730. PubMed: 12797758.
70. Kamara BI, Brand DJ, Brandt EV, Joubert E (2004) Phenolic metabolites from honeybush tea (*Cyclopia subternata*). *J Agric Food Chem* 52: 5391-5395. doi:10.1021/jf040097z. PubMed: 15315375.
71. Louw A, Joubert E, Visser K (2013) Phytoestrogenic potential of *Cyclopia* extracts and polyphenols. *Planta Med* 79: 580-590. doi: 10.1055/s-0032-1328463. PubMed: 23609108.
72. Joubert E, Gelderblom WC, Louw A, de Beer D (2008) South african herbal teas: *Aspalathus linearis*, *Cyclopia* spp. and *Athrixia phyllocoides* - A review. *J Ethnopharmacol* 119: 376-412. doi:10.1016/j.jep.2008.06.014. PubMed: 18621121.
73. Primiano T, Yu R, Kong AT (2001) Signal transduction events elicited by natural products that function as cancer chemopreventive agents. *Pharm Biol* 39: 83. doi:10.1076/phbi.39.2.83.6256.
74. Lee SO, Nadiminty N, Wu XX, Lou W, Dong Y et al. (2005) Selenium disrupts estrogen signaling by altering estrogen receptor expression and ligand binding in human breast cancer cells. *Cancer Res* 65: 3487-3492. PubMed: 15833885.
75. de Beer D, Schulze AE, Joubert E, de Villiers A, Malherbe CJ et al. (2012) Food ingredient extracts of *Cyclopia subternata* (honeybush): Variation in phenolic composition and antioxidant capacity. *Molecules* 17: 14602-14624. doi:10.3390/molecules171214602. PubMed: 23222906.
76. Villalobos M, Olea N, Brotons JA, Olea-Serrano MF, Ruiz de Almodovar JM et al. (1995) The E-screen assay: A comparison of different MCF7 cell stocks. *Environ Health Perspect* 103: 844-850. doi: 10.1289/ehp.95103844. PubMed: 7498097.
77. Flouriot G, Brand H, Denger S, Metivier R, Kos M et al. (2000) Identification of a new isoform of the human estrogen receptor- α (hER α) that is encoded by distinct transcripts and that is able to repress hER α activation function 1. *EMBO J* 19: 4688-4700. doi:10.1093/emboj/19.17.4688. PubMed: 10970861.
78. Denger S, Reid G, Brand H, Kos M, Gannon F (2001) Tissue-specific expression of human ER α and ER β in the male. *Mol Cell Endocrinol* 178: 155-160. doi:10.1016/S0303-7207(01)00417-8. PubMed: 11403905.
79. Hall JM, McDonnell DP, Korach KS (2002) Allosteric regulation of estrogen receptor structure, function, and coactivator recruitment by different estrogen response elements. *Mol Endocrinol* 16: 469-486. doi: 10.1210/me.16.3.469. PubMed: 11875105.
80. Plaisance S, Vanden Berghe W, Boone E, Fiers W, Haegeman G (1997) Recombination signal sequence binding protein jkappa is constitutively bound to the NF- κ B site of the interleukin-6 promoter and acts as a negative regulatory factor. *Mol Cell Biol* 17: 3733-3743. PubMed: 9199307.
81. Bradford MM (1976) A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* 72: 248-254. doi: 10.1016/0003-2697(76)90527-3. PubMed: 942051.
82. Kanno J, Onyon L, Haseman J, Fenner-Crisp P, Ashby J et al. (2001) The OECD program to validate the rat uterotrophic bioassay to screen compounds for *in vivo* estrogenic responses: Phase 1. *Environ Health Perspect* 109: 785-794. doi:10.1289/ehp.01109785. PubMed: 11564613.
83. de Lima; Toccafondo Vieira M, Duarte RF, Campos LM, Nunan Ede A (2008) Comparison of the estrogenic potencies of standardized soy extracts by immature rat uterotrophic bioassay. *Phytomedicine* 15: 31-37. doi:10.1016/j.phymed.2007.06.006. PubMed: 17689940. doi: 10.1016/j.phymed.2007.06.006. PubMed: 17689940.
84. Duncan AM, Phipps WR, Kurzer MS (2003) Phyto-oestrogens. *Best Pract Res Clin Endocrinol Metab* 17: 253-271. doi:10.1016/S1521-690X(02)00103-3. PubMed: 12787551.
85. Murkies AL, Wilcox G, Davis SR (1998) Clinical review 92: Phytoestrogens. *J Clin Endocrinol Metab* 83: 297-303. doi:10.1210/jc.83.2.297. PubMed: 9467531.
86. Schreurs R, Lanser P, Seinen W, van der Burg B (2002) Estrogenic activity of UV filters determined by an in vitro reporter gene assay and an *in vivo* transgenic zebrafish assay. *Arch Toxicol* 76: 257-261. doi: 10.1007/s00204-002-0348-4. PubMed: 12107642.
87. Kulling SE, Lehmann L, Metzler M (2002) Oxidative metabolism and genotoxic potential of major isoflavone phytoestrogens. *J Chromatogr B Anal Technol Biomed Life Sci* 777: 211-218. doi:10.1016/S1570-0232(02)00215-5. PubMed: 12270214.
88. Riggs BL, Khosla S, Melton LJ 3rd (2002) Sex steroids and the construction and conservation of the adult skeleton. *Endocr Rev* 23: 279-302. doi:10.1210/er.23.3.279. PubMed: 12050121.
89. Nilsson S, Mäkelä S, Treuter E, Tujague M, Thomsen J et al. (2001) Mechanisms of estrogen action. *Physiol Rev* 81: 1535-1565. PubMed: 11581496.
90. Romas E, Martin TJ (1997) Cytokines in the pathogenesis of osteoporosis. *Osteoporos Int* 7 Suppl 3: S47-S53. doi:10.1007/BF03194342. PubMed: 9536302.
91. Caquevel M, Lebeurrier N, Chéenne S, Vivien D (2004) Cytokines in neuroinflammation and alzheimer's disease. *Curr Drug Targets* 5: 529-534. doi:10.2174/1389450043345308. PubMed: 15270199.
92. Pacifici R (1996) Estrogen, cytokines, and pathogenesis of postmenopausal osteoporosis. *J Bone Miner Res* 11: 1043-1051. PubMed: 8854239.
93. Cao Y, Karin M (2003) NF- κ B in mammary gland development and breast cancer. *J Mammary Gland Biol Neoplasia* 8: 215-223. doi: 10.1023/A:1025905008934. PubMed: 14635796.
94. Baud V, Karin M (2009) Is NF- κ B a good target for cancer therapy? hopes and pitfalls. *Nat Rev Drug Discov* 8: 33-40. doi: 10.1038/nrd2781. PubMed: 19116625.
95. Grivnenkov SI, Greten FR, Karin M (2010) Immunity, inflammation, and cancer. *Cell* 140: 883-899. doi:10.1016/j.cell.2010.01.025. PubMed: 20303878.
96. Libermann TA, Baltimore D (1990) Activation of interleukin-6 gene expression through the NF- κ B transcription factor. *Mol Cell Biol* 10: 2327-2334. PubMed: 2183031.
97. Shirakawa F, Mizel SB (1989) *In vitro* activation and nuclear translocation of NF- κ B catalyzed by cyclic AMP-dependent protein kinase and protein kinase C. *Mol Cell Biol* 9: 2424-2430. PubMed: 2548081.
98. Nicholson RI, McClelland RA, Robertson JF, Gee JM (1999) Involvement of steroid hormone and growth factor cross-talk in endocrine response in breast cancer. *Endocr Relat Cancer* 6: 373-387. doi:10.1677/erc.0.0060373. PubMed: 10516852.
99. Knabbe C, Lippman ME, Wakefield LM, Flanders KC, Kasid A et al. (1987) Evidence that transforming growth factor-beta is a hormonally regulated negative growth factor in human breast cancer cells. *Cell* 48: 417-428. doi:10.1016/0092-8674(87)90193-0. PubMed: 2879636.
100. Hanahan D, Weinberg RA (2011) Hallmarks of cancer: The next generation. *Cell* 144: 646-674. doi:10.1016/j.cell.2011.02.013. PubMed: 21376230.
101. Harris HA, Katzenellenbogen JA, Katzenellenbogen BS (2002) Characterization of the biological roles of the estrogen receptors, ER α and ER β , in estrogen target tissues in vivo through the use of an ER α -selective ligand. *Endocrinology* 143: 4172-4177. doi: 10.1210/en.2002-220403. PubMed: 12399409.
102. Odum J, Lefevre PA, Tittensor S, Paton D, Routledge EJ et al. (1997) The rodent uterotrophic assay: Critical protocol features, studies with nonyl phenols, and comparison with a yeast estrogenicity assay. *Regul Toxicol Pharmacol* 25: 176-188. doi:10.1006/rtph.1997.1100. PubMed: 9185893.
103. Vogel VG, Costantino JP, Wickerham DL, Cronin WM, Cecchini RS et al. (2006) Effects of tamoxifen vs raloxifene on the risk of developing invasive breast cancer and other disease outcomes: The NSABP study of tamoxifen and raloxifene (STAR) P-2 trial. *JAMA* 295: 2727-2741. doi:10.1001/jama.295.23.joc60074. PubMed: 16754727.
104. Schmidt C (2010) Third-generation SERMs may face uphill battle. *J Natl Cancer Inst* 102: 1690-1692. doi:10.1093/jnci/djq477. PubMed: 21060064.
105. Hall J, McDonnell D (2008) Selective estrogen receptor modulators: From bench, to bedside, and back again. *Menopausal Med* 16.
106. Maximov PY, Lee TM, Jordan VC (2013) The discovery and development of selective estrogen receptor modulators (SERMs) for clinical practice. *Curr Clin Pharmacol* 8: 135-155.
107. Hiroi H, Tsutsumi O, Momoeda M, Takai Y, Osuga Y et al. (1999) Differential interactions of bisphenol A and 17 β -estradiol with estrogen receptor alpha (ER α) and ER β . *Endocr J* 46: 773-778. doi:10.1507/endocrj.46.773. PubMed: 10724352.
108. Kushner PJ, Agard DA, Greene GL, Scanlan TS, Shiau AK et al. (2000) Estrogen receptor pathways to AP-1. *J Steroid Biochem Mol Biol* 74: 311-317. doi:10.1016/S0960-0760(00)00108-4. PubMed: 11162939.
109. Frasar J, Weaver A, Pradhan M, Dai Y, Miller LD et al. (2009) Positive cross-talk between estrogen receptor and NF- κ B in breast cancer.

- Cancer Res 69: 8918-8925. doi:10.1158/0008-5472.CAN-09-2608. PubMed: 19920189.
110. Safe S, Kim K (2008) Non-classical genomic estrogen receptor (ER)/specificity protein and ER/activating protein-1 signaling pathways. *J Mol Endocrinol* 41: 263-275. doi:10.1677/JME-08-0103. PubMed: 18772268.
 111. Scafonas A, Reszka AA, Kimmel DB, Hou XS, Su Q et al. (2008) Agonist-like SERM effects on ER α -mediated repression of MMP1 promoter activity predict in vivo effects on bone and uterus. *J Steroid Biochem Mol Biol* 110: 197-206. doi:10.1016/j.jsmb.2007.10.013. PubMed: 18508261.
 112. Gruber CJ, Gruber DM, Gruber IM, Wieser F, Huber JC (2004) Anatomy of the estrogen response element. *Trends Endocrinol Metab* 15: 73-78. doi:10.1016/j.tem.2004.01.008. PubMed: 15036253.
 113. Galien R, Garcia T (1997) Estrogen receptor impairs interleukin-6 expression by preventing protein binding on the NF-kappaB site. *Nucleic Acids Res* 25: 2424-2429. doi:10.1093/nar/25.12.2424. PubMed: 9171095.
 114. Ray P, Ghosh SK, Zhang DH, Ray A (1997) Repression of interleukin-6 gene expression by 17 beta-estradiol: Inhibition of the DNA-binding activity of the transcription factors NF-IL6 and NF-kappa B by the estrogen receptor. *FEBS Lett* 409: 79-85. doi:10.1016/S0014-5793(97)00487-0. PubMed: 9199508.
 115. Cvoro A, Tzagarakis-Foster C, Tatomer D, Paruthiyil S, Fox MS et al. (2006) Distinct roles of unliganded and liganded estrogen receptors in transcriptional repression. *Mol Cell* 21: 555-564. doi:10.1016/j.molcel.2006.01.014. PubMed: 16483936.
 116. Gosselin D, Rivest S (2011) Estrogen receptor transrepresses brain inflammation. *Cell* 145: 495-497. doi:10.1016/j.cell.2011.04.018. PubMed: 21565607.
 117. Harnish DC, Scicchitano MS, Adelman SJ, Lyttle CR, Karathanasis SK (2000) The role of CBP in estrogen receptor cross-talk with nuclear factor-kappaB in HepG2 cells. *Endocrinology* 141: 3403-3411. doi:10.1210/en.141.9.3403. PubMed: 10965913.
 118. Nettles KW, Gil G, Nowak J, Métiévir R, Sharma VB et al. (2008) CBP is a dosage-dependent regulator of nuclear factor-kappaB suppression by the estrogen receptor. *Mol Endocrinol* 22: 263-272. PubMed: 17932106.
 119. Ghisletti S, Meda C, Maggi A, Vegeto E (2005) 17beta-estradiol inhibits inflammatory gene expression by controlling NF-kappaB intracellular localization. *Mol Cell Biol* 25: 2957-2968. doi:10.1128/MCB.25.8.2957-2968.2005. PubMed: 15798185.
 120. Shang Y, Hu X, DiRenzo J, Lazar MA, Brown M (2000) Cofactor dynamics and sufficiency in estrogen receptor-regulated transcription. *Cell* 103: 843-852. doi:10.1016/S0092-8674(00)00188-4. PubMed: 11136970.
 121. Nettles KW, Greene GL (2005) Ligand control of coregulator recruitment to nuclear receptors. *Annu Rev Physiol* 67: 309-333. doi:10.1146/annurev.physiol.66.032802.154710. PubMed: 15709961.
 122. Hall JM, Couse JF, Korach KS (2001) The multifaceted mechanisms of estradiol and estrogen receptor signaling. *J Biol Chem* 276: 36869-36872. doi:10.1074/jbc.R100029200. PubMed: 11459850.
 123. Aronica SM, Kraus WL, Katzenellenbogen BS (1994) Estrogen action via the cAMP signaling pathway: Stimulation of adenylate cyclase and cAMP-regulated gene transcription. *Proc Natl Acad Sci U S A* 91: 8517-8521. doi:10.1073/pnas.91.18.8517. PubMed: 8078914.
 124. Powell E, Shanle E, Brinkman A, Li J, Keles S et al. (2012) Identification of estrogen receptor dimer selective ligands reveals growth-inhibitory effects on cells that co-express ER α and ER β . *PLOS ONE* 7: e30993. doi:10.1371/journal.pone.0030993. PubMed: 22347418.
 125. Chen X, Garner SC, Quarles LD, Anderson JJ (2003) Effects of genistein on expression of bone markers during MC3T3-E1 osteoblastic cell differentiation. *J Nutr Biochem* 14: 342-349. doi:10.1016/S0955-2863(03)00056-1. PubMed: 12873716.
 126. Choi EM (2012) Liquiritigenin isolated from glycyrrhiza uralensis stimulates osteoblast function in osteoblastic MC3T3-E1 cells. *Int Immunopharmacol* 12: 139-143. doi:10.1016/j.intimp.2011.11.003. PubMed: 22116056.
 127. Helguero LA, Faulds MH, Gustafsson JA, Haldosén LA (2005) Estrogen receptors alpha (ER α) and beta (ER β) differentially regulate proliferation and apoptosis of the normal murine mammary epithelial cell line HC11. *Oncogene* 24: 6605-6616. doi:10.1038/sj.onc.1208807. PubMed: 16007178.
 128. Harris HA, Albert LM, Leathurby Y, Malamas MS, Mewshaw RE et al. (2003) Evaluation of an estrogen receptor-beta agonist in animal models of human disease. *Endocrinology* 144: 4241-4249. doi:10.1210/en.2003-0550. PubMed: 14500559.
 129. Shang Y, Brown M (2002) Molecular determinants for the tissue specificity of SERMs. *Science* 295: 2465-2468. doi:10.1126/science.1068537. PubMed: 11923541.
 130. Gronemeyer H, Gustafsson JA, Laudet V (2004) Principles for modulation of the nuclear receptor superfamily. *Nat Rev Drug Discov* 3: 950-964. doi:10.1038/nrd1551. PubMed: 15520817.
 131. Li H, Huang J, Yang B, Xiang T, Yin X et al. (2013) Mangiferin exerts antitumor activity in breast cancer cells by regulating matrix metalloproteinases, epithelial to mesenchymal transition, and beta-catenin signaling pathway. *Toxicol Appl Pharmacol*.
 132. Han DH, Denison MS, Tachibana H, Yamada K (2002) Relationship between estrogen receptor-binding and estrogenic activities of environmental estrogens and suppression by flavonoids. *Biosci Biotechnol Biochem* 66: 1479-1487. doi:10.1271/bbb.66.1479. PubMed: 12224631.
 133. Zhu JT, Choi RC, Chu GK, Cheung AW, Gao QT et al. (2007) Flavonoids possess neuroprotective effects on cultured pheochromocytoma PC12 cells: A comparison of different flavonoids in activating estrogenic effect and in preventing beta-amyloid-induced cell death. *J Agric Food Chem* 55: 2438-2445. doi:10.1021/jf063299z. PubMed: 17323972.
 134. Collins-Burow BM, Burow ME, Duong BN, McLachlan JA (2000) Estrogenic and antiestrogenic activities of flavonoid phytochemicals through estrogen receptor binding-dependent and -independent mechanisms. *Nutr Cancer* 38: 229-244. doi:10.1207/S15327914NC382_13. PubMed: 11525602.
 135. Setchell KD, Brown NM, Zimmer-Nechemias L, Brashear WT, Wolfe BE et al. (2002) Evidence for lack of absorption of soy isoflavone glycosides in humans, supporting the crucial role of intestinal metabolism for bioavailability. *Am J Clin Nutr* 76: 447-453. PubMed: 12145021.
 136. Németh K, Plumb GW, Berrin JG, Juge N, Jacob R et al. (2003) Deglycosylation by small intestinal epithelial cell beta-glucosidases is a critical step in the absorption and metabolism of dietary flavonoid glycosides in humans. *Eur J Nutr* 42: 29-42. doi:10.1007/s00394-003-0397-3. PubMed: 12594539.
 137. Tu SH, Ho CT, Liu MF, Huang CS, Chang HW et al. (2013) Luteolin sensitises drug-resistant human breast cancer cells to tamoxifen via the inhibition of cyclin E2 expression. *Food Chem* 141: 1553-1561. doi:10.1016/j.foodchem.2013.04.077. PubMed: 23790951.
 138. Slavin JL, Karr SC, Hutchins AM, Lampe JW (1998) Influence of soybean processing, habitual diet, and soy dose on urinary isoflavonoid excretion. *Am J Clin Nutr* 68: 1492S-1495S. PubMed: 9848522.
 139. Friend RH, Chang GW (1984) A colon-specific drug-delivery system based on drug glycosides and the glycosidases of colonic bacteria. *J Med Chem* 27: 261-266. doi:10.1021/jm00369a005. PubMed: 6699871.
 140. Hiremath SP, Badami S, Hunasagatta SK, Patil SB (2000) Antifertility and hormonal properties of flavones of striga orobanchioides. *Eur J Pharmacol* 391: 193-197. doi:10.1016/S0014-2999(99)00723-2. PubMed: 10720651.
 141. Stroheker T, Chagnon MC, Pinnert MF, Berges R, Canivenc-Lavier MC (2003) Estrogenic effects of food wrap packaging xenoestrogens and flavonoids in female wistar rats: A comparative study. *Reprod Toxicol* 17: 421-432. doi:10.1016/S0890-6238(03)00044-3. PubMed: 12849853.
 142. Snijman PW, Swanevelder S, Joubert E, Green IR, Gelderblom WC (2007) The antimutagenic activity of the major flavonoids of rooibos (*Aspalathus linearis*): Some dose-response effects on mutagen activation-flavonoid interactions. *Mutat Res* 631: 111-123. doi:10.1016/j.mrgentox.2007.03.009. PubMed: 17537670.
 143. Fong CJ, Burgoon LD, Williams KJ, Forgacs AL, Zacharewski TR (2007) Comparative temporal and dose-dependent morphological and transcriptional uterine effects elicited by tamoxifen and ethynylestradiol in immature, ovariectomized mice. *BMC Genomics* 8: 151. doi:10.1186/1471-2164-8-151. PubMed: 17555576.