

# Evaluation of the visual prostate symptom score in a male population with great language diversity and limited education: A study from Namibia

C F Heyns,<sup>1</sup> MB ChB, MMed (Urol) PhD, FCSSA (Urol); B A Steenkamp,<sup>1</sup> BSc, MB ChB; J Chiswo,<sup>2</sup> MB ChB; G A Stellmacher,<sup>2</sup> MB ChB, MMed (Urol); H E A Förtsch,<sup>2</sup> MB ChB, MMed (Urol), FCSSA (Urol); A van der Merwe,<sup>1</sup> MB ChB, MMed (Urol) FCUrol (SA)

<sup>1</sup> Department of Urology, Faculty of Health Sciences, Stellenbosch University and Tygerberg Hospital, Tygerberg, Cape Town, South Africa

<sup>2</sup> Department of Urology, Windhoek Central Hospital, Namibia

Corresponding author: C F Heyns (cfh2@sun.ac.za)

**Background.** A visual prostate symptom score (VPSS) using pictograms was developed to assess the force of the urinary stream, urinary frequency, nocturia and quality of life (QoL).

**Objective.** To compare the VPSS with the international prostate symptom score (IPSS) and maximum (Qmax) and average (Qave) urinary flow rates in men from diverse language groups with limited schooling.

**Methods.** Men with lower urinary tract symptoms admitted to the urology ward at Windhoek Central Hospital, Namibia, were evaluated. Patients who were unable to complete the questionnaires alone were assisted by a doctor or nurse. Local ethics committee approval was obtained. Statistical analysis was performed using Student's *t*-test and Spearman's rank correlation test.

**Results.** One hundred men (mean age 56.3 years, range 20.1 - 95.4) were evaluated over a period of one year. All the men understood one or more of 15 languages, and 30 were illiterate; 32 had <5 years of schooling, 34 had 5 - 9 years and 34 had >9 years. The VPSS took significantly less time to complete than the IPSS. There were statistically significant correlations between the total VPSS and IPSS scores, between the four VPSS questions and the corresponding IPSS questions, and between Qmax and Qave and the VPSS total and VPSS questions on the force of the urinary stream and QoL.

**Conclusion.** The VPSS pictograms depicting the force of the urinary stream and QoL correlated significantly with Qmax and Qave, indicating that they can be used as single-item questions to rapidly assess bladder outflow obstruction in men with limited education.

*S Afr Med J* 2014;104(5):353-357. DOI:10.7196/SAMJ.7917



The international prostate symptom score (IPSS) (Fig. 1) is used widely to assess lower urinary tract symptoms (LUTS) in men before and after treatment.<sup>[1]</sup> The IPSS is relatively complicated, especially for patients with low education levels, and elderly patients with visual impairment may find it difficult to read the small print.<sup>[2-4]</sup>

Illiteracy is a major problem with administration of the IPSS in developing countries.<sup>[5]</sup> Translating the IPSS into various languages requires extensive validation studies and constant surveillance to ensure consistency.<sup>[6]</sup>

A drawback of the IPSS is that it does not evaluate all lower urinary tract symptoms, such as incontinence, pain or macroscopic haematuria. More comprehensive questionnaires have been developed, such as the Core LUTS Score with 10 questions, and the EpiLUTS tool evaluating 18 symptoms.<sup>[7,8]</sup> While providing more information, these require a high level of patient education, are time-consuming to administer and analyse, and may not be useful for clinical application.

Some authors have proposed using fewer questions that may be more useful in clinical practice. A previous study has indicated that an abbreviated form of the IPSS consisting of three or four questions only may be used to assess patients more rapidly.<sup>[3]</sup> It has also been suggested that the IPSS quality of life (QoL) question alone is as useful as the more complicated Benign Prostatic Hypertrophy Impact Index (BPH-II).<sup>[9]</sup>

Others have proposed the use of single-item questionnaires, which have the advantages of speed and ease of administration and interpretation, but the disadvantage that a single question cannot quantify the variety of symptoms present, their intensity, duration and frequency, or the distress they cause.<sup>[1]</sup> A recent study indicated that patients' perceptions of the severity of and changes in their symptoms can be captured with single-item questions that are simple, valid and easily administered.<sup>[10]</sup>

We developed a visual prostate symptom score (VPSS) that uses pictograms to assess the force of the urinary stream, daytime urinary frequency, nocturia and QoL.<sup>[11-13]</sup> At the time we were unaware that Peeling<sup>[14]</sup> had described a pictogram representing the force of the urinary stream in 1989, and that in 2006 Ushijima *et al.*<sup>[15]</sup> had described a visual analogue scale for the IPSS, consisting of facial pictograms with expressions ranging from a broad smile (delighted) to a tearful face (distressed).

## Objective

To compare the IPSS and VPSS with urinary flow rate measurements in men with LUTS who were illiterate or had very limited schooling, and who spoke a diversity of languages.

## Methods

Men with LUTS admitted to the urology ward at Windhoek Central Hospital, Namibia, were evaluated with the IPSS (Fig. 1) and the VPSS (Fig. 2). Maximum (Qmax) and average (Qave) urinary flow

rates were measured using a Dantec Urodyn flowmeter. The original version of the VPSS<sup>[11,12]</sup> was modified in a subsequent study.<sup>[13]</sup> The pictogram of the force of the urinary stream (question 3 in the original version) was moved to become question A, because the pictogram of urination made it easier for the patient to understand the pictograms of frequency and nocturia (questions 2 and 3 in the original, and questions B and C in the modified VPSS).<sup>[11-13]</sup>

The following demographic data were collected: the patient's age, his mother's language, his home language, which languages he was able to understand, speak and read or write, years of schooling, occupation and monthly income. The patient was requested to fill in the IPSS and VPSS, with assistance if necessary from a physician and/or a nurse able to speak the patient's home language. The time taken to fill in the IPSS and the VPSS was recorded using a stopwatch.

The following clinical data were collected: findings on digital rectal examination, urine dipstick analysis, Qmax and Qave, voided volume (VV), serum creatinine, serum prostate-specific antigen, findings on cystoscopy, treatment given and histological results. Patients were requested to return for follow-up at 3 and 6 months after treatment.

The study protocol was approved by the Ministry of Health and Social Services of Namibia. Statistical analysis was performed with Student's *t*-test (paired or unpaired, as appropriate) for parametric data, Spearman's rank test for correlations and Fisher's exact test for contingency tables, using GraphPad InStat software.

**Results**

During the period February 2012 - February 2013, a total of 100 new patients were evaluated; 39 returned for follow-up, with a total of 155 visits (some patients had more than one follow-up visit). The distance between the patient's home and the hospital was <100 km in 44 cases, 100 - 500 km in 32, 500 - 1 000 km in 22 and >1 000 km in

two. Overall, 25 patients had never attended school. The number of years at school was <5 years in 32 cases, 5 - 9 years in 34 and >9 years in 34. Overall, 30 patients were not able to read or write.

Table 1 shows the mother languages of the patients, and the languages they spoke at home and were able to understand, speak and read or write. The number of languages understood by the patients was one in 21% of cases, two in 35%, three in 23%, four in 16% and five in 5%. Table 2 shows the number of languages understood by the patients in the five most common language groups.

On admission to hospital, 19 men had an indwelling suprapubic catheter and 14 an indwelling transurethral catheter. The mean duration of catheterisation was 3.7 months (range 0.1 - 41.3). For these men, the symptom scores for the month before catheterisation were obtained; uroflowmetry could not be undertaken. The final diagnosis was urethral stricture in 61 men, benign prostatic hyperplasia (BPH) in 32, no abnormality on cystoscopy in five, prostate cancer in three, prostatitis in three, bladder stones in two and bladder neck stenosis in one (some patients had more than one diagnosis). The surgical treatment involved direct-vision internal urethrotomy in 60 cases, transurethral resection of the prostate in 17, open prostatectomy in six, bladder neck incision in one and urethroplasty in one.

Both the IPSS and VPSS were filled in without assistance by only 18 of patients at the first visit, 82 requiring assistance. At follow-up the IPSS and VPSS were filled in without assistance by 7/55 (12.7%) and 8/55 (14.5%) of patients. For all visits, the mean time taken to complete the IPSS and VPSS was 278 seconds (range 80 - 690) and 124 seconds (30 - 556), respectively (*p*<0.0001). Comparing the first with follow-up visits, the mean time taken to complete the IPSS was 306 seconds (80 - 690) v. 227 seconds (80 - 572), and the mean time taken to complete the VPSS was 173 seconds (30 - 556) v. 100 seconds (30 - 270) (*p*<0.001). Table 3 compares the groups of men with schooling <5 v. >9 years.

**International Prostate Symptom Score (I-PSS)<sup>1</sup>**

Date: \_\_\_\_\_

Circle your score for each below.

	Not at all	Less than 1 time in 5	Less than half the time	About half the time	More than half the time	Almost always
1 Over the past month or so, how often have you had a sensation of not emptying your bladder completely after you finished urinating?	0	1	2	3	4	5
2 Over the past month or so, how often have you had to urinate again less than two hours after you finished urinating?	0	1	2	3	4	5
3 Over the past month or so, how often have you found that you stopped and started again several times when you urinated?	0	1	2	3	4	5
4 Over the past month or so, how often have you found it difficult to postpone urination?	0	1	2	3	4	5
5 Over the past month or so, how often have you had a weak urinary stream?	0	1	2	3	4	5
6 Over the past month or so, how often have you had to push or strain to begin urination?	0	1	2	3	4	5
7 Over the last month, how many times did you most typically get up to urinate from the time you went to bed at night until the time you got up in the morning?	None	1 time	2 times	3 times	4 times	5 or more times
	0	1	2	3	4	5

Total Symptom Score = \_\_\_\_\_  
Sum of Questions 1 to 7 = \_\_\_\_\_/35

---

**Quality of Life Due to Urinary Symptoms**

1 If you were to spend the rest of your life with your urinary condition just the way it is now, how would you feel about that?

	Delighted	Pleased	Mostly satisfied	Mixed about equally satisfied and dissatisfied	Mostly dissatisfied	Unhappy	Terrible
	0	1	2	3	4	5	6

Quality of Life Assessment Index I. = \_\_\_\_\_


Fig. 1. International prostate symptom score.<sup>[1]</sup>

**VPSS (Visual Prostate Symptom Score)**


© Stellenbosch University

**Patient's Name:** \_\_\_\_\_ **Date:** \_\_\_\_\_  
**Pasiënt se Naam:** \_\_\_\_\_ **Datum:** \_\_\_\_\_


**A:**




**B:**



**C:**



**D:**



**A =** ..... **B =** ..... **C =** ..... **D =** ..... **A + B + C =** .....

Fig. 2. Visual prostate symptom score consisting of pictograms to evaluate (A) force of the urinary stream, (B) daytime frequency, (C) nocturia, and (D) quality of life.

**Table 1. Languages of the study group (N=100)**

	Mother's language* n	Home language n	Able to understand n	Able to speak n	Able to read/write n
Oshiwambo	27	24	36	34	23
Otjiherero	23	23	41	38	12
Afrikaans	16	17	67	67	49
Damara	12	11	20	19	8
Nama	11	11	11	11	4
Himba	3	3	3	3	0
Nyemba	3	3	3	2	2
Portuguese	2	2	3	3	2
Rukwangali	2	1	4	2	1
Shiloi	2	2	3	3	3
Dhemba	1	1	1	1	0
English	1	1	50	50	40
Mbukushu	1	1	1	1	1
Sifwe	1	1	1	1	1
German	0	0	5	5	0

\*Some subjects had a mother who spoke more than one language.

**Table 2. Number of languages understood by members of the study group with the most common mother languages**

Mother's language	Languages understood (N, n (%))				
	1	2	3	4	5
Oshiwambo (N=27)	6 (22.2)	9 (33.3)	8 (29.6)	3 (11.1)	1 (3.7)
Otjiherero (N=23)	11 (47.8)	6 (26.1)	0	5 (21.7)	1 (4.3)
Afrikaans (N=15)	0	10 (66.7)	4 (26.7)	1 (6.7)	0
Damara (N=11)	0	0	4 (36.4)	5 (45.5)	2 (18.2)
Nama (N=9)	0	6 (66.7)	2 (22.2)	1 (11.1)	0

**Table 3. Comparison of patient groups with <5 v. >9 years of schooling**

	Whole group (N=100)	Schooling <5 years (N=32)	Schooling >9 years (N=34)	p-value
Age (years), mean (range)	56.3 (20.1 - 95.4)	71 (38.8 - 95.4)	43.2 (20.1 - 87.5)	<0.0001
Able to read and write, %	70	22	100	<0.0001
Income per month (Namibian \$), mean (range)	3 205 (0 - 30 000)	1 732 (0 - 30 000)	5 926 (0 - 25 000)	<0.0001
IPSS completed without assistance, %	18	0	38	<0.0001
VPSS completed without assistance, %	18	0	44	<0.0001
Time to complete IPSS (seconds), mean (range)	278 (80 - 690)	324 (148 - 690)	294 (80 - 630)	0.366
Time to complete VPSS (seconds), mean (range)	124 (30 - 556)	178 (58 - 556)	107 (30 - 330)	0.003
Correlations				
VPSS total v. IPSS total	r=0.863 p<0.0001	r=0.775 p<0.0001	r=0.673 p<0.0001	
VPSS QA (force of stream) v. IPSS Q5 (weak stream)	r=0.657 p<0.0001	r=0.589 p=0.0007	r=0.295 p=0.09	
VPSS QB (frequency) v. IPSS Q2 (frequency)	r=0.522 p<0.0001	r=0.379 p=0.036	r=0.451 p=0.008	
VPSS QC (nocturia) v. IPSS Q7 (nocturia)	r=0.898 p<0.0001	r=0.844 p<0.0001	r=0.851 p<0.0001	
VPSS QD (QoL) v. IPSS QoL	r=0.806 p<0.0001	r=0.635 p=0.0001	r=0.387 p=0.024	

IPSS = international prostate symptom score; VPSS = visual prostate symptom score; Q = question; QoL = quality of life.

Table 4 shows the correlations between the various VPSS and IPSS questions, Qmax and Qave in the study group as a whole, and Qmax and Qave in patients with VV >150 ml and <150 ml. Comparing the groups with VV >150 ml and <150 ml, there were significant correlations between the total VPSS and total IPSS (0.934 v. 0.741, respectively) and between the VPSS and IPSS questions related to the force of the stream (0.775 v. 0.541) and QoL (0.902 v. 0.692). In the group with VV >150 ml there were significant correlations between Qmax and the IPSS and VPSS questions on the force of the urinary stream (-0.365 v. -0.456) and QoL (-0.284 v. -0.223). In the group with VV <150 ml there were even stronger correlations between Qmax and the IPSS and VPSS questions on the force of the stream (-0.434 v. -0.425) and QoL (-0.501 v. -0.387). The same was true for correlations of Qave with the IPSS and VPSS questions about the force of the urinary stream and QoL.

## Discussion

In a previous study of 96 men with LUTS referred to a urology outpatient clinic, Van der Walt *et al.*<sup>[11]</sup> found that a greater proportion of patients with <7 v. >10 years of schooling could complete the VPSS without assistance compared with the IPSS (68% v. 13% and 92% v. 76%, respectively). There were statistically significant correlations between the total VPSS and IPSS scores, and between the total VPSS, Qmax and Qave. In

**Table 4. Correlations between the visual prostate symptom score, international prostate symptom score and maximum and average urinary flow rates**

Spearman's rank correlation	All follow-up visits (N=155)		VV >150 ml (N=57)		VV <150 ml (N=44)	
	Coefficient (r)	p-value	Coefficient (r)	p-value	Coefficient (r)	p-value
VPSS v. IPSS						
Total	0.863	<0.0001	0.934	<0.0001	0.741	<0.0001
VPSS QA (force of stream) v. IPSS	0.657	<0.0001	0.775	<0.0001	0.541	0.0002
Q5 (weak stream)						
VPSS QB (frequency) v. IPSS	0.522	<0.0001	0.525	<0.0001	0.552	0.0001
Q2 (frequency)						
VPSS QC (nocturia) v. IPSS	0.896	<0.0001	0.913	<0.0001	0.907	<0.0001
Q7 (nocturia)						
VPSS QD (QoL) v. IPSS QoL	0.806	<0.001	0.902	<0.0001	0.692	<0.001
Qmax v. IPSS						
Total	-0.409	<0.0001	-0.284	0.038	-0.371	0.022
Qmax v. IPSS Q2 (frequency)	-0.159	0.121	-0.007	0.962	-0.275	0.095
Qmax v. IPSS Q5 (weak stream)	-0.458	<0.0001	-0.365	0.007	-0.434	0.006
Qmax v. IPSS Q7 (nocturia)	-0.082	0.427	0.142	0.304	-0.186	0.263
Qmax v. IPSS QoL	-0.489	<0.0001	-0.284	0.038	-0.501	0.001
Qmax v. VPSS						
Total	-0.340	0.0007	-0.174	0.208	-0.362	0.026
Qmax v. VPSS QA (force of stream)	-0.522	<0.0001	-0.456	0.0005	-0.425	0.008
Qmax v. VPSS QB (frequency)	-0.109	0.109	-0.005	0.973	-0.246	0.136
Qmax v. VPSS QC (nocturia)	-0.072	0.488	-0.152	0.272	-0.125	0.455
Qmax v. VPSS QD (QoL)	-0.368	0.0002	-0.223	0.105	-0.350	0.031
Qave v. IPSS						
Total	-0.479	<0.0001	-0.441	0.0006	-0.337	0.025
Qave v. IPSS Q2 (frequency)	-0.227	0.020	-0.095	0.481	-0.307	0.043
Qave v. IPSS Q5 (weak stream)	-0.481	<0.0001	-0.507	<0.0001	-0.289	0.057
Qave v. IPSS Q7 (nocturia)	-0.194	0.047	0.016	0.097	-0.287	0.059
Qave v. IPSS QoL	-0.549	<0.0001	-0.437	0.0007	-0.447	0.002
Qave v. VPSS						
Total	-0.425	<0.0001	-0.334	0.011	-0.371	0.013
Qave v. VPSS QA (force of stream)	-0.522	<0.0001	-0.572	<0.0001	-0.413	0.005
Qave v. VPSS QB (frequency)	-0.165	0.109	-0.077	0.570	-0.222	0.147
Qave v. VPSS QC (nocturia)	-0.072	0.488	0.022	0.869	-0.193	0.209
Qave v. VPSS QD (QoL)	-0.440	<0.0001	-0.379	0.004	-0.333	0.027

VPSS = visual prostate symptom score; IPSS = international prostate symptom score; VV = voided volume; Q = question; QoL = quality of life; Qmax = maximum urinary flow rate; Qave = average urinary flow rate.

a further analysis of the uroflowmetry data, Heyns *et al.*<sup>[12]</sup> found that the VPSS question on the subject's assessment of his urinary stream showed a significant negative correlation with Qmax ( $r=-0.37$ ,  $p=0.002$ ) and Qave ( $r=-0.31$ ,  $p=0.011$ ), whereas the IPSS question on the subject's urinary stream did not correlate significantly with Qmax or Qave. In a study of 100 men followed up after a diagnosis of urethral stricture, Wessels and Heyns<sup>[13]</sup> found that the VPSS correlated significantly with the IPSS, Qmax and urethral diameter and took significantly less time to complete than the IPSS.

In the current study, the mean patient age (56.3 years) was considerably lower than in other studies of men with LUTS (64 - 70 years).<sup>[5,9,11]</sup> This is explained by the fact that the underlying pathology was urethral stricture in 61%, with BPH in only 32% of the study cohort.

The majority of the patients (66%) had <10 years of schooling, 25% had never attended school and 30% were illiterate. The patients

with <5 years of schooling were significantly older than those with >9 years of schooling (mean age 71 v. 43.2 years) (Table 3), indicating that the older men grew up in a time when access to school education was less readily available.

The study patients all spoke one or more of 15 languages, so it is not surprising that 82% overall, and 100% in the group with <5 years of schooling, required assistance to complete the IPSS and VPSS. The time taken to complete the VPSS was significantly less than that for the IPSS for the group as a whole (124 v. 278 seconds) and also in the groups with <5 years of schooling (178 v. 324 seconds) v. >9 years of schooling (107 v. 294 seconds), confirming that the VPSS takes less time to complete than the IPSS, especially in men with limited education.<sup>[11]</sup> The mean time to complete both the IPSS and VPSS was shorter at follow-up (173 and 100 seconds) than at the first visit (306 and 227 seconds), indicating that patients learn how to answer the questions more quickly.

The group with lower schooling (<5 years v. >9 years) showed stronger correlations between the total VPSS and total IPSS, between VPSS question A and IPSS question 5 (force of the urinary stream), and between the VPSS and IPSS QoL questions (Table 3). This indicates that the correlations between the VPSS and IPSS scores may be even better in men with very limited schooling than in those with a higher level of schooling.

Although it is generally accepted that Qmax and Qave are only reliable if the VV is >150 ml, in a study of 1 271 men Reynard *et al.*<sup>[16]</sup> found that low-volume uroflowmetry can provide useful diagnostic information and the data from such voids should therefore not be discarded. In their study, those voiding <150 ml had a 72% likelihood of bladder outlet obstruction, while in those voiding >150 ml the likelihood was 56%.<sup>[16]</sup>

Comparing the groups with VV >150 ml v. <150 ml in the current study, there were significant correlations between the total VPSS and total IPSS and between the VPSS and IPSS questions related to the force of the stream and QoL (Table 4). In the group with VV >150 ml there were significant correlations between Qmax and the IPSS and VPSS questions on the force of the urinary stream and QoL. In the group with VV <150 ml there were even stronger correlations between Qmax and the IPSS and VPSS questions on the force of the stream and QoL. The same was true for correlations of Qave with the IPSS and VPSS questions about the force of the urinary stream and QoL (Table 4).

## Conclusion

The VPSS takes significantly less time to complete than the IPSS, especially in men with limited schooling. The VPSS correlates significantly with the IPSS with regard to total score, the questions about the force of the urinary stream, and overall QoL. Since the VPSS pictograms correlate significantly with the corresponding IPSS questions and Qmax and Qave, they can be used as single-item questions to rapidly assess the degree of bladder outflow obstruction in men who are illiterate or have limited education.

**Source of funding,** South African Urological Association.

## References

1. Griffith JW. Self-report measurement of lower urinary tract symptoms: A commentary on the literature since 2011. *Curr Urol Rep* 2012;13(6):420-426. [http://dx.doi.org/10.1007/s11934-012-0286-5]
2. Cam K, BPH: How useful is a visual prostate symptom score for patients? *Nat Rev Urol* 2011;8(10):536-537. [http://dx.doi.org/10.1038/nrurol.2011.137]
3. Cam K, Senel F, Akman Y, Erol A. The efficacy of an abbreviated model of the International Prostate Symptom Score in evaluating benign prostatic hyperplasia. *BJU Int* 2003;91(3):186-189. [http://dx.doi.org/10.1046/j.1464-410X.2003.04055.x]
4. Oztürk Mİ, Koca O, Keleş MO, Güneş M, Kaya C, Karaman Mİ. International prostate symptom score: Really appreciated by all patients or not? *Urol J* 2011;8(3):227-230.
5. Ogwuche EI, Dakum NK, Amu CO, Dung ED, Udeh E, Ramyil VM. Problems with administration of international prostate symptom score in a developing community. *Ann Afr Med* 2013;12(3):171-173. [http://dx.doi.org/10.4103/1596-3519.117628]
6. Kim JH, Doo SW, Yang WJ, Song YS. Homogeneity among the Korean international prostate symptom score questionnaires used in real practice. *Korean J Urol* 2013;54(4):249-251. [http://dx.doi.org/10.4111/kju.2013.54.4.249]
7. Fujimura T, Kume H, Nishimatsu H, et al. Assessment of lower urinary tract symptoms in men by international prostate symptom score and core lower urinary tract symptom score. *BJU Int* 2012;109(10):1512-1516. [http://dx.doi.org/10.1111/j.1464-410X.2011.10445.x]
8. Coyne KS, Barsdorf AI, Thompson C, et al. Moving towards a comprehensive assessment of lower urinary tract symptoms (LUTS). *Neurourol Urodyn* 2012;31(4):448-454. [http://dx.doi.org/10.1002/nau.21202]
9. O'Leary MP, Wei JT, Roehrborn CG, Miner M; BPH Registry and Patient Survey Steering Committee. Correlation of the International Prostate Symptom Score bother question with the Benign Prostatic Hyperplasia Impact Index in a clinical practice setting. *BJU Int* 2008;101(12):1531-1535. [http://dx.doi.org/10.1111/j.1464-410X.2008.07574.x]
10. Viktrup L, Hayes RP, Wang P, Shen W. Construct validation of patient global impression of severity (PGI-S) and improvement (PGI-I) questionnaires in the treatment of men with lower urinary tract symptoms secondary to benign prostatic hyperplasia. *BMC Urol* 2012;12(1):30. [http://dx.doi.org/10.1186/1471-2490-12-30]
11. Van der Walt CL, Heyns CF, Groeneveld AE, Edlin RS, van Vuuren SP. Prospective comparison of a new visual prostate symptom score versus the international prostate symptom score in men with lower urinary tract symptoms. *Urology* 2011;78(1):17-20. [http://dx.doi.org/10.1016/j.urolgy.2011.01.065]
12. Heyns CF, Van der Walt CL, Groeneveld AE. Correlation between a new visual prostate symptom score (VPSS) and uroflowmetry parameters in men with lower urinary tract symptoms. *S Afr Med J* 2012;102(4):237-240.
13. Wessels SG, Heyns CF. Prospective evaluation of a new visual prostate symptom score, the international prostate symptom score, and uroflowmetry in men with urethral stricture disease. *Urology* 2014;83(1):220-224. [http://dx.doi.org/10.1016/j.urolgy.2013.08.058]
14. Peeling WB. Diagnostic assessment of benign prostatic hyperplasia. *Prostate Suppl* 1989;2:51-68.
15. Ushijima S, Ukimura O, Okihara K, Mizutani Y, Kawachi A, Miki T. Visual analog scale questionnaire to assess quality of life specific to each symptom of the International Prostate Symptom Score. *J Urol* 2006;176(2):665-671. [http://dx.doi.org/10.1016/j.juro.2006.03.031]
16. Reynard JM, Yang Q, Donovan JL, et al. The ICS-'BPH' Study: Uroflowmetry, lower urinary tract symptoms and bladder outlet obstruction. *Br J Urol* 1998;82(5):619-623. [http://dx.doi.org/10.1046/j.1464-410X.1998.00813.x]

Accepted 18 February 2014.