



REVIEW

The current global situation for tuberculous meningitis: epidemiology, diagnostics, treatment and outcomes [version 1; peer review: 2 approved]

James A Seddon ^{1,2}, Lillian Tugume ³, Regan Solomons ⁴,
Kameshwar Prasad⁵, Nathan C Bahr ⁶,
Tuberculous Meningitis International Research Consortium

¹Department of Infectious Diseases, Imperial College London, London, W2 1PG, UK

²Desmond Tutu TB Centre, Department of Paediatrics and Child Health, Stellenbosch University, Cape Town, South Africa

³Infectious Diseases Institute, Makerere University, Kampala, Uganda

⁴Department of Paediatrics and Child Health, Stellenbosch University, Cape Town, South Africa

⁵Department of Neurology, All India Institute of Medical Sciences, New Delhi, India

⁶Department of Infectious Diseases, University of Kansas, Kansas City, KS, USA

V1 First published: 05 Nov 2019, 4:167
<https://doi.org/10.12688/wellcomeopenres.15535.1>
Latest published: 05 Nov 2019, 4:167
<https://doi.org/10.12688/wellcomeopenres.15535.1>

Abstract

Tuberculous meningitis (TBM) results from dissemination of *M. tuberculosis* to the cerebrospinal fluid (CSF) and meninges. Ischaemia, hydrocephalus and raised intracranial pressure frequently result, leading to extensive brain injury and neurodisability. The global burden of TBM is unclear and it is likely that many cases are undiagnosed, with many treated cases unreported. Untreated, TBM is uniformly fatal, and even if treated, mortality and morbidity are high. Young age and human immunodeficiency virus (HIV) infection are potent risk factors for TBM, while Bacillus Calmette–Guérin (BCG) vaccination is protective, particularly in young children. Diagnosis of TBM usually relies on characteristic clinical symptoms and signs, together with consistent neuroimaging and CSF parameters. The ability to confirm the TBM diagnosis via CSF isolation of *M. tuberculosis* depends on the type of diagnostic tests available. In most cases, the diagnosis remains unconfirmed. GeneXpert MTB/RIF and the next generation Xpert Ultra offer improved sensitivity and rapid turnaround times, and while roll-out has scaled up, availability remains limited. Many locations rely only on acid fast bacilli smear, which is insensitive. Treatment regimens for TBM are based on evidence for pulmonary tuberculosis treatment, with little consideration to CSF penetration or mode of drug action required. The World Health Organization recommends a 12-month treatment course, although data on which to base this duration is lacking. New treatment regimens and drug dosages are under evaluation, with

Open Peer Review

Reviewer Status

Invited Reviewers

1

2

version 1

05 Nov 2019

report

report


1. **Kiran T. Thakur**, Columbia University Irving Medical Center, New York City, USA
2. **Tamilarasu Kadhiravan** , Jawaharlal Institute of Postgraduate Medical Education and Research, Puducherry, India

Any reports and responses or comments on the article can be found at the end of the article.

much higher dosages of rifampicin and the inclusion of fluoroquinolones and linezolid identified as promising innovations. The inclusion of corticosteroids at the start of treatment has been demonstrated to reduce mortality in HIV-negative individuals but whether they are universally beneficial is unclear. Other host-directed therapies show promise but evidence for widespread use is lacking. Finally, the management of TBM within health systems is sub-optimal, with drop-offs at every stage in the care cascade.

Keywords

Tuberculosis, Meningitis, Tuberculous meningitis, diagnosis, TBM



This article is included in the [Tuberculous Meningitis International Research Consortium](#) collection.

Corresponding author: James A Seddon (james.seddon@imperial.ac.uk)

Author roles: **Seddon JA:** Conceptualization, Writing – Original Draft Preparation, Writing – Review & Editing; **Tugume L:** Conceptualization, Writing – Original Draft Preparation, Writing – Review & Editing; **Solomons R:** Conceptualization, Writing – Original Draft Preparation, Writing – Review & Editing; **Prasad K:** Conceptualization, Writing – Original Draft Preparation, Writing – Review & Editing; **Bahr NC:** Conceptualization, Writing – Original Draft Preparation, Writing – Review & Editing;

Competing interests: No competing interests were disclosed.

Grant information: This work was supported by the Wellcome Trust through funding to the Tuberculous Meningitis International Research Consortium. JAS is supported by a Clinician Scientist Fellowship jointly funded by the UK Medical Research Council (MRC) and the UK Department for International Development (DFID) under the MRC/DFID Concordat agreement [MR/R007942/1]. RS is supported by the National Research Foundation of South Africa [109437].

Copyright: © 2019 Seddon JA *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Seddon JA, Tugume L, Solomons R *et al.* **The current global situation for tuberculous meningitis: epidemiology, diagnostics, treatment and outcomes [version 1; peer review: 2 approved]** Wellcome Open Research 2019, 4:167 <https://doi.org/10.12688/wellcomeopenres.15535.1>

First published: 05 Nov 2019, 4:167 <https://doi.org/10.12688/wellcomeopenres.15535.1>

Introduction

Tuberculous meningitis (TBM) is an extra-pulmonary form of tuberculosis (TB) characterised by sub-acute or chronic inflammation of the meninges as a result of invasion of the sub-arachnoid space by the bacilli *M. tuberculosis*. Other forms of central nervous system (CNS) TB such as tuberculoma, cerebral abscess, and spinal TB are not categorised as TBM, even though treatment is similar. TBM most commonly affects young children and individuals with human immunodeficiency virus (HIV). In the absence of TB treatment, TBM is uniformly fatal¹ and even with treatment, outcomes are poor and chronic neurodisability is common. Much of the damage caused by TBM is due to host-derived inflammatory responses, yet our understanding of these processes is limited. In this review we describe the current global situation for TBM in terms of our understanding of the pathogenesis and natural history, epidemiology, diagnosis, and treatment. We also explore how TBM is managed within health systems. Challenges in the field of TBM are shown in [Table 1](#).

Pathogenesis

Primary infection occurs via inhalation of *M. tuberculosis*-containing aerosolised droplets, followed by activation of

neutrophils, dendritic cells and alveolar macrophages, which engulf the mycobacteria in the terminal alveoli. Infected cells then migrate to lymphoid tissue, resulting in activation of Th1 cells and production of pro-inflammatory cytokines, with resultant inflammatory changes in the lung parenchyma and vasculature². If the mycobacteria reach the vasculature, haematogenous dissemination can occur, with the potential to invade the CNS³⁻⁵.

The pathogenesis of TBM continues to be debated. A key virulence feature of mycobacteria is the ability to invade the blood-brain and blood-cerebrospinal fluid (CSF) barriers. The invasion mechanisms are not clear, although *in vitro* and animal data suggest that *M. tuberculosis* may rearrange the actin of the layers^{4,6}. It is also possible that a “Trojan horse” mechanism, by which *M. tuberculosis* is brought across the blood-brain barrier by infected macrophages and neutrophils, may occur⁷. Rich and McCordock described the development of a caseating “Rich” focus in the context of TBM pathogenesis^{8,9}. It is suggested that the Rich focus is formed via activation of microglial cells and astrocytes once the bacilli have gained access to the brain. Once formed, the Rich foci may become activated rapidly or months to years later, resulting in release of

Table 1. Challenges in tuberculous meningitis.

	Challenges	Implication
Pathogenesis	Mechanism of dissemination from lungs to CSF unclear Mechanisms of CSF invasion unclear	Possible preventative steps/interventions are unclear
Natural History	Why some individuals develop TBM unclear	Focussed prevention therapy not possible
Epidemiology	No good estimates of global TBM incidence No good estimates of overall mortality/morbidity Unclear differentiation between adult and childhood TBM	Inadequate research funding, inadequate local, country and international health system investment in treatment and diagnosis of TB meningitis Management tailored for children/adults
Diagnosis	Inadequate ability to rule-out TBM Incomplete market penetration of the most effective tests Relatively expensive tests	Many missed cases of TBM Late diagnosis and potentially preventable long-term neurologic disability or mortality Testing unavailable in some settings due to cost
Treatment	Pulmonary TB treatment regimens adapted (with minimal change) to TBM Incomplete understanding of the best treatment for host immune inflammation	Likely inadequate TBM treatment regimens Uncertainty regarding dosing, optimal anti-tuberculous medications Uncertain efficacy/harm of corticosteroids in some populations
Health Systems	Poor community awareness Inadequate diagnostic device access	Missed diagnoses Inadequate resources committed to TBM Missed diagnoses and poor outcomes

CSF, cerebrospinal fluid; TBM, tuberculous meningitis; TB, tuberculosis.

M. tuberculosis into the subarachnoid space, triggering an inflammatory cascade⁹. The resulting inflammatory changes may explain some of the clinical features associated with TBM. First, peri-vascular inflammation, particularly of the middle cerebral artery, results in decreased perfusion and cerebral infarct. Second, extension of exudative material to the basal cisterns and the mid-brain leads to disruption of CSF flow, hydrocephalus and raised intracranial pressure. Third, exudates encase cranial nerves, resulting in cranial nerve palsies. Finally, expanding parenchymal tubercles may form tuberculomas and, less frequently, brain abscesses. In contrast (or in addition) to the Rich hypothesis, other researchers have suggested that bacilli reach the CSF during miliary dissemination. Donald and colleagues have proposed a pathogenic mechanism, based on more recent clinical, post mortem and epidemiological data, that articulates the central role of the Rich focus but with an additional component that includes miliary TB⁵.

Natural history

The risk and severity of TBM are altered by the status of the host immune response and pathogen virulence, consistent with the damage response framework¹⁰. Multiple host factors such as age, HIV co-infection, Bacillus Calmette–Guérin (BCG) immunisation, malnutrition, and helminth co-infection may lead to either a deficient or exaggerated immune response to mycobacterial infection.

Age

Several studies conducted in the pre-chemotherapy era identified children who had been exposed to TB and followed them for tuberculin skin test conversion, chest radiograph changes, clinical progression, and mycobacterial culture positivity. Marais and colleagues reviewed these studies and proposed a timetable for TB, in the absence of drug therapy¹¹. Following infection with *M. tuberculosis*, young children are at high risk of progression to TB disease including disseminated forms such as TBM. Both the risk of disease progression and risk of dissemination fall rapidly through childhood, reaching a nadir in the primary school age. In more recent cohorts of children with TBM, the median age is between two and four years¹². Marais and colleagues also found that children who develop miliary TB or TBM generally do so from one to four months after exposure.

HIV

HIV co-infection is the most significant risk factor for TBM in adults and is believed to attenuate the host immune response in the CSF. The impact of HIV on the clinical presentation of TBM is debated, with some studies suggesting a higher rate of extra-meningeal disease and of TB drug resistance^{13–16}. Other studies report that the clinical course of TBM is unaltered by HIV infection^{17,18}.

BCG, helminth infections and nutrition

Neonatal BCG immunization is effective in preventing childhood TB, particularly TBM and miliary TB^{19–21}. Efficacy varies from as high as 80% near the poles, to <20% near the

equator, possibly due to masking or blocking from environmental mycobacteria or modulation due to helminth infections^{20,22,23}. Helminth infections activate a T helper type 2 and/or regulatory T cell response and may down-regulate T helper type 1 and T helper type 17 responses to mycobacterial antigens, potentially decreasing BCG vaccine immunogenicity. Whether this is clinically important is unclear^{24–31}. Several epidemiological, clinical and laboratory studies have demonstrated that malnutrition, a form of acquired immunodeficiency, increases susceptibility to TB. In particular, vitamin D and vitamin D receptor polymorphisms are critical for function of macrophages in the context of TBM^{32–34}. Vitamin D deficiency was significantly more common in TBM compared to controls and pulmonary TB in an Indian study³⁵.

Strain type

There are seven main *M. tuberculosis* lineages, each associated with varying degrees of proinflammatory host response. The modern lineage x (Beijing) is associated with disseminated extra-pulmonary disease, drug resistance, and possibly poor treatment outcomes^{36–39}. One plausible mechanism of Beijing strain survival is a dampened IFN- γ host inflammatory response, promoting high bacterial load.

Epidemiology

Our understanding of the global burden of TBM is poor. Because of inadequate diagnostic test performance and availability, many cases of TBM remain undiagnosed. Autopsy studies in adults with HIV have found high proportions of TB co-infection, commonly with extrapulmonary disease, including meningitis, which is frequently undiagnosed and untreated prior to death⁴⁰. Even those cases diagnosed may not be appropriately reported as individuals with TBM are usually diagnosed in hospitals, which in some contexts, are less likely to be reporting units for TB⁴¹.

The proportion of TBM in cohorts of TB varies dramatically by the local TB prevalence, age, HIV prevalence, and whether the cohort is describing all or only hospitalized persons (which tend to have a higher proportion of TBM cases). A population-based estimate in a low TB-prevalence settings suggested that around 1% of TB cases were TBM⁴², while a paediatric hospital-based cohort of confirmed TB in a high TB-burden setting suggested that over 10% of TB cases were TBM⁴³. Given that the World Health Organization (WHO) estimates 10.4 million new TB cases each year (of which 1 million are children), we suggest that at least 100,000 individuals develop TBM annually, but this figure may be much higher.

Untreated, TBM is uniformly fatal. A study from the pre-chemotherapy literature in children demonstrated that the median time to death was 19 and a half days⁴⁴. Even if treated, up to 20% of children will die and, of survivors, over half have severe neurodevelopmental sequelae⁴⁵. Mortality in HIV-infected adults treated for TBM can be as high as 60% in some cohorts, dependent on degree of immunosuppression, use of antiretroviral therapy and stage of TBM at presentation^{14,46}. The

total mortality impact of this condition is unclear, and likely underestimated.

Morbidity in TBM survivors is heavily dependent on the age at which the disease occurs and the context in which the individual is living. Survivors require extensive support from medical staff, physiotherapy, occupational therapy, pharmacists and dieticians. Many of these services are not widely available in high TB-burden settings. They also require vast commitments from family and community members, with implications for their quality of life and economic circumstances. Given that they have more potential life years ahead, children lose more disability adjusted life years than older individuals. Even in children with limited neuromotor disability following treatment, cognitive disability, attention-deficit and behavioural problems are more common with TBM survivors than with community controls^{47,48}.

Diagnosics

TBM remains difficult to diagnose. Clinical features such as fever, headache, neck stiffness, or a 'typical' CSF pattern (elevated protein, lymphocytic pleocytosis, and low glucose) cannot reliably distinguish TBM from other forms of sub-acute meningitis^{49,50}. Neuroimaging is frequently unavailable in high TB burden contexts. Where characteristics features of TBM are present on neuroimaging, combinations of basal enhancement, tuberculomas, infarction or hydrocephalus support (but do not confirm) the diagnosis. Many cases of TBM, however, are associated with neuroimaging that does not assist with the diagnosis. Traditional tests to detect bacilli include acid fast bacilli (AFB) smear and culture. AFB smear is rapid and widely available but only 10–15% sensitive in most settings^{49,51}. Culture is more sensitive (~50–60%) but results return too slowly to allow for clinical intervention (two-six weeks depending on the media)^{50–53}. Death frequently occurs prior to culture results if empiric therapy has not been initiated in the interim period⁵⁰. Culture also requires a biosafety level three laboratory, limiting use to referral centres in many settings.

In recent years, nucleic acid amplification tests (NAATs) have been utilised for TBM diagnosis. The most notable tests are GeneXpert MTB/RIF (Xpert) and the re-engineered GeneXpert MTB/RIF Ultra (Xpert Ultra), though other commercial and 'in-house' NAATs have been used as well⁵⁴. Xpert and Xpert Ultra (Cepheid, Inc, Sunnyvale, CA, USA) are cartridge-based, rapid, fully automated polymerase chain reaction (PCR) tests. In 2013, the WHO recommended Xpert as the initial test for diagnosis of TBM based on a systematic review of 13 studies^{55–57}. Sensitivity is generally similar to culture (50–60%), but with a run-time of two hours^{51,53,58}. Xpert and culture frequently detect cases that the other modality misses and Xpert's negative predictive value is only 84–94%, meaning that while Xpert is helpful if positive, it cannot effectively rule out TBM^{51,59,60}. One way to maximise Xpert's utility is to centrifuge larger volumes of CSF (>5ml), although this effect has not been universally observed⁵¹. At present, access to Xpert is increasing although still not adequate⁶¹.

Xpert Ultra utilizes the same instrument as Xpert with a software upgrade. WHO has recommended Xpert Ultra in place of Xpert for TB meningitis based on one study of Ugandan HIV-infected adults with suspected TBM, which found 95% sensitivity versus a composite microbiological end point, and 70% sensitivity versus consensus research case definitions^{52,62}. This was as opposed to inferior sensitivities for both Xpert and culture versus the same reference standards^{52,63}. A recent cohort reported 86% sensitivity for Xpert Ultra for definite TBM versus 36% for Xpert and 14% for culture⁶⁴. Negative predictive values remain inadequate to 'rule-out' TBM and the publication of additional studies of Xpert Ultra is required to fully evaluate this technology.

Although there is hope that immunodiagnostics will bridge the gap from Ultra to a definitive diagnostic tool-kit for TBM, thus far, adenosine deaminase, INF- γ release assays, and antibody tests, among others, have not yet proven adequate adjunctive tests for TBM^{54,65}. Thus, diagnostic tests remain imperfect (though improving) and a low threshold to start empiric therapy remains a key to effective management of TB meningitis.

Treatment

The evidence to guide treatment in TBM is limited, based mainly on experience with pulmonary TB⁶⁶. The first treatment option for TBM, streptomycin became available in 1946^{67,68}, albeit at the cost of long, painful treatment duration and poor survival^{69–75}. Para-amino salicylic acid (PAS), introduced in the early 1950s, given in combination with streptomycin reduced resistance and mortality, even with poor CSF penetration^{72,76–79}. Isoniazid, with low toxicity and better CNS penetration, was introduced in 1952, whereafter outcomes improved^{76,77,80–83}. Rifampicin and pyrazinamide were used in TBM treatment regimens from 1970 onwards; however, initially there was no significant reduction in mortality compared to regimens containing streptomycin, PAS and isoniazid^{76,84–87}.

Available treatment options for TBM are hampered by poor CSF penetration and toxicity^{76,88}. Of the first-line TB drugs, pharmacokinetic studies have shown that isoniazid and pyrazinamide have the best CSF penetration⁸⁸. In children, modelling suggests that higher dosages of rifampicin may be required⁸⁹, while in adults, doses up to 600mg per day increases CSF penetration^{88,90}. Ethambutol and streptomycin both have poor CSF penetration. Second-line TB drugs with good CSF penetration include ethionamide, the fluoroquinolones (levofloxacin, moxifloxacin and ofloxacin) and terizidone⁸⁸. Linezolid is another potential treatment option, given excellent CSF penetration^{76,91}. Moxifloxacin up to 800mg per day has demonstrated good CSF concentration with low toxicity^{90,92}.

Current WHO treatment guidelines for TBM recommend two months of a four-drug regimen (rifampicin, isoniazid, pyrazinamide and ethambutol) in both adults and children, followed by 7–10 months of rifampicin and isoniazid^{93,94}. Similar dosing to pulmonary TB is advocated based on low quality evidence (Table 2)^{88,95}. Thus, the focus of recent TBM trials has

Table 2. World Health Organization recommended drug dosing for adults and children with tuberculous meningitis^{93,94}.

Drug	Adult dosage and range (mg/kg)	Childhood dosage and range (mg/kg)	Maximum dosage (mg/day)
Isoniazid (H)	5 (4-6)	10 (10-15)	300
Rifampicin (R)	10 (8-12)	15 (10-20)	600
Pyrazinamide (Z)	25 (20-30)	35 (30-40)	
Ethambutol (E)	15 (15-20)	20 (15-25)	

been to optimize CNS penetration while minimizing toxicity using alternative regimens, higher doses and/or shorter treatment. Adjunctive levofloxacin (20mg/kg per day) did not demonstrate increased survival benefit among adults in one Vietnamese randomized controlled trial⁹⁶⁻⁹⁸. Regarding dosing strategies, high-dose intravenous rifampicin for the first two weeks of treatment demonstrated increased CSF drug concentration as well as significant reduction in six-month mortality^{90,99,100}. Similarly, a study evaluating an abbreviated course (six months in HIV-uninfected and nine months in HIV-infected children) of a high dose four-drug regimen (rifampicin 20mg/kg, isoniazid 20mg/kg, pyrazinamide 40mg/kg and ethionamide 20mg/kg) demonstrated mortality of less than 5%¹⁰¹.

Adjuvant immunomodulatory treatment is used in TBM to ameliorate the potentially damaging host response. A 2016 Cochrane systematic review and meta-analysis found that adjunctive corticosteroids reduce mortality in the short term but had no effect on long-term neurological disability in HIV-uninfected patients with TBM; however, the benefit of corticosteroids is unclear in HIV co-infected individuals¹⁰². Corticosteroids may have a differential effect dependent on leukotriene A4 hydrolase genotype, being beneficial in those with a hyper-inflammatory genotype but detrimental in others. Use of adjunctive aspirin has been shown to be beneficial in adults with TBM^{103,104}, with one recent study of high dose aspirin demonstrating prevention of new cerebral infarction in adults¹⁰⁵. This effect, however, has not been demonstrated in children and requires larger randomised trials in adults¹⁰⁶. Currently registered clinical trials for TBM are shown in [Table 3](#).

TB mass lesions, usually in the setting of immune reconstitution inflammatory syndrome (IRIS) and optochiasmatic arachnoiditis, are difficult to treat and often occur near vital structures in the brainstem and spinal cord. Thalidomide, a potent TNF- α inhibitor, has demonstrated clinical benefit in both TB mass lesions and optochiasmatic arachnoiditis^{107,108}. Reports have also demonstrated a clinical role for other agents that cause TNF- α inhibition, including infliximab and IFN- γ ¹⁰⁹⁻¹¹¹. Prophylactic pyridoxine (5–10 mg per day in children and 10 mg per day in adults) is recommended to prevent isoniazid related peripheral neuropathy^{93,94}. Neuropathy is more severe during pregnancy, HIV co-infection, alcohol dependency, malnutrition, diabetes,

chronic liver disease and renal failure. Treatment of hydrocephalus depends on the level of CSF obstruction. Communicating hydrocephalus, a common complication of TBM, can be successfully treated with one month of acetazolamide 50 mg/kg per day and frusemide 1 mg/kg per day^{112,113}.

Health systems

The care cascade for TBM has not been specifically investigated. For TB more generally, large gaps exist at every stage from incidence to diagnosis, from diagnosis to treatment and from treatment initiation to favourable outcome¹¹⁴⁻¹¹⁶. However, there are specific challenges for health systems in their management of TBM, including rehabilitation following treatment conclusion, that suggest gaps could be even larger than for TB as a whole.

Symptoms of TBM are non-specific and community sensitisation is poor. Even where sensitisation to TB has occurred, this is usually to the symptoms and signs of pulmonary TB, such as cough, weight loss, night sweats and haemoptysis¹¹⁷. If individuals with TBM do present for evaluation, it is commonly to primary care facilities, at which staff training and experience can be limited and appropriate investigations, such as lumbar puncture or cerebral imaging, are often unavailable. Multiple presentations to healthcare workers are common and are associated with delayed diagnosis and treatment initiation^{118,119}. Referral from primary care to hospital can lead to further delays and frequently incur costs to patients and families. These delays can lead to clinical deterioration¹²⁰.

Even when an individual presents to a hospital for evaluation, the availability of Xpert is variable globally⁶¹. Although AFB smear is widely available, it has limited sensitivity and culture, if available, returns results in a timescale that is not clinically useful. Although drug therapy is usually provided freely by national TB programmes and is available at peripheral care settings, the specialist services required to appropriately manage TBM, such as paediatrics, neurology, neurosurgery and rehabilitation, are usually only available in specialist centres in larger urban areas and frequently incur costs for patients.

Surveillance data for TBM is limited, with cases rarely reported if patients die before starting treatment. Many patients with TBM are cared for in hospitals for most of their illness and given that many hospitals are not TB-reporting units, these cases are not reflected in regional, national or global reports.

Table 3. Currently open and registered clinical trials in tuberculous meningitis (from the ClinicalTrials.gov and ISRCTN registries).

Trial Name	Registration Number	Countries Recruiting	Target Patient Group	Study Arms	Primary Outcome	Sample Size
Optimizing Anti-tuberculosis Therapy in Adults with Tuberculous Meningitis	NCT03787940	China	Adults 18–65 years	Standard dose vs. high dose isoniazid for rapid NAT2 acetylators	Death or severe disability at 12 months from enrolment	338 rapid acetylators
Retrospective Real-word Study of Linezolid for the Treatment of Tuberculous Meningitis	NCT03898635	China	Adults 18 years and older	<ol style="list-style-type: none"> No linezolid in treatment regimen at any point Linezolid added during the treatment course Linezolid in treatment regimen from beginning 	Survival within five years	n/s
Linezolid, Aspirin and Enhanced Dose Rifampicin in HIV-TBM (LASER-TBM)	NCT03927313	South Africa	HIV-infected adults (>18 years)	<ol style="list-style-type: none"> Standard of care drug regimen at standard of care dosages Higher dosages of rifampicin and the addition of linezolid for the first 56 days of treatment Higher dosages of rifampicin and the addition of linezolid and aspirin for the first 56 days of treatment 	Treatment related adverse events	100
Adjunctive Corticosteroids for Tuberculous Meningitis in HIV-infected Adults (The ACT HIV Trial)	NCT03092817	Vietnam, Indonesia	HIV-infected adults (>18 years)	Dexamethasone vs. placebo	Survival 12 months post randomisation	520
Leukotriene A4 Hydrolase Stratified Trial of Adjunctive Corticosteroids for HIV-uninfected Adults with Tuberculous Meningitis	NCT03100786	Vietnam	HIV-uninfected adults (>18 years)	<ol style="list-style-type: none"> Open label dexamethasone for all patients with TT genotype for first 6–8 weeks treatment Individuals with CT or CC genotypes randomised to dexamethasone or placebo for first 6–8 weeks treatment 	All-cause mortality or new neurological event by 12 months post randomisation	640
Optimizing Treatment to Improve TBM Outcomes in Children (TBM-KIDS)	NCT02958709	India, Malawi	Children	<ol style="list-style-type: none"> High dose rifampicin in a standard drug regimen High dose rifampicin and levofloxacin instead of ethambutol Standard dose rifampicin in a standard drug regimen 	Characterise the pharmacokinetic parameters of rifampicin and levofloxacin and describe outcomes and safety	120

Trial Name	Registration Number	Countries Recruiting	Target Patient Group	Study Arms	Primary Outcome	Sample Size
Harvest trial - improving outcomes from TB meningitis with high dose oral rifampicin	ISRCTN15668391	Uganda, South Africa, Indonesia	Adults (>18 years)	Standardly dosed regimen compared to a standard drug regimen but with elevated rifampicin dosage	Six-month survival	500
SURE: Short intensive treatment for children with tuberculous meningitis	ISRCTN40829906	India, Uganda, Vietnam, Zambia, Zimbabwe	Children under 15 years (and over 28 days)	Factorial design: <ul style="list-style-type: none"> First randomisation: standard drug regimen vs. drug regimen with high dose rifampicin and substitution of ethambutol with levofloxacin Second randomisation: aspirin vs. placebo 	First randomisation: all-cause mortality by 48 weeks Second randomisation: neurodevelopment at 48 weeks assessed using a Modified Rankin Score	400
Improving diagnosis and treatment of HIV-associated Tuberculous meningitis	ISRCTN42218549	Uganda	Adults (>18 years)	<ol style="list-style-type: none"> Intravenous 20mg/kg/day rifampicin for two weeks (followed by oral rifampicin 35mg/kg/day for six weeks) Oral 35mg/kg/day rifampicin for eight weeks Standard of care oral rifampicin (~10mg/kg/day) for eight weeks 	Pharmacokinetic parameters and safety composite endpoint	60

Disaggregation of data occurs to the level of pulmonary TB and extrapulmonary TB, without specifying the site of disease. Some electronic registers allow the recording of International Classification of Diseases 10th revision (ICD-10) codes, but these are often poorly completed.

Conclusions

TBM is the most severe form of TB and is associated with high mortality and morbidity. Our understanding of the global burden is limited but it is likely that most cases are not diagnosed and appropriately treated. At least 100,000 cases are likely to occur each year. The current diagnostic tools are largely inadequate, and treatment is usually started only once substantial neurological damage has occurred. Treatment, both anti-mycobacterial and anti-inflammatory, require further investigation and optimisation, and improvements at every stage along the care cascade are urgently needed.

Data availability

No data are associated with this article.

Acknowledgements

Tuberculous Meningitis International Research Consortium

Rob E. Aarnoutse; Suzanne T. B. Anderson; Nathan C. Bahr; Nguyen D. Bang; David R. Boulware; Tom Boyles; Lindsey H. M. te Brake; Satish Chandra; Felicia C. Chow; Fiona V. Cresswell; Reinout van Crevel; Angharad G. Davis; Sofiati Dian; Joseph Donovan; Kelly E. Dooley; Anthony Figaji; A. Rizal Ganiem; Ravindra Kumar Garg; Diana M. Gibb; Raph L. Hamers; Nguyen T. T. Hiep; Darma Imran; Akhmad Imron; Sanjay K. Jain; Sunil K. Jain; Byramee Jeejeebhoy; Jayantee Kalita; Rashmi Kumar; Vinod Kumar; Arjan van Laarhoven; Rachel P-J. Lai; Abi Manesh; Suzaan Marais; Vidya Mave; Graeme Meintjes; David B. Meyers; Usha K. Misra; Manish Modi; Alvaro A. Ordonez; Nguyen H. Phu; Sunil Pradhan; Kameshwar Prasad; Alize M. Proust; Lalita Ramakrishnan; Ursula Rohlwick; Rovina Ruslami; Johannes F. Schoeman; James A. Seddon; Kusum Sharma; Omar Siddiqi; Regan S. Solomons; Nguyen T. T. Thuong; Guy E. Thwaites; Ronald van Toorn; Elizabeth W. Tucker; Sean A. Wasserman; Robert J. Wilkinson.

References

- Jullien S, Ryan H, Modi M, *et al.*: **Six months therapy for tuberculous meningitis.** *Cochrane Database Syst Rev.* 2016; 9: CD012091. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- O'Garra A, Redford PS, McNab FW, *et al.*: **The immune response in tuberculosis.** *Annu Rev Immunol.* 2013; 31: 475–527. [PubMed Abstract](#) | [Publisher Full Text](#)
- Krishnan N, Robertson BD, Thwaites G: **The mechanisms and consequences of the extra-pulmonary dissemination of *Mycobacterium tuberculosis*.** *Tuberculosis (Edinb).* 2010; 90(6): 361–366. [PubMed Abstract](#) | [Publisher Full Text](#)
- Jain SK, Paul-Satyaseela M, Lamichhane G, *et al.*: ***Mycobacterium tuberculosis* invasion and traversal across an *in vitro* human blood-brain barrier as a pathogenic mechanism for central nervous system tuberculosis.** *J Infect Dis.* 2006; 193(9): 1287–1295. [PubMed Abstract](#) | [Publisher Full Text](#)
- Donald PR, Schaaf HS, Schoeman JF: **Tuberculous meningitis and miliary tuberculosis: the Rich focus revisited.** *J Infect.* 2005; 50(3): 193–195. [PubMed Abstract](#) | [Publisher Full Text](#)
- Be NA, Kim KS, Bishai WR, *et al.*: **Pathogenesis of central nervous system tuberculosis.** *Curr Mol Med.* 2009; 9(2): 94–99. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Nguyen L, Pieters J: **The Trojan horse: survival tactics of pathogenic mycobacteria in macrophages.** *Trends Cell Biol.* 2005; 15(5): 269–276. [PubMed Abstract](#) | [Publisher Full Text](#)
- Hektoen L: **The Vascular Changes of Tuberculous Meningitis, Especially the Tuberculous Endarteritis.** *J Exp Med.* 1896; 1(1): 112–163. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Rich AR, McCordock HA: **The pathogenesis of tuberculous meningitis.** *Bull Johns Hopkins Hospital.* 1933; 52: 5–37.
- Pirofski LA, Casadevall A: **Immune-Mediated Damage Completes the Parabola: *Cryptococcus neoformans* Pathogenesis Can Reflect the Outcome of a Weak or Strong Immune Response.** *mBio.* 2017; 8(6): pii: e02063–17. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Marais BJ, Gie RP, Schaaf HS, *et al.*: **The natural history of childhood intrathoracic tuberculosis: a critical review of literature from the pre-chemotherapy era.** *Int J Tuberc Lung Dis.* 2004; 8(4): 392–402. [PubMed Abstract](#)
- van Well GT, Paes BF, Terwee CB, *et al.*: **Twenty years of pediatric tuberculous meningitis: a retrospective cohort study in the western cape of South Africa.** *Pediatrics.* 2009; 123(1): e1–8. [PubMed Abstract](#) | [Publisher Full Text](#)
- Torok ME, Chau TT, Mai PP, *et al.*: **Clinical and microbiological features of HIV-associated tuberculous meningitis in Vietnamese adults.** *PLoS One.* 2008; 3(3): e1772. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Thwaites GE, Duc Bang N, Huy Dung N, *et al.*: **The influence of HIV infection on clinical presentation, response to treatment, and outcome in adults with Tuberculous meningitis.** *J Infect Dis.* 2005; 192(12): 2134–2141. [PubMed Abstract](#) | [Publisher Full Text](#)
- Karstaedt AS, Valtchanova S, Barriere R, *et al.*: **Tuberculous meningitis in South African urban adults.** *QJM.* 1998; 91(11): 743–747. [PubMed Abstract](#) | [Publisher Full Text](#)
- Thwaites GE, Lan NT, Dung NH, *et al.*: **Effect of antituberculosis drug resistance on response to treatment and outcome in adults with tuberculous meningitis.** *J Infect Dis.* 2005; 192(1): 79–88. [PubMed Abstract](#) | [Publisher Full Text](#)
- Katrak SM, Shembalkar PK, Bijwe SR, *et al.*: **The clinical, radiological and pathological profile of tuberculous meningitis in patients with and without human immunodeficiency virus infection.** *J Neurol Sci.* 2000; 181(1–2): 118–126. [PubMed Abstract](#) | [Publisher Full Text](#)
- Karande S, Gupta V, Kulkarni M, *et al.*: **Tuberculous meningitis and HIV.** *Indian J Pediatr.* 2005; 72(9): 755–760. [PubMed Abstract](#) | [Publisher Full Text](#)
- Trunz BB, Fine P, Dye C: **Effect of BCG vaccination on childhood tuberculous meningitis and miliary tuberculosis worldwide: a meta-analysis and assessment of cost-effectiveness.** *Lancet.* 2006; 367(9517): 1173–1180. [PubMed Abstract](#) | [Publisher Full Text](#)
- Mangtani P, Abubakar I, Ariti C, *et al.*: **Protection by BCG vaccine against tuberculosis: a systematic review of randomized controlled trials.** *Clin Infect Dis.* 2014; 58(4): 470–480. [PubMed Abstract](#) | [Publisher Full Text](#)
- Kelekçi S, Karabel M, Karabel D, *et al.*: **Bacillus Calmette-Guérin is a preventive factor in mortality of childhood tuberculous meningitis.** *Int J Infect Dis.* 2014; 21: 1–4. [PubMed Abstract](#) | [Publisher Full Text](#)
- Fine PE: **Variation in protection by BCG: implications of and for heterologous immunity.** *Lancet.* 1995; 346(8986): 1339–1345. [PubMed Abstract](#) | [Publisher Full Text](#)
- Arregui S, Sanz J, Marinova D, *et al.*: **On the impact of masking and blocking hypotheses for measuring the efficacy of new tuberculosis vaccines.** *PeerJ.* 2016; 4: e1513. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

24. Roy A, Eisenhut M, Harris RJ, *et al.*: **Effect of BCG vaccination against *Mycobacterium tuberculosis* infection in children: systematic review and meta-analysis.** *BMJ.* 2014; **349**: g4643.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
25. Babu S, Nutman TB: **Helminth-Tuberculosis Co-infection: An Immunologic Perspective.** *Trends Immunol.* 2016; **37**(9): 597–607.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
26. Gaze S, Bethony JM, Periago MV: **Immunology of experimental and natural human hookworm infection.** *Parasite Immunol.* 2014; **36**(8): 358–366.
[PubMed Abstract](#) | [Publisher Full Text](#)
27. King CL, Mahanty S, Kumaraswami V, *et al.*: **Cytokine control of parasite-specific energy in human lymphatic filariasis. Preferential induction of a regulatory T helper type 2 lymphocyte subset.** *J Clin Invest.* 1993; **92**(4): 1667–1673.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
28. Neto LM, Oliveira Rde V, Totino PR, *et al.*: **Enteroparasitosis prevalence and parasitism influence in clinical outcomes of tuberculosis patients with or without HIV co-infection in a reference hospital in Rio de Janeiro (2000-2006).** *Braz J Infect Dis.* 2009; **13**(6): 427–432.
[PubMed Abstract](#) | [Publisher Full Text](#)
29. Tristão-Sá R, Ribeiro-Rodrigues R, Johnson LT, *et al.*: **Intestinal nematodes and pulmonary tuberculosis.** *Rev Soc Bras Med Trop.* 2002; **35**(5): 533–535.
[PubMed Abstract](#) | [Publisher Full Text](#)
30. Elias D, Mengistu G, Akuffo H, *et al.*: **Are intestinal helminths risk factors for developing active tuberculosis?** *Trop Med Int Health.* 2006; **11**(4): 551–558.
[PubMed Abstract](#) | [Publisher Full Text](#)
31. Brown M, Miro G, Nkurunziza P, *et al.*: **Schistosoma mansoni, nematode infections, and progression to active tuberculosis among HIV-1-infected Ugandans.** *Am J Trop Med Hyg.* 2006; **74**(5): 819–825.
[PubMed Abstract](#)
32. Bar-Shavit Z, Teitelbaum SL, Reitsma P, *et al.*: **Induction of monocytic differentiation and bone resorption by 1,25-dihydroxyvitamin D3.** *Proc Natl Acad Sci U S A.* 1983; **80**(19): 5907–5911.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
33. Chandra G, Selvaraj P, Jawahar MS, *et al.*: **Effect of vitamin D₃ on phagocytic potential of macrophages with live *Mycobacterium tuberculosis* and lymphoproliferative response in pulmonary tuberculosis.** *J Clin Immunol.* 2004; **24**(3): 249–257.
[PubMed Abstract](#) | [Publisher Full Text](#)
34. Rigby WF, Shen L, Ball ED, *et al.*: **Differentiation of a human monocytic cell line by 1,25-dihydroxyvitamin D3 (calcitriol): a morphologic, phenotypic, and functional analysis.** *Blood.* 1984; **64**(5): 1110–1115.
[PubMed Abstract](#)
35. Rizvi I, Garg RK, Jain A, *et al.*: **Vitamin D status, vitamin D receptor and toll like receptor-2 polymorphisms in tuberculous meningitis: a case-control study.** *Infection.* 2016; **44**(5): 633–640.
[PubMed Abstract](#) | [Publisher Full Text](#)
36. Faksri K, Drobniewski F, Nikolayevskyy V, *et al.*: **Epidemiological trends and clinical comparisons of *Mycobacterium tuberculosis* lineages in Thai TB meningitis.** *Tuberculosis (Edinb).* 2011; **91**(6): 594–600.
[PubMed Abstract](#) | [Publisher Full Text](#)
37. Hang NT, Maeda S, Lien LT, *et al.*: **Primary drug-resistant tuberculosis in Hanoi, Viet Nam: present status and risk factors.** *PLoS One.* 2013; **8**(8): e71867.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
38. Huyen MN, Buu TN, Timersma E, *et al.*: **Tuberculosis relapse in Vietnam is significantly associated with *Mycobacterium tuberculosis* Beijing genotype infections.** *J Infect Dis.* 2013; **207**(10): 1516–1524.
[PubMed Abstract](#) | [Publisher Full Text](#)
39. Hang NT, Maeda S, Keicho N, *et al.*: **Sublineages of *Mycobacterium tuberculosis* Beijing genotype strains and unfavorable outcomes of anti-tuberculosis treatment.** *Tuberculosis (Edinb).* 2015; **95**(3): 336–342.
[PubMed Abstract](#) | [Publisher Full Text](#)
40. Gupta RK, Lucas SB, Fielding KL, *et al.*: **Prevalence of tuberculosis in post-mortem studies of HIV-infected adults and children in resource-limited settings: a systematic review and meta-analysis.** *AIDS.* 2015; **29**(15): 1987–2002.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
41. du Preez K, Schaaf HS, Dunbar R, *et al.*: **Incomplete registration and reporting of culture-confirmed childhood tuberculosis diagnosed in hospital.** *Public Health Action.* 2011; **1**(1): 19–24.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
42. Ducomble T, Tolksdorf K, KaraGiannis I, *et al.*: **The burden of extrapulmonary and meningitis tuberculosis: an investigation of national surveillance data, Germany, 2002 to 2009.** *Euro Surveill.* 2013; **18**(12): pii: 20436.
[PubMed Abstract](#)
43. Seddon JA, Hesseling AC, Marais BJ, *et al.*: **The evolving epidemic of drug-resistant tuberculosis among children in Cape Town, South Africa.** *Int J Tuberc Lung Dis.* 2012; **16**(7): 928–933.
[PubMed Abstract](#) | [Publisher Full Text](#)
44. Lincoln EM: **Tuberculous meningitis in children; with special reference to serous meningitis; serous tuberculous meningitis.** *Am Rev Tuberc.* 1947; **56**(2): 75–94.
[PubMed Abstract](#)
45. Chiang SS, Khan FA, Milstein MB, *et al.*: **Treatment outcomes of childhood tuberculous meningitis: a systematic review and meta-analysis.** *Lancet Infect Dis.* 2014; **14**(10): 947–957.
[PubMed Abstract](#) | [Publisher Full Text](#)
46. Cecchini D, Ambrosioni J, Brezzo C, *et al.*: **Tuberculous meningitis in HIV-infected and non-infected patients: comparison of cerebrospinal fluid findings.** *Int J Tuberc Lung Dis.* 2009; **13**(2): 269–271.
[PubMed Abstract](#)
47. Wait JW, Stanton L, Schoeman JF: **Tuberculosis meningitis and attention deficit hyperactivity disorder in children.** *J Trop Pediatr.* 2002; **48**(5): 294–299.
[PubMed Abstract](#) | [Publisher Full Text](#)
48. Wait JW, Schoeman JF: **Behaviour profiles after tuberculous meningitis.** *J Trop Pediatr.* 2010; **56**(3): 166–171.
[PubMed Abstract](#) | [Publisher Full Text](#)
49. Bahr NC, Boulware DR: **Methods of rapid diagnosis for the etiology of meningitis in adults.** *Biomark Med.* 2014; **8**(9): 1085–1103.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
50. Bahr NC, Meintjes G, Boulware DR: **Inadequate diagnostics: the case to move beyond the bacilli for detection of meningitis due to *Mycobacterium tuberculosis*.** *J Med Microbiol.* 2019; **68**(5): 755–760.
[PubMed Abstract](#) | [Publisher Full Text](#)
51. Bahr NC, Tugume L, Rajasingham R, *et al.*: **Improved diagnostic sensitivity for tuberculous meningitis with Xpert® MTB/RIF of centrifuged CSF.** *Int J Tuberc Lung Dis.* 2015; **19**(10): 1209–1215.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
52. Bahr NC, Nuwagira E, Evans EE, *et al.*: **Diagnostic accuracy of Xpert MTB/RIF Ultra for tuberculous meningitis in HIV-infected adults: a prospective cohort study.** *Lancet Infect Dis.* 2018; **18**(1): 68–75.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
53. Nhu NT, Heemskerck D, Thu do DA, *et al.*: **Evaluation of GeneXpert MTB/RIF for diagnosis of tuberculous meningitis.** *J Clin Microbiol.* 2014; **52**(1): 226–233.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
54. Pormohammad A, Nasiri MJ, McHugh TD, *et al.*: **A Systematic Review and Meta-analysis of the Diagnostic Accuracy of Nucleic Acid Amplification Tests for Tuberculous Meningitis.** *J Clin Microbiol.* 2019; **57**(6): pii: e01113-18.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
55. Denkinger CM, Schumacher SG, Boehme CC, *et al.*: **Xpert MTB/RIF assay for the diagnosis of extrapulmonary tuberculosis: a systematic review and meta-analysis.** *Eur Respir J.* 2014; **44**(2): 435–446.
[PubMed Abstract](#) | [Publisher Full Text](#)
56. **Automated Real-Time Nucleic Acid Amplification Technology for Rapid and Simultaneous Detection of Tuberculosis and Rifampicin Resistance: Xpert MTB/RIF Assay for the Diagnosis of Pulmonary and Extrapulmonary TB in Adults and Children: Policy Update.** Geneva, 2013.
[PubMed Abstract](#)
57. World Health Organization, Geneva, Switzerland: **Global Tuberculosis Report 2014.**
[Reference Source](#)
58. Patel VB, Theron G, Lenders L, *et al.*: **Diagnostic accuracy of quantitative PCR (Xpert MTB/RIF) for tuberculous meningitis in a high burden setting: a prospective study.** *PLoS Med.* 2013; **10**(10): e1001536.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
59. Bahr NC, Marais S, Caws M, *et al.*: **GeneXpert MTB/Rif to Diagnose Tuberculous Meningitis: Perhaps the First Test but not the Last.** *Clin Infect Dis.* 2016; **62**(9): 1133–5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
60. Boyles TH, Thwaites GE: **Appropriate use of the Xpert® MTB/RIF assay in suspected tuberculous meningitis.** *Int J Tuberc Lung Dis.* 2015; **19**(3): 276–277.
[PubMed Abstract](#) | [Publisher Full Text](#)
61. Cazabon D, Pande T, Kik S, *et al.*: **Market penetration of Xpert MTB/RIF in high tuberculosis burden countries: A trend analysis from 2014 - 2016 [version 2; peer review: 4 approved].** *Gates Open Res.* 2018; **2**: 35.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
62. World Health Organization, Geneva, Switzerland: **WHO Meeting Report of a Technical Expert Consultation: Non-inferiority analysis of Xpert MTB/RIF Ultra compared to Xpert MTB/RIF.** (accessed 11 April 2019). 2017.
[Reference Source](#)
63. Marais S, Thwaites G, Schoeman JF, *et al.*: **Tuberculous meningitis: a uniform case definition for use in clinical research.** *Lancet Infect Dis.* 2010; **10**(11): 803–812.
[PubMed Abstract](#) | [Publisher Full Text](#)
64. Wang G, Wang S, Jiang G, *et al.*: **Xpert MTB/RIF Ultra improved the diagnosis of paucibacillary tuberculosis: A prospective cohort study.** *J Infect.* 2019; **78**(4): 311–316.
[PubMed Abstract](#) | [Publisher Full Text](#)
65. Bahr NC, Halupnick R, Linder G, *et al.*: **Delta-like 1 protein, vitamin D binding protein and fetuin for detection of *Mycobacterium tuberculosis* meningitis.** *Biomark Med.* 2018; **12**(7): 707–716.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
66. Heemskerck D, Day J, Chau TT, *et al.*: **Intensified treatment with high dose rifampicin and levofloxacin compared to standard treatment for adult patients with tuberculous meningitis (TBM-IT): protocol for a randomized controlled trial.** *Trials.* 2011; **12**: 25.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
67. Cooke RE, Dunphy DL, Blake FG: **Streptomycin in tuberculous meningitis; a**

- report of its use in a one-year-old infant. *Yale J Biol Med.* 1946; **18**: 221–226.
[PubMed Abstract](#) | [Free Full Text](#)
68. Tucker WB: **Evaluation of streptomycin regimens in the treatment of tuberculosis; an account of the study of the Veterans Administration, Army, and Navy, July 1946 to April 1949.** *Am Rev Tuberc.* 1949; **60**(6): 715–754.
[PubMed Abstract](#)
69. Smith HV, Vollum RL, Cairns H: **Treatment of tuberculous meningitis with streptomycin.** *Lancet.* 1948; **1**(6504): 627–636.
[PubMed Abstract](#) | [Publisher Full Text](#)
70. Cairns H, Smith HV, Vollum RL: **Tuberculous meningitis.** *J Am Med Assoc.* 1950; **144**(2): 92–96.
[PubMed Abstract](#) | [Publisher Full Text](#)
71. McSweeney CJ: **The treatment of tuberculous meningitis.** *Tubercle.* 1950; **31**(9): 210–213.
[PubMed Abstract](#) | [Publisher Full Text](#)
72. Bernard E, Drouot A, Lotte A, *et al.*: **[Results of therapy of tuberculous meningitis in adults; study of 255 cases].** *Bull Mem Soc Med Hop Paris.* 1954; **70**(30–31): 1118–1127.
[PubMed Abstract](#)
73. Lorber J: **The results of treatment of 549 cases of tuberculous meningitis.** *Am Rev Tuberc.* 1954; **69**(1): 13–25.
[PubMed Abstract](#)
74. Debre R: **The prognosis of tuberculous meningitis.** *Am Rev Tuberc.* 1952; **65**(2): 168–180.
[PubMed Abstract](#)
75. Lepper MH, Spies HW: **The present status of the treatment of tuberculosis of the central nervous system.** *Ann N Y Acad Sci.* 1963; **106**: 106–123.
[PubMed Abstract](#) | [Publisher Full Text](#)
76. Donald PR: **The chemotherapy of tuberculous meningitis in children and adults.** *Tuberculosis (Edinb).* 2010; **90**(6): 375–392.
[PubMed Abstract](#) | [Publisher Full Text](#)
77. Fitzpatrick MJ: **The treatment of tuberculous meningitis; a study of forty-six adult patients.** *Am Rev Tuberc.* 1954; **69**(3): 370–382.
[PubMed Abstract](#)
78. Lorber J: **Tuberculous meningitis in children treated with streptomycin and P.A.S.** *Lancet.* 1954; **266**(6822): 1104–1107.
[PubMed Abstract](#) | [Publisher Full Text](#)
79. Debre R, Brissaud HE: **Present method and results of treatment of tuberculous meningitis.** *Am Rev Tuberc.* 1956; **74**(2 Part 2): 221–224.
[PubMed Abstract](#)
80. Elmendorf DF Jr, Cawthon WU, Muschenheim C, *et al.*: **The absorption, distribution, excretion, and short-term toxicity of isonicotinic acid hydrazide (hydrazid) in man.** *Am Rev Tuberc.* 1952; **65**(4): 429–442.
[PubMed Abstract](#)
81. Barclay WR, Ebert RH, Le Roy GV, *et al.*: **Distribution and excretion of radioactive isoniazid in tuberculous patients.** *J Am Med Assoc.* 1953; **151**(16): 1384–1388.
[PubMed Abstract](#)
82. Fletcher AP: **C.S.F.–isoniazid levels in tuberculous meningitis.** *Lancet.* 1953; **265**(6788): 694–696.
[PubMed Abstract](#) | [Publisher Full Text](#)
83. Pellegrino ED, Petrik FG, Horton R: **The treatment of tuberculous meningitis in infants with streptomycin and isonicotinic acid hydrazide (isoniazid); a preliminary report of six patients under the age of two years treated without intrathecal medication.** *Dis Chest.* 1954; **26**(2): 146–165.
[PubMed Abstract](#) | [Publisher Full Text](#)
84. Rahajoe NN, Rahajoe N, Boediman I, *et al.*: **The treatment of tuberculous meningitis in children with A combination of isoniazid, rifampicin and streptomycin (a preliminary report).** *Paediatr Indones.* 1979; **19**(11–12): 285–294.
[PubMed Abstract](#) | [Publisher Full Text](#)
85. Doğanay M, Bakır M, Dökmetaş I: **Treatment of tuberculous meningitis in adults with a combination of isoniazid, rifampicin and streptomycin: a prospective study.** *Scand J Infect Dis.* 1989; **21**(1): 81–85.
[PubMed Abstract](#) | [Publisher Full Text](#)
86. Girgis NI, Yassin MW, Laughlin LW, *et al.*: **Rifampicin in the treatment of tuberculous meningitis.** *J Trop Med Hyg.* 1978; **81**(12): 246–247.
[PubMed Abstract](#)
87. Sunakorn P, Wongsarop J: **Rifampin in treatment of tuberculous meningitis in children.** *J Med Assoc Thai.* 1978; **61**(2): 93–98.
[PubMed Abstract](#)
88. Donald PR: **Cerebrospinal fluid concentrations of antituberculosis agents in adults and children.** *Tuberculosis (Edinb).* 2010; **90**(5): 279–292.
[PubMed Abstract](#) | [Publisher Full Text](#)
89. Savic RM, Ruslami R, Hibma JE, *et al.*: **Pediatric tuberculous meningitis: Model-based approach to determining optimal doses of the anti-tuberculosis drugs rifampin and levofloxacin for children.** *Clin Pharmacol Ther.* 2015; **98**(6): 622–629.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
90. Ruslami R, Ganiem AR, Dian S, *et al.*: **Intensified regimen containing rifampicin and moxifloxacin for tuberculous meningitis: an open-label, randomised controlled phase 2 trial.** *Lancet Infect Dis.* 2013; **13**(1): 27–35.
[PubMed Abstract](#) | [Publisher Full Text](#)
91. Davis A, Meintjes G, Wilkinson RJ: **Treatment of Tuberculous Meningitis and Its Complications in Adults.** *Curr Treat Options Neurol.* 2018; **20**(3): 5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
92. Donald PR: **Chemotherapy for Tuberculous Meningitis.** *N Engl J Med.* 2016; **374**(2): 179–181.
[PubMed Abstract](#) | [Publisher Full Text](#)
93. In: th ed. **Treatment of Tuberculosis: Guidelines.** Geneva. 2010.
94. In: nd ed. **Guidance for National Tuberculosis Programmes on the Management of Tuberculosis in Children.** Geneva. 2014.
95. In: **Rapid Advice: Treatment of Tuberculosis in Children.** Geneva. 2010.
96. Thwaites GE, Bhavnani SM, Chau TT, *et al.*: **Randomized pharmacokinetic and pharmacodynamic comparison of fluoroquinolones for tuberculous meningitis.** *Antimicrob Agents Chemother.* 2011; **55**(7): 3244–3253.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
97. Heemskerck AD, Bang ND, Mai NT, *et al.*: **Intensified Antituberculosis Therapy in Adults with Tuberculous Meningitis.** *N Engl J Med.* 2016; **374**(2): 124–134.
[PubMed Abstract](#) | [Publisher Full Text](#)
98. Rizvi I, Malhotra HS, Garg RK, *et al.*: **Fluoroquinolones in the management of tuberculous meningitis: Systematic review and meta-analysis.** *J Infect.* 2018; **77**(4): 261–275.
[PubMed Abstract](#) | [Publisher Full Text](#)
99. Ruslami R, Ganiem AR, Aarnoutse RE, *et al.*: **Rifampicin and moxifloxacin for tuberculous meningitis—authors' reply.** *Lancet Infect Dis.* 2013; **13**(7): 570.
[PubMed Abstract](#) | [Publisher Full Text](#)
100. Te Brake L, Dian S, Ganiem AR, *et al.*: **Pharmacokinetic/pharmacodynamic analysis of an intensified regimen containing rifampicin and moxifloxacin for tuberculous meningitis.** *Int J Antimicrob Agents.* 2015; **45**(5): 496–503.
[PubMed Abstract](#) | [Publisher Full Text](#)
101. van Toorn R, Schaaf HS, Laubscher JA, *et al.*: **Short intensified treatment in children with drug-susceptible tuberculous meningitis.** *Pediatr Infect Dis J.* 2014; **33**(3): 248–252.
[PubMed Abstract](#) | [Publisher Full Text](#)
102. Prasad K, Singh MB, Ryan H: **Corticosteroids for managing tuberculous meningitis.** *Cochrane Database Syst Rev.* 2016; **4**: CD002244.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
103. Misra UK, Kalita J, Nair PP: **Role of aspirin in tuberculous meningitis: a randomized open label placebo controlled trial.** *J Neurol Sci.* 2010; **293**(1–2): 12–17.
[PubMed Abstract](#) | [Publisher Full Text](#)
104. Misra UK, Kalita J, Sagar B, *et al.*: **Does adjunctive corticosteroid and aspirin therapy improve the outcome of tuberculous meningitis?** *Neurol India.* 2018; **66**(6): 1672–1677.
[PubMed Abstract](#) | [Publisher Full Text](#)
105. Mai NT, Dobbs N, Phu NH, *et al.*: **A randomised double blind placebo controlled phase 2 trial of adjunctive aspirin for tuberculous meningitis in HIV-uninfected adults.** *eLife.* 2018; **7**: pii: e33478.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
106. Schoeman JF, Janse van Rensburg A, Laubscher JA, *et al.*: **The role of aspirin in childhood tuberculous meningitis.** *J Child Neurol.* 2011; **26**(8): 956–962.
[PubMed Abstract](#) | [Publisher Full Text](#)
107. van Toorn R, du Plessis AM, Schaaf HS, *et al.*: **Clinicoradiologic response of neurologic tuberculous mass lesions in children treated with thalidomide.** *Pediatr Infect Dis J.* 2015; **34**(2): 214–218.
[PubMed Abstract](#) | [Publisher Full Text](#)
108. Schoeman JF, Andronikou S, Stefan DC, *et al.*: **Tuberculous meningitis-related optic neuritis: recovery of vision with thalidomide in 4 consecutive cases.** *J Child Neurol.* 2010; **25**(7): 822–828.
[PubMed Abstract](#) | [Publisher Full Text](#)
109. Wilkinson RJ, Rohlwin U, Misra UK, *et al.*: **Tuberculous meningitis.** *Nat Rev Neurol.* 2017; **13**(10): 581–598.
[PubMed Abstract](#) | [Publisher Full Text](#)
110. Molton JS, Huggan PJ, Archuleta S: **Infliximab therapy in two cases of severe neurotuberculosis paradoxical reaction.** *Med J Aust.* 2015; **202**(3): 156–157.
[PubMed Abstract](#) | [Publisher Full Text](#)
111. Lee JY, Yim JJ, Yoon BW: **Adjuvant interferon- γ treatment in two cases of refractory tuberculosis of the brain.** *Clin Neurol Neurosurg.* 2012; **114**(6): 732–734.
[PubMed Abstract](#) | [Publisher Full Text](#)
112. Bruwer GE, Van der Westhuizen S, Lombard CJ, *et al.*: **Can CT predict the level of CSF block in tuberculous hydrocephalus?** *Childs Nerv Syst.* 2004; **20**(3): 183–187.
[PubMed Abstract](#) | [Publisher Full Text](#)
113. van Toorn R, Solomons R: **Update on the diagnosis and management of tuberculous meningitis in children.** *Semin Pediatr Neurol.* 2014; **21**(1): 12–18.
[PubMed Abstract](#) | [Publisher Full Text](#)
114. Cazabon D, Alsdurf H, Satyanarayana S, *et al.*: **Quality of tuberculosis care in high burden countries: the urgent need to address gaps in the care cascade.** *Int J Infect Dis.* 2017; **56**: 111–116.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
115. Kim J, Keshavjee S, Atun R: **Health systems performance in managing tuberculosis: analysis of tuberculosis care cascades among high-burden and non-high-burden countries.** *J Glob Health.* 2019; **9**(1): 010423.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
116. Naidoo P, Theron G, Rangaka MX, *et al.*: **The South African Tuberculosis Care**

- Cascade: Estimated Losses and Methodological Challenges.** *J Infect Dis.* 2017; **216**(suppl_7): S702–S713.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
117. Parija D, Patra TK, Kumar AM, *et al.*: **Impact of awareness drives and community-based active tuberculosis case finding in Odisha, India.** *Int J Tuberc Lung Dis.* 2014; **18**(9): 1105–1107.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
118. He Y, Han C, Chang KF, *et al.*: **Total delay in treatment among tuberculous meningitis patients in China: a retrospective cohort study.** *BMC Infect Dis.* 2017; **17**(1): 341.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
119. Solomons R, Grantham M, Marais BJ, *et al.*: **IMCI indicators of childhood TBM at primary health care level in the Western Cape Province of South Africa.** *Int J Tuberc Lung Dis.* 2016; **20**(10): 1309–1313.
[PubMed Abstract](#) | [Publisher Full Text](#)
120. Sheu JJ, Yuan RY, Yang CC: **Predictors for outcome and treatment delay in patients with tuberculous meningitis.** *Am J Med Sci.* 2009; **338**(2): 134–139.
[PubMed Abstract](#) | [Publisher Full Text](#)

Open Peer Review

Current Peer Review Status:  

Version 1

Reviewer Report 17 February 2020

<https://doi.org/10.21956/wellcomeopenres.17007.r37823>

© 2020 Kadhiraivan T. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Tamilarasu Kadhiraivan 

Department of Medicine, Jawaharlal Institute of Postgraduate Medical Education and Research, Puducherry, Puducherry, India

In this article, Seddon *et al.* have provided a comprehensive overview of the pathogenesis, epidemiology, and diagnostics of tuberculous meningitis. Further, this article traces the historical progress made in the treatment of tuberculous meningitis, and also provides a succinct summary of the recently concluded as well as ongoing clinical trials of intensified/adjunctive treatments for tuberculous meningitis. Overall, this is a well written article, and it provides a contextual background for the other articles in the collection.

I have two suggestions. First, one of the important barriers to early diagnosis of tuberculous meningitis could be the (incorrect) notion that tuberculous meningitis is a subacute disease process. Contrary to this notion, nearly half the patients with tuberculous meningitis seen in many settings have less than 2 weeks of symptoms at presentation. However, clinicians still fail to consider tuberculous meningitis in such patients with less than 2 weeks of symptoms.

Second, I am not sure how many of the national TB control programmes worldwide provide dexamethasone free-of-cost to tuberculous meningitis patients. My suspicion is that current health systems are unable to reliably deliver the only effective adjunctive treatment available to treat tuberculous meningitis.

Is the topic of the review discussed comprehensively in the context of the current literature?

Yes

Are all factual statements correct and adequately supported by citations?

Yes

Is the review written in accessible language?

Yes

Are the conclusions drawn appropriate in the context of the current research literature?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Clinical research; clinical epidemiology; randomised controlled trials; systematic reviews

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Reviewer Report 09 December 2019

<https://doi.org/10.21956/wellcomeopenres.17007.r36945>

© 2019 Thakur K. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Kiran T. Thakur

Department of Neurology, Columbia University Irving Medical Center, New York City, NY, USA

Experts in TB and TB meningitis in the Tuberculous Meningitis International Research Consortium provide an overview on the global situation for tuberculous meningitis and discuss current knowledge around epidemiology, diagnostic testing and treatment for TB meningitis. The authors provide a well-written article around the basic aspects of TBM which provides an important overview on the topic, including knowledge gaps in our fundamental understanding of TBM pathogenesis, diagnosis and treatments. Though much of the review of data has been previously published, authors provide important novel aspects including implications of our lack of knowledge in key aspects of TBM, and aspects of healthcare system infrastructure in current TBM care.

The authors discuss the debate in TBM pathogenesis including historical work by Roch and McCordock. Additional commentary on miliary TB pathogenesis would be useful to the readership.

The natural history of TBM is discussed around host factors. The discussion focuses on risk factors including age, HIV, BCG vaccine, helminth infections, nutrition and strain type though importantly does not mention other causes of immunocompromised state including immunomodulatory treatments and possible environmental factors which are of growing importance globally (overcrowding in inner city environments, pollution, etc). A brief discussion on the immunological aspects which may children particularly susceptible to disseminated disease would also be educational to the reader. The discussion of strain type as a risk factor is important, though only touched upon very briefly in this article and worthy of expansion.

The unknowns in terms of current epidemiology of TBM is also focused upon. It would be

worthwhile for authors to summarize past epidemiological data regionally, and also note ongoing studies on epidemiology of TBM. An important milestone would be a multicenter international study on epidemiology of TBM, and the consortium may be planning such work.

Epidemiological studies are limited by current diagnostics in TBM, and the authors importantly note the limited availability of gene xpert and gene xpert ultra. It would be important to identify where this testing is actually available and the barriers to implementation of such testing at the healthcare system level in those where the testing does not exist.

The treatment discussion is nicely delineated, with a historical and current perspective discussion. Missing from the discussion is commentary on treatment in the pediatric population and treatment initiation of antiretroviral medications in patients with TBM, both challenges to those at the bedside treating such patients. It would be worthwhile to expand the treatment section to include supportive care given the importance of neurointensive care efforts including ICP monitoring, treatment of hyponatremia, when to consider shunting etc.

The healthcare systems section of the paper is important and novel and would benefit perhaps from a visual of the current system at the primary, secondary and tertiary care health systems levels and major knowledge gaps/policy gaps in clinical practice.

Expansion of Table one is also important in terms of future directions beyond implications of challenges in TBM--how do we study and fill this knowledge gap is essential in TBM.

Overall, while the article does not provide new data on TBM, the review provides an important summary on our current state of knowledge, with an emphasis on knowledge gaps. I applaud the authors for their work on this important disease, and look forward to further fruitful work from the consortium.

Is the topic of the review discussed comprehensively in the context of the current literature?

Yes

Are all factual statements correct and adequately supported by citations?

Yes

Is the review written in accessible language?

Yes

Are the conclusions drawn appropriate in the context of the current research literature?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Clinical research in neuroinfectious diseases, Studies on access to neurological care in resource-limited settings

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.