

CORRECTION

Open Access



# Correction to: Impact of district mental health care plans on symptom severity and functioning of patients with priority mental health conditions: the Programme for Improving Mental Health Care (PRIME) cohort protocol

Emily C. Baron<sup>1\*</sup>, Sujit D. Rathod<sup>2</sup>, Charlotte Hanlon<sup>3,4</sup>, Martin Prince<sup>4</sup>, Abebaw Fedaku<sup>5,6</sup>, Fred Kigozi<sup>7</sup>, Mark Jordans<sup>8,9</sup>, Nagendra P. Luitel<sup>10</sup>, Girmay Medhin<sup>11</sup>, Vaibhav Murhar<sup>12</sup>, Juliet Nakku<sup>7</sup>, Vikram Patel<sup>12,13,14</sup>, Inge Petersen<sup>15</sup>, One Selohilwe<sup>15</sup>, Rahul Shidhaye<sup>16,17</sup>, Joshua Ssebunnya<sup>7</sup>, Mark Tomlinson<sup>1,18</sup>, Crick Lund<sup>1,9</sup> and Mary De Silva<sup>19</sup>

**Correction to: BMC Psychiatry 18, 61 (2018)**  
<https://doi.org/10.1186/s12888-018-1642-x>

Following publication of the original article [1], the authors identified an error in Table 2. The correct table is given below.

## Author details

<sup>1</sup>Alan J Flisher Centre for Public Mental Health, Department of Psychiatry and Mental Health, University of Cape Town, 46 Sawkins Road 7700 Rondebosch, Cape Town, South Africa. <sup>2</sup>Department of Population Health, London School of Hygiene and Tropical Medicine, London, UK. <sup>3</sup>College of Health Sciences, School of Medicine, Department of Psychiatry, Addis Ababa University, Addis Ababa, Ethiopia. <sup>4</sup>Centre for Global Mental Health, Health Services and Population Research Department, King's College London, London, UK. <sup>5</sup>Centre for Innovative Drug Development and Therapeutic Trials for Africa, College of Health Sciences, Addis Ababa University, Addis Ababa, Ethiopia. <sup>6</sup>Global Health and Infection Department, Brighton and Sussex Medical School, University of Sussex, Brighton, UK. <sup>7</sup>Butabika National Referral and Teaching Mental Hospital, Makerere University, Kampala, Uganda. <sup>8</sup>Research and Development Department, HealthNet TPO, Amsterdam, the Netherlands. <sup>9</sup>Center for Global Mental Health, Institute of Psychiatry, Psychology and

Neuroscience, King's College London, London, UK. <sup>10</sup>Research Department, Transcultural Psychosocial Organization (TPO) Nepal, Baluwatar, Kathmandu, Nepal. <sup>11</sup>Aklilu Lemma Institute of Pathobiology, Addis Ababa University, Addis Ababa, Ethiopia. <sup>12</sup>Sangath, Goa, India. <sup>13</sup>Harvard Medical School, Boston, USA. <sup>14</sup>Public Health Foundation of India, New Delhi, India. <sup>15</sup>Centre for Rural Health, School of Nursing and Public Health and School of Applied Human Sciences, University of KwaZulu-Natal, KwaZulu-Natal, South Africa. <sup>16</sup>Centre for Chronic Conditions and Injuries, Public Health Foundation of India, New Delhi, India. <sup>17</sup>CAPHRI School for Public Health and Primary Care, Maastricht University, Maastricht, the Netherlands. <sup>18</sup>Alan J Flisher Centre for Public Mental Health, Department of Psychology, Stellenbosch University, Stellenbosch, South Africa. <sup>19</sup>Department of Population Health, Wellcome Trust, London, UK.

Published online: 29 September 2020

## Reference

1. Baron, et al. Impact of district mental health care plans on symptom severity and functioning of patients with priority mental health conditions: the Programme for Improving Mental Health Care (PRIME) cohort protocol. *BMC Psychiatry*. 2018;18:61. <https://doi.org/10.1186/s12888-018-1642-x>.

The original article can be found online at <https://doi.org/10.1186/s12888-018-1642-x>.

\* Correspondence: [emily.baron@uct.ac.za](mailto:emily.baron@uct.ac.za)

<sup>1</sup>Alan J Flisher Centre for Public Mental Health, Department of Psychiatry and Mental Health, University of Cape Town, 46 Sawkins Road 7700 Rondebosch, Cape Town, South Africa

Full list of author information is available at the end of the article



© The Author(s). 2020 **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

**Table 2** Assessment schedule for PRIME Cohorts

Data collected by questionnaire	Depression			Alcohol use disorders			Psychosis			Epilepsy		
	0	3/6	12	0	3	12	0	6	12	0	6	12
DEMOGRAPHICS CHARACTERISTICS	✓			✓			✓			✓		
CLINICAL MEASURES												
WHO Disability Assessment Schedule (WHODAS 2.0) [34]	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Patient Health Questionnaire (PHQ-9) [26]	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Alcohol Use Disorder Identification Test (AUDIT) [27]	✓			✓	✓	✓	✓	✓	✓	✓		
Short Inventory of Problems – Recent (SIP 2-R) [75, 76]				✓	✓	✓						
Suicidality (Composite International Diagnostic Interview - suicidality module) [77]	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Epilepsy severity (developed by PRIME)											✓	✓
Brief Psychiatric Rating Scale (BPRS-E) [30]							✓	✓	✓			
Positive and Negative Syndrome Scale (PANSS) [78]							✓	✓	✓			
HEALTH SERVICE USE												
Group/community interventions (developed by PRIME)	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Mental health services received (developed by PRIME)		✓	✓		✓	✓		✓	✓		✓	✓
Health Service use and costs (adapted from the Client Service Receipt Inventory)	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
MEDICATION ADHERENCE												
Morisky, Green and Levine Adherence Scale [79]	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Medication adherence (adapted from Care for People with Schizophrenia in India)	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
SOCIAL AND ECONOMIC MEASURES												
Economic activity (adapted from WHODAS 2.0, added items by PRIME)	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Severe Adverse Events (developed by PRIME)		✓	✓		✓	✓		✓	✓		✓	✓
Oslo 3-item Social Support Scale [80]	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Caregiver work burden - WHO Family Interview Schedule (Impact) [81]							✓	✓	✓	✓	✓	✓
Caregiver economic activity (adapted from WHODAS 2.0, items added by PRIME)							✓	✓	✓			
STIGMA AND DISCRIMINATION												
Discrimination and Stigma Scale [82]							✓		✓	✓		✓
Caregiver stigma & discrimination - WHO Family Interview Schedule (Stigma) [81]										✓		✓
Human rights abuse by caregiver (developed by PRIME)							✓	✓	✓	✓	✓	✓

\* 6 months for depression in Ethiopia