Invasion, alien control and restoration: Legacy effects linked to folivorous insects and phytopathogenic fungi

MALEBAJOA A. MAOELA,^{1*} SHAYNE M. JACOBS,¹ FRANCOIS ROETS^{1,2} AND KAREN J. ESLER^{1,3}

¹*Department of Conservation Ecology and Entomology, Stellenbosch University, Private Bag X1, Matieland 7602 (E-mail: malebajoam@gmail.com), ²DST/NRF Centre of Excellence in Tree Health Biotechnology (CTHB), Forestry and Agricultural Biotechnology Institute (FABI), University of Pretoria, Pretoria and ³DST/NRF Centre of Excellence for Invasion Biology, Stellenbosch University, Matieland, South Africa*

Abstract Invasive alien trees increase native tree stress and may increase attack by herbivores and pathogenic fungi. Alien tree removal should ameliorate such impacts. Here we compared the levels of damage by phytopathogenic fungi and folivorous insects on *Brabejum stellatifolium* and *Metrosideros angustifolia* (native trees) and *Acacia mearnsii* (invasive tree species) among near-pristine, invaded and restored sites. Generally, foliar damage levels were higher at invaded than at near-pristine sites. Damage levels at restored sites were similar, or even higher than those at invaded sites. Decreased native tree species richness did not explain these patterns, as restored sites had native tree species richness levels similar to those of near-pristine sites. Increased host abundance and leaf nitrogen content did not significantly correlate with increased damage in most cases. Therefore, plant species richness recovers following restoration, but native trees still experience increased pressure from folivores and phytopathogenic fungi, which may even exceed levels experienced at invaded sites, thus impacting recovery trajectories.

Key words: *Acacia mearnsii*, fynbos, leaf damage, leaf nutrient, pathogen, riparian system.

INTRODUCTION

Pest and pathogen loads in plant communities vary with plant stress levels (Gilbert 2002), native plant species richness (Scherber *et al.* 2010), host abundance (Mitchell *et al.* 2002; Bell *et al.* 2006) and plant nutritional status (Forkner & Hunter 2000; Moran & Scheidler 2002). Fragmentation of tropical forests, for example, leads to increased plant stress and, in turn, increased damage by pathogenic fungi and folivores (Benítez-Malvido *et al.* 1999). Invasive alien plants may also increase native plant stress by altering environmental conditions such as water availability, amount of sunlight and nutrient availability (Brooks *et al.* 2004). Because these factors are known to affect host–plant/pest (and pathogen) dynamics (White *et al.* 2006), we hypothesize that invasive alien plants would increase the susceptibility of native plants to natural enemies (folivorous insects and pathogenic fungi) in invaded systems.

In South Africa, woody invasive alien trees have been targeted for removal in riparian areas in a 15-year-old landscape level restoration initiative led by the extended public works programme, Working for Water. Although this initiative is credited with the creation of thousands of jobs, a body of evidence relating to the lack of effectiveness of the Working for Water initiative for curbing the spread of invasive alien trees is emerging

(van Wilgen *et al.* 2012). In addition, clearing of invasive alien plants results in gap formation, alters the ecosystem microenvironment (including light, moisture and temperature regimes) and increases patchiness and habitat heterogeneity (Torchin & Mitchell 2004). As folivore and plant pathogen populations may directly depend on these factors, their impact on host plants is also likely to change in response to these altered conditions (Folgarait *et al.* 1995).

The influence of plant species diversity on performance of natural enemies has been a focus of several experimental studies (Pfisterer *et al.* 2003; Agrawal *et al.* 2006; van Hezewijk *et al.* 2008; Scherber *et al.* 2010). Specifically, many studies found that altered native plant diversity can alter the stability, diversity and performance of their natural enemies (Mitchell *et al.* 2002; Johnson *et al.* 2008). Two hypotheses have been proposed to explain such effects (Root 1973). The ‘resource concentration’ hypothesis predicts that folivores, especially specialist folivores, increase in abundance with increasing concentration of preferred resources in more homogenous stands of host plants compared with heterogeneous stands (Grez & Gonzalez 1995). The ‘enemy’ hypothesis states that the lower folivory levels in heterogeneous communities are a result of increased effectiveness of predators and parasitoids in controlling insect pest densities (Bach 1980; Riihimäki *et al.* 2005).

Apart from plant species diversity and leaf nutrient levels (Peeters 2002a, 2002b) host abundance can also influence the prevalence of folivorous arthropods and

*Corresponding author.

Accepted for publication March 2016.

phytopathogenic fungi (Agrawal *et al.* 2006). Hypotheses relating to host abundance and folivore attack rates generally predict increased rates as host abundance increases (Agrawal *et al.* 2006; Cook & Holt 2006). Some studies, however, have found evidence that natural enemy attack is reduced on plants in dense stands compared with plants in sparse stands (Bach 1980; Garrett & Mundt 2000). Changes in host abundance might influence: (i) the quality and quantity of resources (host plant individuals) available for pathogenic fungi or folivores (Bach 1980; Cook & Holt 2006); (ii) the spatial arrangement of host plants (Folgarait *et al.* 1995), where increased host density reduces search time and increases occupation time for some folivore species (Cook & Holt 2006); and (iii) the microclimatic conditions under which the folivore/pathogen and host interact (Folgarait *et al.* 1995).

How native tree species richness, host plant abundance and leaf nutrient levels affect plant exposure to attack by pathogenic fungi and folivorous insects in a restoration scenario has received little focussed attention (Covington *et al.* 1997). Here, we investigate whether differences exist in the levels of leaf damage caused by folivorous insects and phytopathogenic fungi on an invasive alien plant (*Acacia mearnsii* De Wild (L.)) and two native plants (*Brabejum stellatifolium* (L.) and *Metrosideros angustifolia* (L.)) in riparian sites that differ in invasion status. Three categories were chosen: (i) near-pristine (few *A. mearnsii* individuals); (ii) heavily invaded (*A. mearnsii*-dominated communities); and (iii)

restored sites that were cleared of invasive *A. mearnsii* more than 7 years ago. The roles of variation in native tree species richness, host abundance and leaf nutrients in explaining observed differences in damage levels were evaluated. These data may provide tools for evaluating restoration success in the future.

METHODS

Study sites and focal species

The study was conducted in mountain stream and foothill sections of three riparian systems within south-western Cape, South Africa (Fig. 1). The area is characterized by a Mediterranean-type climate with cool rainy winters (June–August) and hot dry summers (December–February) (Deacon *et al.* 1992). The riparian ecosystems contain some fynbos vegetation (dominated by Ericaceae, Proteaceae and Restionaceae) (Mucina & Rutherford 2006) and include a variety of tree taxa that form forest pockets (Goldblatt & Manning 2000). Nine study sites in three different catchments were selected based on invasion status: three near-pristine sites (reference sites), three heavily invaded sites (dominated by *A. mearnsii*) and three restored sites (formerly invaded sites that had been cleared of invasive plants more than 7 years prior to this study) (Fig. 1). Site categorization was based on visual scoring of *A. mearnsii* cover within two transects measuring 50 m in length (parallel to the river) and 5 m in width (perpendicular to the river crossing both wet and dry bank zones). For heavily invaded sites, the *A. mearnsii* canopy cover was greater than 75%, and for near-pristine sites, the cover

Fig. 1. Location of the three Western Cape rivers: Dwars River (1), Molenaars River (2) and Wit River (3) and the nine sites identified in this study. Circles, near-pristine sites; squares, heavily invaded sites; triangles, restored sites.

was greater than 5%. For restored sites, site categorization was based on historical records and the cover was greater than 5%. Reference sites are frequently used in restoration projects to provide direction for restoration goals (Blanchard & Holmes 2008). Potential sites were located by using information from previous studies (Blanchard & Holmes 2008) and through discussions with Western Cape nature conservation managers, members of Water for Water and private landowners.

Two species native to fynbos riparian zones, *B. stellatifolium* (Proteaceae) and *M. angustifolia* (Myrtaceae) (Crous *et al.* 2012), were selected. These evergreen trees are common components of these riparian habitats and considered key species in south-western Cape Mediterranean-type riparian systems (Galatowitsch & Richardson 2005). Within the riparian ecosystems studied, invasive taxa include numerous Australian acacias such as *Acacia cyclops* (A. Cunn. ex G. Don), *Acacia longifolia* ((Andrew) Wild), *Acacia saligna* ((Labill.) H.L. Wendl.) and *Acacia melanoxylon* (R.Br.). However, an evergreen *A. mearnsii* is by far the most abundant and was subsequently selected for comparisons with the selected native species. It is also considered the most problematic species in these riparian systems with individuals also commonly found scattered in near-pristine areas (Dye & Jarman 2004).

Extent of folivorous insect and pathogen damage

At each of the nine sites, five individuals of each of the three focal tree taxa were randomly selected for comparisons of the extent of damage caused by folivorous insects and phytopathogenic fungi. Selected individuals had similar heights and stem diameters for each species respectively across all sites (to minimize possible difference in susceptibility due to plant age), and all were within 5 m of the water's edge (to minimize possible differences in susceptibility due to water stress) (Crous *et al.* 2012).

During August and October 2011, five small branches (about 5 cm in diameter and 30 cm long) were randomly collected from each individual tree at the canopy level. All leaves were detached from branches, and mature leaves were combined into a single bulk sample per individual tree in a brown paper bag. From these bulk samples, 100 leaves for each tree individual were randomly chosen (blindly drawing from brown paper bag) in the laboratory. Symptom assessments were conducted for each leaf in the laboratory by using a dissecting microscope, allowing careful confirmation of damage type.

Leaves were assigned to one of the following damage types: (i) fungal damage alone; (ii) folivore damage alone; and (iii) intact leaves (García-Guzmán & Dirzo 2004). Wounds created by folivorous insects are often entry points for pathogenic fungi (Agrios 2005). Therefore, leaves with both fungal and insect damages were not included in analyses because this would bias enumeration of leaves with fungal pathogen damage. Insect damage was categorized into two types – that caused by insects with chewing mouthparts and those with piercing and sucking mouthparts. Damage caused by leaf mining insects was negligible and not included in the analyses. Leaves colonized by phytopathogenic fungi were separated according to four

predominant disease symptoms (blight, chlorotic areas, necrotic spots and black mould). Identifications of the phytopathogenic fungi that caused these main symptoms were based on a combination of characters including the symptoms themselves, fungal micromorphology (studied with the aid of a light and dissecting microscope), culture characteristics (as grown for 2 weeks in the dark at 25 °C on malt extract agar, Biolab, South Africa) and published literature (Doidge & Bottomley 1931; Morris *et al.* 1988; Arzanlou *et al.* 2007).

One leaf of each of the five sampled individual trees per tree species per site from each the three catchments was used for DNA extraction and amplification. Therefore, for each disease type on each tree species, we sequenced nine separate fungal individuals. DNA extraction and PCR procedures followed Musvuugwa *et al.* (2015) without modification. The primers ITS1-f (Gardes & Bruns 1993) and ITS4 (White *et al.* 1990) were used for amplification and sequencing reactions. Preliminary identifications were confirmed by comparisons with those of taxa published on GenBank (<http://www.ncbi.nlm.nih.gov/genbank/>) by using standard BLAST searches. Pathogenicity tests were not performed because repeated isolations always resulted in the isolation of the same fungus species from the same lesion type, and we therefore assumed that the identified fungal species were the causal agents of the necrotic lesions.

Severity of pest and pathogen damage

Damage severity was defined as the percentage of leaf area visibly infected by a pathogen or by feeding activities of arthropods. The percentage of area lost to disease or folivory was determined by using a subjective estimate of lamina damage (1 = intact; 2 = 1–11%; 3 = 12–49% and 4 = 50–100% of leaf area damaged) based on previous studies (Benítez-Malvido *et al.* 1999; García-Guzmán & Dirzo 2004; Eichhorn *et al.* 2010). Leaf damage for all tree individuals was scored by the same person. The estimates of leaf area lost were based on comparisons with a standard leaf of the same size as that being scored. Using these methods, we could therefore not account for leaves that were completely removed from the plants, and our estimations may therefore be lower than actual damage levels.

To test the effects of invasion status on the numbers of leaves damaged (extent of damage) and the categories of damage severity, for each host tree, generalized linear models were performed. These variables were fitted to a Poisson distribution model with a log-link function using generalized estimating equations by using the R software (R Core Team 2014). The Poisson distribution model was selected to minimize the deviance statistic. For significant terms, the individual levels were compared by using Tukey's honestly significant difference test (García-Guzmán & Dirzo 2004). Significant differences are reported where $P < 0.05$.

Leaf nutrient analyses

In the present study, the term 'leaf' denotes the primary photosynthetic organ of the plant. Leaf nutrients were measured for

B. stellatifolium, *M. angustifolia*, and *A. mearnsii*. Only mature leaf samples were taken, and leaves were considered mature when fully expanded. These were collected from five replicate trees of each tree species from each of the nine sites selected. Samples from each tree individual were removed separately with scissors and placed in plastic resealable bags with moistened paper towelling. Hands were gloved to avoid contamination of leaf material. Samples were placed in insulated containers with ice and were transferred to the Institute for Plant Production, Western Cape Department of Agriculture, Stellenbosch, South Africa, for processing.

Leaves were dried, ground in a Wiley mill (Thomas Scientific, Swedesboro, New Jersey, USA) and analysed for foliar phosphorus (P) (expressed as percentage of dry mass) by using a CE Instruments NC2100 CN analyser (CE Elantech, Incorporated, Lakewood, New Jersey, USA). Total nitrogen (expressed as percentage of dry mass) was determined by the semimicro Kjeldahl method (Ofosu-Budu *et al.* 1992) by using a Tecator Kjeltac Auto 1030 analyser (Tecator, Höganäs, Sweden). Mean leaf nutrient levels (as percentage) were compared for each plant species across sites within three different invasion treatments by using ANOVA followed by Tukey's multiple comparison tests in STATISTICA.

Effect of native tree species richness, host abundance and leaf nutrients on host damage

At each of the nine sites, two transects were established measuring 50 m in length (parallel to the river) and 5 m in width (perpendicular to the river crossing both wet and dry bank zones; for description of wet and dry bank zones, see Reinecke *et al.* 2015). In these, native tree species richness and host abundance were determined. Data from the two transects were combined per site and compared among sites with different invasion statuses as similarly done for leaf nutrient content.

To investigate possible interactions between host species abundance, native plant species richness and leaf nutrients on the extent of folivorous insect and phytopathogen damage, regression analyses were performed with the percentage of leaves damaged by the respective organism groups as the response variables and native tree species richness, host abundance and nutrient levels as predictors. Analyses were performed by using STATISTICA. Unadjusted $P < 0.05$ values are reported.

RESULTS

Differences in folivorous insect and fungal damage levels

All three tree species exhibited significant differences in levels of folivory in relation to the invasion status of the site. Levels of folivory on *B. stellatifolium* were significantly higher at invaded sites than at near-pristine sites and increased significantly from invaded sites at restored sites ($Z_{[2,42]} = 78.53$, $P < 0.001$) (Fig. 2a). For *M. angustifolia*, folivore damage showed a similar pattern,

but the difference was significant only for comparisons between restored and near-pristine sites ($Z_{[2,42]} = 72.53$, $P < 0.001$) (Fig. 2a). *Acacia mearnsii* showed a different pattern, with significantly more leaves damaged by folivorous insects at heavily invaded sites compared with near-pristine sites and restored sites ($Z_{[2,42]} = 49.42$, $P < 0.001$), which were statistically similar (Fig. 2a).

In addition to damage caused by folivorous insects, the only other organisms causing significant leaf damage to the study plants were fungi. Each host species had at least one common phytopathogenic species. *Brabejum stellatifolium* was infected by the biotrophic pathogen *Periconiella velutina* (G. Winter), which causes black mould, and the necrotrophic pathogen *Phyllosticta owaniana* (G. Winter), which causes leaf spot. *Metrosideros angustifolia* was infected by an *Alternaria* sp. that caused necrotic spots. *Acacia mearnsii* was infected by the biotrophic heteroecious pathogen *Uromycladium alpinum* (McAlpine) that caused chlorotic spots.

For *B. stellatifolium* the extent of fungal disease was significantly higher at restored sites and heavily invaded sites ($Z_{[2,42]} = 17.59$, $P < 0.001$) compared with near-pristine sites (Fig. 2b). For *M. angustifolia*, the extent of fungal disease was significantly higher in restored riparian vegetation ($Z_{[2,42]} = 31.82$, $P < 0.001$) than in near-pristine and heavily invaded sites, which did not differ significantly (Fig. 2b). For *A. mearnsii*, fungal disease was higher in heavily invaded sites ($Z_{[2,42]} = 80.37$, $P < 0.001$) than in restored and near-pristine sites, which were statistically similar (Fig. 2b).

Brabejum stellatifolium mostly incurred damage caused by folivorous insects with chewing mouthparts ($Z_{[2,42]} = 77.79$, $P < 0.001$) (Fig. 3a), while *M. angustifolia* and *A. mearnsii* were almost equally damaged by insects with chewing and those with piercing-sucking mouthparts (*M. angustifolia*: $Z_{[2,42]} = 53.75$, $P < 0.001$, and *A. mearnsii*: $Z_{[2,42]} = 25.71$, $P < 0.001$) (Fig. 3b,c). Damage levels caused by both chewing and sucking insects followed a similar pattern to those observed for the combined data presented in Figure 2a. Similarly, for *B. stellatifolium*, both fungal taxa showed the same pattern (Fig. 3d) as was found for their combined effect shown in Figure 2b.

Severity of pest and pathogen damage

Severity of leaf damage is presented in Figure 4. For all three species, the proportion of damaged leaves in each severity category varied significantly among study sites as denoted by different letters. For *B. stellatifolium* and *M. angustifolia*, damage severity caused by both folivorous insects and phytopathogenic fungi was higher at heavily invaded and restored sites than at near-pristine sites, but for both taxa, damage severity tended to be highest at the restored sites. *Acacia mearnsii* tended to be more severely damaged by both folivorous

Fig. 2. Mean percentage (+SE) of leaves damaged by either (a) folivorous insects or (b) pathogens (for *Brabejum stellatifolium*, *Metrosideros angustifolia* and *Acacia mearnsii* from riparian sites with different invasion status: near-pristine (clear), heavily invaded (dark grey) and restored (light grey). Significant differences are indicated above error bars by different lower case letters for *B. stellatifolium*, different uppercase letters for *M. angustifolia* and differences in the number of asterisks for *A. mearnsii* ($n = 15$ for each tree species in each invasion status).

insects and phytopathogenic fungi at heavily invaded sites (Fig. 4c,f).

Effect of native tree species richness, host abundance and leaf nutrients on host damage

Mean native tree species richness was significantly lower at heavily invaded sites than at near-pristine sites ($F_{[2, 6]} = 6.86$, $P = 0.03$) but returned to preinvasion levels after alien removal. Abundance of *B. stellatifolium* was highest at the near-pristine sites ($F_{[2, 6]} = 16.89$; $P = 0.003$), with no difference detected between the restored and invaded sites (Table 1). Abundance of *M. angustifolia* was highest at the near-pristine sites and restored sites, while at heavily invaded sites, abundance was significantly lower ($F_{[2, 6]} = 6.31$; $P = 0.03$) (Table 1). *Acacia mearnsii* and *M. angustifolia* showed differences in size of individuals from sites that differ in invasion status, with individuals at restored sites significantly smaller than those at near-pristine or invaded sites (Table 1). The size of *B.*

stellatifolium did not vary across sites that differ in invasion status (Table 1).

Regressions between native plant species richness and percentage of leaves damaged by phytopathogenic fungi and insects were not significant in all cases except for damage caused by phytopathogenic fungi on *A. mearnsii* (Fig. 5); in this case, damage caused by phytopathogenic fungi was negatively correlated with native plant species richness. Native host tree densities were always negatively correlated with the number of leaves damaged by phytopathogenic fungi or folivorous insects, although only significantly so in the case of damage by folivorous insects (Fig. 5). The invasive *A. mearnsii* showed the opposite, with both damage types significantly and positively correlated with host abundance (Fig. 5).

Leaf P content did not vary among sites for any plant species tested (Fig. 6). Leaf N varied significantly among sites only for *B. stellatifolium* and *A. mearnsii* (Fig. 6). For *B. stellatifolium*, levels of leaf N were significantly higher in restored sites compared with heavily invaded and

Fig. 3. Frequency of leaves damaged (a–c) by folivorous insects with chewing and sucking mouthparts and (d–f) diseases caused by four phytopathogenic fungi for *Brabejum stellatifolium*, *Metrosideros angustifolia* and *Acacia mearnsii* at sites that differed in their invasion status: near-pristine (clear), heavily invaded (dark grey) and restored riparian zones (light grey). Different letters indicate a significant difference between groups within each damage type.

near-pristine sites ($F_{[2,42]} = 68.12$, $P < 0.00001$), which did not significantly differ from each other (Fig. 6). Significantly higher levels of N for *A. mearnsii* were found in heavily invaded sites as compared with near-pristine sites ($F_{[2,42]} = 8.39$, $P = 0.0009$). Leaves of *A. mearnsii* at restored sites had intermediate N levels (Fig. 6). No significant correlations were found between leaf N content and the number of leaves damaged by phytopathogenic fungi for any tree species (Table 2). However, leaf N content was significantly correlated with the number of leaves damaged by folivorous insects for *B. stellatifolium* and *A. mearnsii*. This was in contrast to *M. angustifolia*, which showed no significant correlation between folivore damage and nitrogen content of leaves (Table 2).

DISCUSSION

In this study, both native and alien trees in riparian zones varied in their susceptibility towards pathogenic fungal colonization and folivorous insect (natural enemies) damage across sites, in accordance with differences in invasive alien plant invasion status. For the native species *B. stellatifolium*, both folivore and pathogen damages were significantly higher at invaded sites than at near-pristine sites, and these levels did not return to their original levels after restoration. In fact, folivore damage levels were even higher at restored sites than at heavily invaded sites. Similarly, damage caused by leaf phytopathogenic fungi and folivorous insects was also higher for the other native species *M. angustifolia* at

Fig. 4. Frequency distributions of percentage of leaf area damaged by (a–c) folivorous insects and (d–f) phytopathogens. The darker the bars, the more severe the damage caused by the various organisms were. Categories of leaf area damaged were as follows: 0% (light grey), 1–11% (mid grey), 12–49% (dark grey) and 50–100% (black). Significant differences across sites for each category are denoted by different letters.

Table 1. Differences in native tree species richness among studied sites that vary in their invasion status and size-classes (diameter (cm) measured at breast height) and host abundance of the three tree species considered in this study

Invasion status	Mean native tree species richness $\pm$ standard error	Plant species	Host abundance	Mean stem diameter (cm) $\pm$ standard error
Near-pristine	20 ± 0.9^a	<i>Brabejum stellatifolium</i>	25 ± 1.7^a	36 ± 1.1^a
		<i>Metrosideros angustifolia</i>	26 ± 1.0^A	31 ± 2.4^B
		<i>Acacia mearnsii</i>	$6 \pm 0.3^*$	$30 \pm 1.3^*$
Heavily invaded	7 ± 2.3^b	<i>B. stellatifolium</i>	11 ± 2.1^b	29 ± 0.95^a
		<i>M. angustifolia</i>	8 ± 2.0^B	19 ± 1.1^A
		<i>A. mearnsii</i>	$63 \pm 30.9^{**}$	$36 \pm 2.4^*$
Restored	17 ± 0.7^a	<i>B. stellatifolium</i>	15 ± 1.3^b	32 ± 1.3^a
		<i>M. angustifolia</i>	17 ± 3.3^{AC}	22 ± 2.2^A
		<i>A. mearnsii</i>	$6 \pm 0.3^*$	$13 \pm 0.9^{**}$

Significant differences in native tree species richness, host abundance and size-classes are indicated by different lower case letters for *B. stellatifolium*, different uppercase letters for *M. angustifolia* and differences in the number of asterisks for *A. mearnsii*.

Fig. 5. The effects of host abundance on the percentage of leaves damaged by (a,c,e) folivorous insects or (b,d,f) phytopathogens for three tree species (*Brabejum stellatifolium*, *Metrosideros angustifolia* and *Acacia mearnsii*) at sites that differ in their invasion status: near-pristine (clear square), heavily invaded (dark grey square) and restored (light-grey square) riparian zones. Reported P -values are for partial regression coefficients; reported r^2 is for each overall regression. All trends were linear.

restored sites than it was at heavily invaded and near-pristine sites. The pattern was different for the invasive *A. mearnsii*, where damage caused by both folivorous insects and phytopathogenic fungi increased at heavily invaded sites but returned to near-pristine site levels at restored sites. The high levels of folivory and fungal diseases on *A. mearnsii* could not be credited to biological control agents because the two known biocontrol agents (*Melanterius maculatus* and *Dasineura rubiformis*) do not damage the vegetative parts of *A. mearnsii* but rather cause reductions in seed availability (Impson *et al.* 2011). The lower levels of folivory and fungal disease in near-pristine sites concur with the literature, which shows that higher heterogeneity in plant species composition and structure of near-pristine sites possibly explains the observed lower levels of damage by natural enemies (Bach 1980; Agrawal *et al.* 2006). The higher levels of damage by folivorous insects and phytopathogenic fungi on native plants in restored areas are intriguing, particularly because this study was undertaken 7 years after restoration efforts.

After removal of invasive alien plants, host abundance of *B. stellatifolium* and *M. angustifolia* (two native trees) was similar to that in near-pristine sites. However, abundance of these species was low at heavily invaded sites. Abundance of *A. mearnsii* was greater in heavily invaded sites than in restored and near-pristine sites. Apart from

decreased host abundance of the two native trees in heavily invaded sites, native plant species richness was also lower than that in near-pristine sites. Results therefore support previous observations that alien plant species have a negative impact on native species richness and abundance by replacing native plant species in the communities they invade (French *et al.* 2008; Vilà *et al.* 2011). Most invasive Australian *Acacia* spp. in south Africa are transformers, posing serious threat to the structure and function of pristine fynbos areas by elevating soil nitrogen levels and converting them into species-poor ecosystems (Richardson *et al.* 2000). Despite these severe impacts, native plant species richness can return to its original level once the invasive alien plants are removed (Holmes & Cowling 1997; Blanchard & Holmes 2008; Marchante *et al.* 2011 and this study). The recovery of native plant species in riparian areas referred to in these studies suggests that in each case, no 'threshold of irreversibility' (as defined by Aronson *et al.* 1993) had been reached (see Holmes & Cowling 1997), thereby allowing regeneration after clearing.

There is a debate over whether or not the levels of leaf damage by folivorous insects and phytopathogenic fungi depend on native plant species richness, with homogeneous tree stands being more damage prone than heterogeneous stands (Mitchell *et al.* 2002; Pfisterer *et al.* 2003; Haas *et al.* 2011). Consequently, one would expect to

Fig. 6. Mean + SE of (a) foliar nitrogen content and (b) phosphorus content measured as percentage of dry mass of *Brabejum stellatifolium*, *Metrosideros angustifolia* and *Acacia mearnsii* from riparian sites with different invasion status: near-pristine (clear), heavily invaded (dark grey) and restored (light grey). Significant differences are indicated above bars by different lower case letters for each tree species.

Table 2. Correlations for foliar damage caused by phytopathogenic fungi and folivorous insects as response variables *versus* native tree species richness and leaf nitrogen as the predictors ($n = 9$)

Plant species	Number of leaves damaged by phytopathogenic fungi		Number of leaves damaged by folivorous insects	
	Tree species richness	Leaf nitrogen	Tree species richness	Leaf nitrogen
<i>Brabejum stellatifolium</i>	0.21n.s.	0.21n.s.	0.02n.s.	0.51*
<i>Metrosideros angustifolia</i>	0.00n.s.	0.03n.s.	0.13n.s.	-0.11n.s.
<i>Acacia mearnsii</i>	0.59*	0.21n.s.	0.25n.s.	0.52*

* $P \leq 0.001$; n.s., not significant, $P > 0.05$. Data presented are for the fitted intercept, the coefficient of determination (r^2).

see decreasing levels of folivory and fungal disease with increasing native plant species richness. However, results from this study suggest that activities of folivores and phytopathogenic fungi in diverse riparian habitats (near-pristine and restored) and simple habitats (heavily invaded) are independent of plant species richness, a pattern observed in other ecosystems (e.g. Scherber

et al. 2010). Bach (1980) suggested that folivore activity is linked to host abundance/density rather than plant species richness. An area that has a high abundance/density of a preferred host favours folivore survival more so than a diverse area where access to the preferred host is more limited, a phenomenon known as the 'resource concentration' hypothesis (Grez & González 1995).

Apart from host abundance, Mitchell *et al.* (2002) highlighted the importance of understorey richness in maintaining and controlling natural enemies of plants. We did not include understorey plant species in our analyses, which might have influenced the results obtained for restored and near-pristine sites. Therefore, even though tree species richness was similar, understorey plant richness could have been less diverse in restored sites, leading to a decrease in natural enemies and increased levels of folivory. Future studies should therefore include understorey plant richness as an explanatory variable for altered herbivore pressure.

Several studies have tested relationships between levels of folivory, fungal diseases and the abundance of hosts, with some studies showing positive (Mabry & Wayne 1997), some negative (Underwood & Halpern 2012; Halpern *et al.* 2014) and some neutral (Rhainds & English-Loeb 2003) effects. In this study, increased host abundance was negatively correlated with native tree folivory and positively correlated with the levels of both folivory and fungal diseases for *A. mearnsii*. The negative relationship between folivory and host abundance for the two native trees could be explained by dilution of folivore effects with increased abundance (e.g. see Halpern *et al.* 2014). Alternatively, at low abundance, when host plants are thinly dispersed, folivores may show fidelity to a host, as finding another host increases both energy use and exposure to predaceous arthropods (Schuldt *et al.* 2010).

Levels of damage by folivores and pathogens on *A. mearnsii* seemed to follow a more classic pattern as is well known in agricultural monotypic stands. *Acacia mearnsii* forms thick near-monocultures that make it easy for insects to find hosts (if preadapted) and for spores of pathogens to land on susceptible hosts (e.g. Bell *et al.* 2006). It is also possible that predator communities may have been disrupted. There is some evidence that in their introduced range, alien plants may assemble communities of folivores as abundant and rich as those of native hosts (Frenzel & Brandl 2003), especially if the invasive alien plants is closely related to several native trees (Procheş *et al.* 2008). The Fabaceae are well represented in South Africa (18% of the 198 listed trees species in South Africa belong to Fabaceae; Coates-Palgrave 2002). Hence, it is possible that a rich regional pool of preadapted folivores is available to counteract chemical defences of *A. mearnsii* (e.g. Litt *et al.* 2014).

Apart from the influences of host abundance and native plant species richness, the observed higher levels of folivory and fungal diseases may be related to changes in leaf chemical components due to high rates of nutrient inputs to the ecosystems through the presence of nitrogen-fixing invasive alien plants (e.g. *A. mearnsii* is a putative nitrogen-fixing legume; Tye & Drake 2012), which are known to elevate soil nutrient levels (van der Waal 2009). A study by Naudé (2012), at some of the

same riparian sites used in this study, showed that a legacy effect remained after clearing invasive alien plants in that levels of available nitrogen in soil remained high. This implies that plants in restored sites may be more vigorous than in near-pristine sites because of abundant available resources (such as seen for the native trees in this study). This, in turn, may make these plants more attractive to folivorous insects and/or phytopathogenic fungi (Moran & Scheidler 2002), thereby increasing their abundance (and causing an increase in damage) in riparian systems after alien clearing.

Recorded levels of foliar nitrogen of the host trees fell within the range of values reported for other Mediterranean trees (Fife *et al.* 2008) and for other evergreen trees such as Australian *Acacia* spp. (Peeters 2002b). Likewise, leaf phosphorus levels were within the lower and upper ranges of reported values for these hosts (Fife *et al.* 2008). Foliar nutrient content data confirmed that *B. stellatifolium* had greater levels of foliar nitrogen at restored sites, as did *A. mearnsii*. Phosphorus levels did not vary significantly for any plant species tested across all invasion statuses. Of the covariates tested, leaf nitrogen was the only factor that was positively correlated with folivores and leaf chewers for *B. stellatifolium* and with folivores for *A. mearnsii*. This result suggests that nitrogen content is an important driver of arthropod population dynamics for *B. stellatifolium* and *A. mearnsii*.

In conclusion, the results of this study suggest that, although plant communities often recover after invasive alien removal, native trees continue to be damaged by folivorous insects and pathogenic fungi, resulting in damage levels in excess of those seen at invaded sites. This implies that restoration efforts of invaded riparian zones do not remove the legacy effect of the invasion, with many unexpected and lasting ecological consequences. Of the explanatory factors assessed, host abundance (resource concentration) and nitrogen availability (resource quality) seemed to be the most influential on damage caused by arthropods. Neither of these two factors explained the similar observed patterns for damage caused by phytopathogenic fungi. Although these correlative analyses cannot ascertain the exact mechanism by which host plant abundance affected folivory, it is highly possible that the interplay between folivore densities, predaceous arthropod densities and/or plant nutritional status may be important (see Forkner & Hunter 2000; Moran & Scheidler 2002). Investigating the effect of these factors on pest and pathogen populations under controlled conditions is therefore a necessary next step for future research.

ACKNOWLEDGEMENTS

The authors are grateful to Marno Fourie and Oliver Kambaj for field assistance. They thank private landowners and CapeNature for access to field sites. This

study was funded by the DST/NRF Centre of Excellence in Tree Health Biotechnology. The authors are also thankful to Dr Kenneth Oberlander for assistance with molecular techniques for identification of phytopathogenic fungi isolates in this study.

REFERENCES

- Agrawal A., Lau J. & Hamback P. (2006) Community heterogeneity and the evolution of interactions between plants and insect herbivores. *Q. Rev. Biol.* **81**, 349–76.
- Agrios G. N. (2005) *Plant Pathology*. Elsevier Academic Press, London, United Kingdom.
- Aronson J., Floret C., Le Floch E., Ovalle C. & Pontanier R. (1993) Restoration and rehabilitation of degraded ecosystems in arid and semi-arid lands. *I. A view from the South. Restoration Ecol.* **1**, 8–17.
- Arzanlou M., Groenewald J. Z., Gams W., Braun U., Shin H. D. & Crous P. W. (2007) Phylogenetic and morphotaxonomic revision of *Ramichloridium* and allied genera. *Stud. Mycol.* **58**, 57–93.
- Bach C. E. (1980) Effects of plant density and diversity on the population dynamics of a specialist herbivore, the striped cucumber beetle, *Acalymma vittata* (Fab). *Ecology* **61**, 1515–30.
- Bell T., Freckleton R. P. & Lewis O. T. (2006) Plant pathogens drive density-dependent seedling mortality in a tropical tree. *Ecol. Lett.* **9**, 569–74.
- Benítez-Malvido J., García-Guzmán G. & Kossmann-Ferraz I. D. (1999) Leaf-fungal incidence and herbivory on tree seedlings in tropical rainforest fragments: An experimental study. *Biol. Conserv.* **91**, 143–50.
- Blanchard R. & Holmes P. M. (2008) Riparian vegetation recovery after invasive alien tree clearance in the Fynbos Biome. *S. Afr. J. Bot.* **74**, 421–31.
- Brooks T. M., daFonseca G. S. B. & Rodrigues A. S. L. (2004) Protected areas and species. *Conserv. Biol.* **18**, 616–8.
- Coates-Palgrave K. (2002) *Trees of Southern Africa*. Struik, Cape Town, South Africa.
- Cook W. M. & Holt R. D. (2006) Influence of multiple factors on insect colonization of heterogeneous landscapes: A review and case study with periodical cicadas (Homoptera: Cicadidae). *Ecol. popul. biol.* **99**, 809–20.
- Covington W., Fule P., Moore M., et al. (1997) Restoring ecosystem health in ponderosa pine forests of the southwest. *J. For.* **95**, 23–9.
- Crous C. J., Jacobs S. M. & Esler K. J. (2012) Drought-tolerance of an invasive alien tree. *Acacia mearnsii* and two native competitors in Fynbos riparian ecotones. *Biol. Invasions* **14**, 619–31.
- Deacon H. J., Jury M. R. & Ellis F. (1992) Selective time and regime. In: *Fynbos: Nutrients, Fire and Diversity* (ed R. M. Cowling) pp. 6–23. Oxford University Press, London.
- Doidge E. M. & Bottomley A. M. (1931) A revised list of diseases occurring in South Africa. *Bot. Surv. S. Afr.* **11**, 1–78.
- Dye P. & Jarmain C. (2004) Water use by black wattle (*Acacia mearnsii*): Implications for the link between removal of invading trees and catchment streamflow response. *S. Afr. J. Sci.* **100**, 40–5.
- Eichhorn M. P., Nilus R., Compton S. G., Hartley S. E. & Burslem D. (2010) Herbivory of tropical rain forest tree seedlings correlates with future mortality. *Ecology* **91**, 1092–101.
- Fife D. N., Nambiar E. K. S. & Saur E. (2008) Retranslocation of foliar nutrients in evergreen tree species planted in a Mediterranean environment. *Tree Physiol.* **28**, 187–96.
- French K., Ens E., Gosper C. R., et al. (2008) Management implications of recent research into the effect of bitou bush invasion. *Plant Prot. Q.* **23**, 4–28.
- Frenzel M. & Brandl R. (2003) Diversity and abundance patterns of phytophagous insect communities on alien and native host plants in the Brassicaceae. *Ecography* **26**, 723–30.
- Folgarait P., Marquis R., Ingvarsson P., Braker H. & Arguedas M. (1995) Patterns of attack by insect herbivores and a fungus on saplings in a tropical tree plantation. *Environ. Entomol.* **24**, 1487–94.
- Forkner R. E. & Hunter M. D. (2000) What goes up must come down? Nutrient addition and predation pressure on oak herbivores. *Ecology* **81**, 1588–600.
- Galatowitsch S. & Richardson D. M. (2005) Riparian scrub recovery after clearing of invasive alien trees in headwater streams of the Western Cape, South Africa. *Biol. Conserv.* **12**, 509–21.
- García-Guzmán G. & Dirzo R. (2004) Incidence of leaf pathogens in the canopy of a Mexican tropical wet forest. *Plant Ecol.* **172**, 31–50.
- Gardes M. & Bruns T. D. (1993) ITS primers with enhanced specificity for basidiomycetes application to the identification of mycorrhizae and rusts. *Mol. Ecol.* **2**, 113–8.
- Garrett K. A. & Mundt C. C. (2000) Effects of planting density and the composition of wheat cultivar mixtures on stripe rust: An analysis taking into account limits to the replication of controls. *Phytopathol.* **90**, 1313–21.
- Gilbert G. S. (2002) Evolutionary ecology of plant diseases in natural ecosystems. *Annu. Rev. Phytopathol.* **40**, 13–43.
- Goldblatt P. & Manning J. C. (2000) Cape plants: A conspectus of the Cape flora of South Africa. *Strelitzia* **7**, 1–743.
- Greze A. A. & Gonzalez R. H. (1995) Resource concentration hypothesis-effect of host-plant patch size on density of herbivorous insects. *Oecologia* **103**, 471–74.
- Haas S. E., Hooten M. B., Rizzo D. M. & Meentemeyer R. K. (2011) Forest species diversity reduces disease risk in a generalist plant pathogen invasion. *Ecol. Lett.* **14**, 1108–16.
- Halpern S. I., Bednar D., Chisholm A. & Underwood N. (2014) Plant-mediated effects of host plant density on a specialist herbivore of *Solanum carolinense*. *Ecol. Entomol.* **39**, 217–25.
- Holmes P. M. & Cowling R. M. (1997) The effects of invasion by *Acacia saligna* on the guild structure and regeneration capabilities of South African fynbos shrublands. *J. Appl. Ecol.* **34**, 317–32.
- Impson F. A. C., Kleinjan C. A., Hoffmann J. H., Post J. A. & Wood A. R. (2011) Biological control of Australian *Acacia* species and *Paraserianthes lophantha* (Willd.) Nielsen (Mimosaceae), in South Africa. *Afr. Entomol.* **19**, 186–207.
- Johnson P. T. J., Hartson R. B., Larson D. J. & Sutherland D. R. (2008) Diversity and disease: Community structure drives parasite transmission and host fitness. *Ecol. Lett.* **11**, 1017–26.
- Litt A. R., Cord E. E., Fulbright T. E. & Schuster G. L. (2014) Effects of invasive plants on arthropods. *Conserv. Biol.* **28**, 1532–49.
- Mabry C. M. & Wayne P. W. (1997) Defoliation of the annual herb *Abutilon theophrasti*: Mechanisms underlying reproductive compensation. *Oecologia* **111**, 225–32.
- Marchante H., Freitas H. & Hoffmann J. H. (2011) Post-clearing recovery of coastal dunes invaded by *Acacia longifolia*: Is duration of invasion relevant for management success? *J. Appl. Ecol.* **9**, 1780–90.
- Mitchell C. E., Tilman D. & Groth J. V. (2002) Effects of grassland plant species diversity, abundance and composition on foliar fungal disease. *Ecology* **83**, 1713–26.
- Moran M. D. & Scheidler A. R. (2002) Effects of nutrients and predators on an old-field food chain: Interactions of top-down and bottom-up processes. *Oikos* **98**, 116–24.

- Morris M. J., Wingfield M. J. & Walker J. (1988) First record of a rust on *Acacia mearnsii* in Southern Africa. *Trans. Br. Mycol. Soc.* **90**, 324–7.
- Mucina L. & Rutherford M. C. (2006) *The Vegetation of South Africa, Lesotho and Swaziland*. South African National Biodiversity Institute, Pretoria.
- Musvuugwa T., Wilhelm de Beer Z., Duong T. A., Dreyer L. L., Oberlander K. C. & Roets F. (2015) New species of Ophiostomatales from Scolytinae and Platypodinae beetles in Cape Floristic Region including the discovery of the sexual state of *Raffaelea*. *Anton. Leeuw.* **108**, 933–50.
- Naudé M. (2012) *Fynbos riparian biogeochemistry and invasive Australian Acacias*. Stellenbosch University, South Africa, MSc. thesis.
- Ofose-Budu K. G., Ogata S. & Fujita K. (1992) Temperature effects on root nodule activity and nitrogen release in some sub-tropical and temperate legumes. *Soil Sci. Plant Nutr.* **38**, 717–26.
- Peeters P. (2002a) Correlations between leaf structural traits and the densities of herbivorous insect guilds. *Biol. J. Linn. Soc.* **77**, 43–65.
- Peeters P. J. (2002b) Correlations between leaf constituent levels and the densities of herbivorous insect guilds in an Australian forest. *Austral Ecol.* **27**, 658–71.
- Pfisterer A., Diemer M. & Schmid B. (2003) Dietary shift and lowered biomass gain of a generalist herbivore in species-poor experimental plant communities. *Oecologia* **135**, 234–41.
- Procheş Ş., Wilson J. R. U., Richardson D. M. & Chown S. L. (2008) Herbivores, but not other insects, are scarce on alien plants. *Austral Ecol.* **33**, 691–700.
- R Core Team (2014) *R: A language and environment for statistical computing (3.1.0 ed.)*. R Foundation for Statistical Computing, Vienna, Austria.
- Reinecke M. K., Brown C. A., Esler K. J., King J. M., Kleynhans M. T. & Kidd M. (2015) Links between lateral vegetation zones and river flow. *Wetlands* **35**, 473–86.
- Rhainds M. & English-Loeb G. (2003) Testing the resource concentration hypothesis with tarnished plant bug on strawberry: density of hosts and patch size influence the interaction between abundance of nymphs and incidence of damage. *Ecol. Entomol.* **28**, 348–58.
- Richardson D. M., Pyšek P., Rejmánek M., Barbour M. G., Panetta F. D. & West C. J. (2000) Naturalization and invasion of alien plants: Concepts and definitions. *Divers. Distrib.* **6**, 93–107.
- Riihimäki J., Kaitaniemi P., Koricheva J. & Vehviläinen H. (2005) Testing the enemies hypothesis in forest stands: The important role of tree species composition. *Oecologia* **142**, 90–7.
- Root R. B. (1973) Organization of a plant–arthropod association in simple and diverse habitats: The fauna of collards (*Brassica oleracea*). *Ecol. Monogr.* **43**, 95–124.
- Scherber C., Specht J., Köhler G., Mitschunas N. & Weisser W. W. (2010) Functional identity versus species richness: Herbivory resistance in plant communities. *Oecologia* **163**, 707–17.
- Schuldt A., Baruffol M., Böhnke M., et al. (2010) Tree diversity promotes insect herbivory in subtropical forests of South-East China. *Ecology* **98**, 917–26.
- Torchin M. E. & Mitchell C. E. (2004) Parasites, pathogens, and invasions by plants and animals. *Front. Ecol. Environ.* **2**, 183–90.
- Tye D. R. & Drake D. C. (2012) An exotic Australian *Acacia* fixes more N than a coexisting indigenous *Acacia* in a South African riparian zone. *Plant Ecol.* **213**, 251–7.
- Underwood N. & Halpern S. (2012) Insect herbivores, density dependence, and the performance of the perennial herb *Solanum carolinense*. *Ecology* **93**, 1026–35.
- Vilà M., Espinar J. L., Hejda M., et al. (2011) Ecological impacts of invasive alien plants: A meta-analysis of their effects on species, communities and ecosystems. *Ecol. Lett.* **14**, 702–8.
- van der Waal B. W. (2009) *The influence of Acacia mearnsii* invasion on soil properties in the Kouga Mountains, Eastern Cape
- . Rhodes University, Grahamstown, South Africa. MSc thesis.
- van Hezewijk B. H., De Clerck-Floate R. A. & Moyer J. R. (2008) Effect of nitrogen on the preference and performance of a biological control agent for an invasive plant. *Biol. Control* **46**, 332–40.
- van Wilgen B. W., Forsyth G. C., Le Maitre D. C., et al. (2012) An assessment of the effectiveness of a large, national-scale invasive alien plant control strategy in South Africa. *Biol. Conserv.* **148**, 28–38.
- White T. J., Bruns T. D., Lee S. & Taylor J. W. (1990) Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: *PCR Protocols: A Guide to Methods and Applications* (eds M. A. Innis, D. H. Gelfand & J. J. Sninsky) pp. 315–322. New York Academic Press, New York.
- White E. M., Wilson J. C. & Clarke A. R. (2006) Biotic indirect effects: a neglected concept in invasion biology. *Divers. Distrib.* **12**, 443–55.